

# An introduction to asymptotic enumeration

Part 1. Why are asymptotic results interesting?

Part 2. Asymptotics of sequences  $a(n)$

Part 3. Limit laws

## The study of additional statistics

$$\begin{array}{ll} \text{size} : \mathcal{A} & \rightarrow \mathbb{N} \\ a & \mapsto |a| \end{array} \qquad \begin{array}{ll} s : \mathcal{A} & \rightarrow \mathbb{Z} \\ a & \mapsto s(a) \end{array}$$


When objects are taken uniformly in  $\mathcal{A}_n$ , the statistic  $s$  becomes a **random variable**  $S_n$ :

$$\mathbb{P}(S_n = k) = \frac{a(n, k)}{a(n)}$$

where  $a(n, k)$  is the number of objects of size  $n$  for which the additional statistic  $s$  equals  $k$ .

## Examples of additional statistics


- the final height of a 1D walk with  $n$  steps,
- the maximal height of such a walk,
- the width/height of a directed animal with  $n$  cells,
- the area enclosed by a SAP of perimeter  $2n$ ,
- the area enclosed by a staircase polygon of perimeter  $2n$ ...



## Convergence in law (or: in distribution)

**Definition.** The random variables  $S_n$  converge in law to the random variable  $S$ , having distribution function  $F(x)$ , if for all  $x$  [where  $F$  is continuous],


$$\mathbb{P}(S_n \leq x) = F_n(x) \rightarrow F(x) = \mathbb{P}(S \leq x).$$



## A discrete limit law: contacts in a Dyck path

Convergence of the distribution function of  $S_n$ :

$$\mathbb{P}(S_n \leq \ell + 1) \rightarrow \sum_{k=1}^{\ell} \frac{k}{2^{k+1}}$$




## A continuous limit law: The final height of a random walk

As  $\mathbb{E}(S_n^2) = n$ , we consider the random variable  $S_n/\sqrt{n}$ .  
Its distribution function satisfies

$$\mathbb{P}\left(\frac{S_n}{\sqrt{n}} \leq x\right) \rightarrow \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt$$

$\Rightarrow$  Convergence to a Gaussian law



# How to prove convergence in law

- From the distribution function
- Pumping moments
- Discrete limit laws: pointwise convergence of the probability generating function
- Continuous limit laws: pointwise convergence of the Laplace or Fourier transform

# 1. Convergence of the moments and convergence in law

**Theorem.** For  $S_n$  to converge in law to  $S$ , it suffices that

- all moments converge :  $\mathbb{E}(S_n^j) \rightarrow \mathbb{E}(S^j)$  for all  $j$
- the Laplace transform

$$\mathbb{E}(e^{xS}) = \sum_{j \geq 0} \mathbb{E}(S^j) \frac{x^j}{j!}$$

has a positive radius of cv.



## Where do we find the moments of the statistic $s$ ?

Bivariate generating function:

$$A(t, u) = \sum_{a \in \mathcal{A}} t^{|a|} u^{s(a)} = \sum_n t^n \left( \sum_k a(n, k) u^k \right)$$

Specialization  $u = 1$ :

$$A(t, 1) = \sum_{a \in \mathcal{A}} t^{|a|} = \sum_n a(n) t^n$$

• Average value of  $S_n$ :

$$u \frac{\partial}{\partial u} A(t, u) = \sum_n t^n \left( \sum_k k a(n, k) u^k \right) \Rightarrow u \frac{\partial}{\partial u} A(t, u) \Big|_{u=1} = \sum_n t^n a(n) \mathbb{E}(S_n)$$

## Where do we find the moments of the statistic $s$ ?

Bivariate generating function:

$$A(t, u) = \sum_{a \in \mathcal{A}} t^{|a|} u^{s(a)} = \sum_{n, k} a(n, k) t^n u^k$$

Specialization  $u = 1$ :

$$A(t, 1) = \sum_{a \in \mathcal{A}} t^{|a|} = \sum_n a(n) t^n$$

• Average value of  $S_n$ :

$$u \frac{\partial}{\partial u} A(t, u) = \sum_n t^n \left( \sum_k k a(n, k) u^k \right) \Rightarrow u \frac{\partial}{\partial u} A(t, u) \Big|_{u=1} = \sum_n t^n a(n) \mathbb{E}(S_n)$$

• Further moments:

$$\left( u \frac{\partial}{\partial u} \right)^i A(t, u) = \sum_n t^n \left( \sum_k k^i a(n, k) u^k \right) \Rightarrow \left( u \frac{\partial}{\partial u} \right)^i A(t, u) \Big|_{u=1} = \sum_n t^n a(n) \mathbb{E}(S_n^i)$$

## Where do we find the moments of the statistic $s$ ?

Bivariate generating function:

$$A(t, u) = \sum_{a \in \mathcal{A}} t^{|a|} u^{s(a)} = \sum_{n, k} a(n, k) t^n u^k$$

Specialization  $u = 1$ :

$$A(t, 1) = \sum_{a \in \mathcal{A}} t^{|a|} = \sum_n a(n) t^n$$

- Average value of  $S_n$ :

$$u \frac{\partial}{\partial u} A(t, u) = \sum_n t^n \left( \sum_k k a(n, k) u^k \right) \Rightarrow u \frac{\partial}{\partial u} A(t, u) \Big|_{u=1} = \sum_n t^n a(n) \mathbb{E}(S_n)$$

- Further moments:

$$\left( u \frac{\partial}{\partial u} \right)^i A(t, u) = \sum_n t^n \left( \sum_k k^i a(n, k) u^k \right) \Rightarrow \left( u \frac{\partial}{\partial u} \right)^i A(t, u) \Big|_{u=1} = \sum_n t^n a(n) \mathbb{E}(S_n^i)$$

- The exponential moments:

$$\frac{\partial^i A}{\partial u^i}(t, 1) = \sum_n t^n a(n) \mathbb{E}(S_n(S_n - 1) \cdots (S_n - i + 1))$$

## Contacts in bilateral Dyck paths

- GF of paths counted by their half-length ( $t$ ) and number of visits at 0 ( $u$ ):

$$A(t, u) = \frac{u}{1 - 2uP(t)} \quad \text{with} \quad P(t) = \frac{1 - \sqrt{1 - 4t}}{2}$$

$$\Rightarrow A(t, 1) = \frac{1}{\sqrt{1 - 4t}} = \sum_{n \geq 0} \binom{2n}{n} t^n = \sum_n a(n) t^n$$

## Contacts in bilateral Dyck paths

- GF of paths counted by their half-length ( $t$ ) and number of visits at 0 ( $u$ ):

$$A(t, u) = \frac{u}{1 - 2uP(t)} \quad \text{with} \quad P(t) = \frac{1 - \sqrt{1 - 4t}}{2}$$

$$\Rightarrow A(t, 1) = \frac{1}{\sqrt{1 - 4t}} = \sum_{n \geq 0} \binom{2n}{n} t^n = \sum_n a(n) t^n$$

- Take derivatives w.r.t.  $u$

$$\frac{\partial^i A}{\partial u^i}(t, 1) = \frac{i! (2P(t))^i}{(1 - 4t)^{(i+1)/2}} = \sum_n t^n a(n) \mathbb{E}(S_n(S_n - 1) \dots (S_n - i + 1))$$

## Contacts in bilateral Dyck paths

- GF of paths counted by their half-length ( $t$ ) and number of visits at 0 ( $u$ ):

$$A(t, u) = \frac{u}{1 - 2uP(t)} \quad \text{with} \quad P(t) = \frac{1 - \sqrt{1 - 4t}}{2}$$

$$\Rightarrow A(t, 1) = \frac{1}{\sqrt{1 - 4t}} = \sum_{n \geq 0} \binom{2n}{n} t^n = \sum_n a(n) t^n$$

- Take derivatives w.r.t.  $u$

$$\frac{\partial^i A}{\partial u^i}(t, 1) = \frac{i! (2P(t))^i}{(1 - 4t)^{(i+1)/2}} = \sum_n t^n a(n) \mathbb{E}(S_n(S_n - 1) \dots (S_n - i + 1))$$

- Singularity analysis:

$$\mathbb{E}(S_n(S_n - 1) \dots (S_n - i + 1)) \sim \frac{i! \Gamma(1/2)}{\Gamma((i + 1)/2)} n^{i/2}$$

$$\Rightarrow \mathbb{E} \left( \left( \frac{S_n}{\sqrt{n}} \right)^i \right) \sim \frac{i! \Gamma(1/2)}{\Gamma((i + 1)/2)}$$

These are the moments of the **Rayleigh law** of density  $x/2 \exp(-x^2/4)$  for  $x > 0$   
 $\Rightarrow$  Convergence in law to a Rayleigh distribution

## The area of staircase polygons

- GF of staircase polygons by half-perimeter ( $t$ ) and area ( $u$ ):

$$P(t, u) = t^2u + 2tuP(t, u) + P(t, u)P(tu, u)$$

- Repeated differentiations of the equation

$\Rightarrow \frac{\partial^i P}{\partial u^i}(t, 1)$  is always algebraic and

$$\frac{\partial^j P}{\partial u^j}(t, 1) = \sum_n t^n a(n) \mathbb{E}(S_n(S_n - 1) \dots (S_n - j + 1)) \sim \frac{c_j}{(1 - 4t)^{(3j-1)/2}}$$

with

$$c_0 = -1/2, \quad c_j = \frac{j(3j - 4)}{8}c_{j-1} + \sum_{i=1}^{j-1} \binom{j}{i} c_i c_{j-i}$$

- Singularity analysis

$$\mathbb{E} \left( \left( \frac{S_n}{n^{3/2}} \right)^j \right) \sim \frac{c_j}{c_0} \frac{\Gamma(-1/2)}{\Gamma((3j - 1)/2)}$$

Moments of the Airy distribution  $\Rightarrow$  Convergence in law

## 2. Continuity theorem for discrete limit laws

**Theorem.** For  $S_n$  to converge in law to  $S$ , it suffices that the probability generating function of  $S_n$  converges pointwise to that of  $S$ :

$$G_n(u) := \sum_k \mathbb{P}(S_n = k)u^k \rightarrow G(u) := \sum_k \mathbb{P}(S = k)u^k$$

for  $u$  in a set  $\Omega$  having an accumulation point of modulus  $< 1$ .

**Example:**  $\Omega = [0, 1/2]$



## Where do we find the probability g.f. of the statistic $s$ ?

- Probability GF:

$$G_n(u) := \sum_k \mathbb{P}(S_n = k)u^k = \frac{1}{a(n)} \sum_k a(n, k)u^k$$

- Bivariate generating function:

$$A(t, u) = \sum_n t^n \left( \sum_k a(n, k)u^k \right) = \sum_n t^n a(n)G_n(u)$$

Specialization  $u = 1$ :

$$A(t, 1) = \sum_{a \in \mathcal{A}} t^{|a|} = \sum_n a(n)t^n$$

⇒ Estimate the coefficient of  $t^n$  in  $A(t, u)$ , where  $u$  is a fixed real or complex number

⇒ Recycle the methods used for the analysis of  $A(t, 1)$ !

## Contacts in Dyck paths

- Let  $A(t, u)$  count Dyck paths by the half-length ( $t$ ) and contacts ( $u$ ):

$$A(t, u) = \frac{u}{1 - u(1 - \sqrt{1 - 4t})/2}$$

- Fix  $u$  in  $[0, 1]$ . Then, as  $t \rightarrow 1/4$ :

$$A(t, u) = \frac{2}{2 - u} \left( 1 - \frac{u}{2 - u} \sqrt{1 - 4t} \right) + O(1 - 4t)$$

## Contacts in Dyck paths

- Let  $A(t, u)$  count Dyck paths by the half-length ( $t$ ) and contacts ( $u$ ):

$$A(t, u) = \frac{u}{1 - u(1 - \sqrt{1 - 4t})/2}$$

- Fix  $u$  in  $[0, 1]$ . Then, as  $t \rightarrow 1/4$ :

$$A(t, u) = \frac{2}{2 - u} \left( 1 - \frac{u}{2 - u} \sqrt{1 - 4t} \right) + O(1 - 4t)$$

- Singularity analysis:

$$[t^n]A(t, u) = a(n)G_n(u) \sim -\frac{2u}{(2 - u)^2} \frac{4^n}{\Gamma(-1/2)} n^{-3/2}$$

$$G_n(u) \sim \frac{u}{(2 - u)^2} = \sum_{k \geq 1} \frac{k}{2^{k+1}} u^k = G(u)$$

Convergence in law to the distribution given by

$$\mathbb{P}(S = k) = \frac{k}{2^{k+1}}$$

## The degree of the root in a Cayley tree

- Exponential GF of labelled rooted trees by vertices ( $t$ ) and degree of the root ( $u$ )

$$A(t, u) = t \exp(uT(t)) \quad \text{with} \quad T(t) = t \exp(T(t))$$

- For  $u > 0$ ,  $A(t, u)$  has a unique dominant singularity at  $t = 1/e$ , in the vicinity of which

$$A(t, u) = e^{u-1} - ue^{u-1} \sqrt{2} \sqrt{1-te} (1 + o(1))$$

- Singularity analysis:

$$[t^n] A(t, u) = \frac{a(n)}{n!} G_n(u) \sim -ue^{u-1} \sqrt{2} \frac{e^n}{\Gamma(-1/2)} n^{-3/2}$$

$$G_n(u) \sim ue^{u-1} = \sum_{k \geq 1} \frac{u^k}{e(k-1)!} = G(u)$$

Distribution function of a Poisson random variable

### 3. Continuity theorem for continuous limit laws

**Theorem 1.** For the sequence  $S_n$  to converge in law to  $S$ , it suffices that the **Fourier transform** (or characteristic function) converges pointwise

$$\mathbb{E}(e^{ixS_n}) \rightarrow \mathbb{E}(e^{ixS})$$

for all  $x$  in  $\mathbb{R}$ .

**Theorem 2.** For the sequence  $S_n$  to converge in law to  $S$ , it suffices that the **Laplace transform** converges pointwise

$$\mathbb{E}(e^{xS_n}) \rightarrow \mathbb{E}(e^{xS})$$

for all  $x$  in a real neighborhood of 0.

## Where do we find the Laplace transform of the statistic $s$ ?

Bivariate generating function:

$$A(t, u) = \sum_n t^n \left( \sum_k a(n, k) u^k \right) = \sum_n t^n a(n) \mathbb{E} \left( u^{S_n} \right)$$
$$\Rightarrow a(n) \mathbb{E} \left( e^{x S_n} \right) = [t^n] A(t, e^x)$$

$\Rightarrow$  Estimate the coefficient of  $t^n$  in  $A(t, u)$ , for  $u$  close to 1

# The number of cycles in permutations

- From Gilbert's lectures:

$$A(t, u) = \sum_n \frac{t^n}{n!} \sum_{|\sigma|=n} u^{c(\sigma)} = \frac{1}{(1-t)^u}$$

- Moments:

$$\frac{\partial^i A}{\partial u^i}(t, 1) = \sum_n t^n \mathbb{E}(S_n(S_n - 1) \dots (S_n - i + 1)) = \left( \log \frac{1}{1-t} \right)^i \frac{1}{1-t}$$

- Singularity analysis:

$$\mathbb{E}(S_n(S_n - 1) \dots (S_n - i + 1)) \sim (\log n)^i$$

$$\Rightarrow \mathbb{E}(S_n^i) \sim (\log n)^i$$

$$\frac{S_n}{\log n} \xrightarrow{d} 1$$

**Concentration phenomenon.** What are the fluctuations around  $\log n$ ?

## The number of cycles in permutations (contd.)

- Normalize  $S_n$ . As  $\mathbb{E}(S_n) \sim \log n$  and  $\mathbb{E}((S_n - \log n)^2) \sim \log n$ , define

$$\tilde{S}_n := \frac{S_n - \log n}{\sqrt{\log n}}$$



## The number of cycles in permutations (contd.)

- **Normalize  $S_n$ .** As  $\mathbb{E}(S_n) \sim \log n$  and  $\mathbb{E}((S_n - \log n)^2) \sim \log n$ , define

$$\tilde{S}_n := \frac{S_n - \log n}{\sqrt{\log n}}$$

- Study the **Laplace transform** of  $\tilde{S}_n$ :

$$\mathbb{E}(e^{x\tilde{S}_n}) = e^{-x\sqrt{\log n}} \mathbb{E}\left((e^{x/\sqrt{\log n}})^{S_n}\right) = e^{-x\sqrt{\log n}} \mathbb{E}(u^{S_n})$$

with  $u = e^{x/\sqrt{\log n}}$ .

## The number of cycles in permutations (contd.)

- **Normalize  $S_n$ .** As  $\mathbb{E}(S_n) \sim \log n$  and  $\mathbb{E}((S_n - \log n)^2) \sim \log n$ , define

$$\tilde{S}_n := \frac{S_n - \log n}{\sqrt{\log n}}$$

- Study the **Laplace transform** of  $\tilde{S}_n$ :

$$\mathbb{E}(e^{x\tilde{S}_n}) = e^{-x\sqrt{\log n}} \mathbb{E}\left((e^{x/\sqrt{\log n}})^{S_n}\right) = e^{-x\sqrt{\log n}} \mathbb{E}(u^{S_n})$$

with  $u = e^{x/\sqrt{\log n}}$ .

- **Generating function:**

$$A(t, u) = \sum_n t^n \mathbb{E}(u^{S_n}) = \frac{1}{(1-t)^u} \Rightarrow \mathbb{E}(u^{S_n}) = \frac{\Gamma(n+u)}{n! \Gamma(u)}$$

$$\mathbb{E}(e^{x\tilde{S}_n}) \sim \exp(x^2/2)$$

Laplace transform of the gaussian distribution  $\mathcal{N}(0, 1)$

**Conclusion:**

$$\tilde{S}_n := \frac{S_n - \log n}{\sqrt{\log n}} \xrightarrow{d} \mathcal{N}(0, 1)$$

## Patterns and universality in limit laws

“If the position and nature of the singularity do not change when  $u$  varies around 1, one expects a discrete limit law”

### Sub-critical composition schema for alg-log series

If  $A(t, u) = C(uB(t))$  with  $B(\rho_B) < \rho_C$ , the first singularity of  $A(t, u)$  occurs at  $\rho_B$  and we have a discrete limit law with probability g.f.:

$$G(u) = \sum \mathbb{P}(S = k)u^k = \frac{uB'(\tau u)}{B'(\tau)}$$

where  $\tau = B(\rho_B)$ .

### Examples:

- Contacts in Dyck paths

$$A(t, u) = \frac{1}{1 - u(1 - \sqrt{1 - 4t})/2}$$

- When  $C$  is the Sequence species ( $C(z) = 1/(1-z)$ ), convergence to a negative binomial law
- When  $C$  is the Set species ( $C = \exp$ ), convergence to a Poisson law

## Patterns and universality in limit laws

“If the position or nature of the singularity changes smoothly around  $u = 1$ , one expects a gaussian limit law”

### The quasi-power theorem

Assume that, uniformly in  $u$  taken in a neighborhood of 1, the probability g.f. satisfies

$$G_n(u) = \frac{1}{a(n)} [t^n] A(t, u) = \kappa(u) (\mu(u))^{\beta_n} (1 + O(\epsilon_n))$$

where  $\beta_n \rightarrow \infty$  and  $\epsilon_n \rightarrow 0$ ,  $\kappa(u)$  and  $\mu(u)$  are analytic at 1 (+ variability condition)

Let  $\mu_n = \mathbb{E}(S_n)$  and  $\sigma_n^2 = \mathbb{E}((S_n - E(S_n))^2)$ . Then the normalized variable

$$\frac{S_n - \mu_n}{\sigma_n}$$

converges to a gaussian law  $\mathcal{N}(0, 1)$

**Example:** Number of cycles in a permutation

$$G_n(u) \sim \frac{n^{u-1}}{\Gamma(u)} \sim \frac{1}{\Gamma(u)} (\exp(u-1))^{\log n}$$

## Patterns and universality in limit laws

The most interesting cases are those such that the position or nature of the singularity of  $A(t, u)$  changes non-smoothly around  $u = 1$ .

**Example.** The area of staircase polygons

$$A(t, u) = t^2 u + 2tuA(t, u) + A(t, u)A(tu, u)$$

	Type of singularity
$0 < u < 1$	simple pole
$u = 1$	square root
$u > 1$	radius 0

## Patterns and universality in limit laws

The most interesting cases are those such that the position or nature of the singularity of  $A(t, u)$  changes non-smoothly around  $u = 1$ .

**Example.** The area of staircase polygons

$$A(t, u) = t^2 u + 2tuA(t, u) + A(t, u)A(tu, u)$$

	Type of singularity	Average area in polygons weighted by $u^{a(p)}$
$0 < u < 1$	simple pole	$\kappa n$
$u = 1$	square root	$\kappa n^{3/2}$
$u > 1$	radius 0	$n^2/4$

Random staircase polygons of perimeter  $2n$  with probability

$$\text{prob}(p) = \frac{1}{Z_n(u)} u^{a(p)}$$

These cases precisely correspond to phase transitions in statistical mechanics