


Invariants for three-quadrant walks: the Kreweras trilogy

Mireille Bousquet-Mélou, CNRS, Université de Bordeaux, France




Counting walks in (rational) cones

Take a starting point p_0 in \mathbb{Z}^2 , a (finite) step set $\mathcal{S} \subset \mathbb{Z}^2$ and a cone \mathcal{C} .

Questions

- What is the number $c(n)$ of n -step walks starting at p_0 , taking their steps in \mathcal{S} and contained in \mathcal{C} ?
- For $(i,j) \in \mathcal{C}$, what is the number $c_{i,j}(n)$ of such walks that end at (i,j) ?



Counting walks in (rational) cones

Take a starting point p_0 in \mathbb{Z}^2 , a (finite) step set $\mathcal{S} \subset \mathbb{Z}^2$ and a cone \mathcal{C} .

Questions

- What is the number $c(n)$ of n -step walks starting at p_0 , taking their steps in \mathcal{S} and contained in \mathcal{C} ?
- For $(i, j) \in \mathcal{C}$, what is the number $c_{i,j}(n)$ of such walks that end at (i, j) ?

- Generating function:

$$\begin{aligned} C(x, y; t) &= \sum_{i,j,n} c_{i,j}(n) x^i y^j t^n \\ &= \sum_{w \text{ walk}} x^{i(w)} y^{j(w)} t^{|w|} \end{aligned}$$

What is the value/nature of this series?

A hierarchy of formal power series

- Rational series

$$A(t) = \frac{P(t)}{Q(t)}$$

- Algebraic series

$$P(t, A(t)) = 0$$

- Differentially finite series (D-finite)

$$\sum_{i=0}^d P_i(t) A^{(i)}(t) = 0$$

- D-algebraic series

$$P(t, A(t), A'(t), \dots, A^{(d)}(t)) = 0$$



A hierarchy of formal power series

- Rational series

$$A(t) = \frac{P(t)}{Q(t)}$$

- Algebraic series

$$P(t, A(t)) = 0$$

- Differentially finite series (D-finite)

$$\sum_{i=0}^d P_i(t) A^{(i)}(t) = 0$$

- D-algebraic series

$$P(t, A(t), A'(t), \dots, A^{(d)}(t)) = 0$$

Multi-variate series: one DE per variable




Normalizing the cone: 4 cases

- The full space: **rational** series

$$C(x,y;t) = \frac{1}{1 - tS(x,y)} = \sum_{n \geq 0} t^n S(x,y)^n,$$


where $S(x,y)$ is the **step polynomial**:

$$S(x,y) = \sum_{(i,j) \in \mathcal{S}} x^i y^j.$$




Normalizing the cone: 4 cases

- The full space: **rational** series
- A half-space: **algebraic** series
[Gessel 80]; [mbm-Petkovšek 00], [Duchon 00], [Banderier & Flajolet 02]...



Normalizing the cone: 4 cases

- The full space: **rational** series
- A half-space: **algebraic** series
[Gessel 80]; [mbm-Petkovšek 00], [Duchon 00], [Banderier & Flajolet 02]...
- A convex cone \rightarrow walks **in the non-negative quadrant**: $Q(x, y; t)$
- A non-convex cone \rightarrow walks **avoiding** the negative quadrant: $C(x, y; t)$



Walks with *small* steps



- $\mathcal{S} \subset \{\bar{1}, 0, 1\}^2 \setminus \{(0,0)\} \Rightarrow 2^8 = 256$ step sets (or: **models**)

Walks with *small* steps



- $\mathcal{S} \subset \{\bar{1}, 0, 1\}^2 \setminus \{(0,0)\} \Rightarrow 2^8 = 256$ step sets (or: **models**)
- However, some models are equivalent to a half-space problem (hence algebraic) and/or to another model (diagonal symmetry).

Walks with *small* steps



- $\mathcal{S} \subset \{\bar{1}, 0, 1\}^2 \setminus \{(0,0)\} \Rightarrow 2^8 = 256$ step sets (or: **models**)
- However, some models are equivalent to a half-space problem (hence algebraic) and/or to another model (diagonal symmetry).
- In the quadrant, one is left with **79 interesting distinct models** [mbm-Mishna 09].

Walks with *small* steps








- $\mathcal{S} \subset \{\bar{1}, 0, 1\}^2 \setminus \{(0, 0)\} \Rightarrow 2^8 = 256$ step sets (or: **models**)
- However, some models are equivalent to a half-space problem (hence algebraic) and/or to another model (diagonal symmetry).
- In the quadrant, one is left with **79 interesting distinct models** [mbm-Mishna 09].
- In the three-quadrant cone, one is left with only **74 interesting distinct models**: the 5 “singular” models become trivial.

Singular models






Non-singular



Singular

Classification of quadrant walks with small steps



The 4 algebraic models:



Kreweras



Reverse
Kreweras



Double
Kreweras




Gessel

What about walks in the three-quadrant cone?

Outline


I. Kreweras' walks in the quadrant

- Functional equation
- A solution via invariants
[Bernardi, mbm, Raschel 17(a)]



II. Kreweras' walks in three quadrants

- Functional equation
- A solution via invariants



Invariants: a method introduced by W. Tutte to count properly coloured planar triangulations (1973-1984)


Kreweras' walks in the quadrant



- Generating function:

$$Q(x,y) \equiv Q(x,y;t) = \sum_{i,j,n \geq 0} q_{i,j}(n) x^i y^j t^n$$

where $q_{i,j}(n)$ is the number of walks of length n from $(0,0)$ to (i,j) in the quadrant.



Kreweras' walks in the quadrant




- Generating function:

$$Q(x,y) \equiv Q(x,y;t) = \sum_{i,j,n \geq 0} q_{i,j}(n) x^i y^j t^n$$

where $q_{i,j}(n)$ is the number of walks of length n from $(0,0)$ to (i,j) in the quadrant.

- Functional equation (with $\bar{x} = 1/x$, $\bar{y} = 1/y$):

$$Q(x,y) = 1 + t(xy + \bar{x} + \bar{y})Q(x,y) - t\bar{x}Q(0,y) - t\bar{y}Q(x,0).$$



Kreweras' walks in the quadrant



- Generating function:

$$Q(x,y) \equiv Q(x,y;t) = \sum_{i,j,n \geq 0} q_{i,j}(n) x^i y^j t^n$$

where $q_{i,j}(n)$ is the number of walks of length n from $(0,0)$ to (i,j) in the quadrant.

- Functional equation (with $\bar{x} = 1/x$, $\bar{y} = 1/y$):

$$Q(x,y) = 1 + t(xy + \bar{x} + \bar{y})Q(x,y) - t\bar{x}Q(0,y) - t\bar{y}Q(x,0).$$

Equivalently,

$$xy(1 - t(xy + \bar{x} + \bar{y}))Q(x,y) = xy - tyQ(0,y) - txQ(x,0),$$

where

$$K(x,y) := xy(1 - t(xy + \bar{x} + \bar{y}))$$

is the kernel (a **polynomial** in t, x, y). Two catalytic variables: x and y .

What are invariants?

A pair of series $(I(x), J(y))$ in t with coefficients in $\mathbb{Q}(x)$ and $\mathbb{Q}(y)$ (respectively) is a pair of invariants if $I(x) - J(y)$ is divisible by $K(x, y)$.

- Isn't any series divisible by $K(x, y)$? After all,

$$\frac{1}{K(x, y)} = \frac{1}{xy(1 - t(xy + \bar{x} + \bar{y}))} = \bar{x}\bar{y} \sum_{n \geq 0} t^n (xy + \bar{x} + \bar{y})^n$$

is well-defined...

What are invariants?

A pair of series $(I(x), J(y))$ in t with coefficients in $\mathbb{Q}(x)$ and $\mathbb{Q}(y)$ (respectively) is a pair of invariants if $I(x) - J(y)$ is divisible by $K(x, y)$.

- Isn't any series divisible by $K(x, y)$? After all,

$$\frac{1}{K(x, y)} = \frac{1}{xy(1 - t(xy + \bar{x} + \bar{y}))} = \bar{x}\bar{y} \sum_{n \geq 0} t^n (xy + \bar{x} + \bar{y})^n$$

is well-defined...

- We want more: the coefficients (of t^n , $n \geq 0$) in the ratio

$$H(x, y) = \frac{I(x) - J(y)}{K(x, y)},$$

which are rational functions of the form $p(x, y)/(d(x)d'(y))$, must have **poles of bounded order** at $x = 0$ and $y = 0$.

Equivalently, there exists i, j such that the coefficients of $x^i y^j H(x, y)$ have no pole at $x = 0$ nor $y = 0$.

Do such pairs exist?



- Trivial invariants: take $I(x) = J(y) \in \mathbb{Q}((t))$

Do such pairs exist?



- Trivial invariants: take $I(x) = J(y) \in \mathbb{Q}((t))$
- Basic rational invariants: let

$$I_0(x) = \bar{x}^2 - \bar{x}/t - x, \quad J_0(y) = I_0(y).$$

Then

$$K(x, y) \cdot \frac{x-y}{x^2 y^2} \cdot \frac{1}{t} = I_0(x) - J_0(y),$$

and the series $H_0(x, y) = (x-y)/(tx^2y^2)$ has poles of bounded order at $x=0$ and $y=0$.

Do such pairs exist?



- Trivial invariants: take $I(x) = J(y) \in \mathbb{Q}((t))$
- Basic rational invariants: let

$$I_0(x) = \bar{x}^2 - \bar{x}/t - x, \quad J_0(y) = I_0(y).$$

Then

$$K(x, y) \cdot \frac{x-y}{x^2 y^2} \cdot \frac{1}{t} = I_0(x) - J_0(y),$$

and the series $H_0(x, y) = (x-y)/(tx^2y^2)$ has poles of bounded order at $x=0$ and $y=0$.

New invariants from old ones

The componentwise sum (resp. product) of two pairs of invariants $(I_0(x), J_0(y)), (I_1(x), J_1(y))$ is another pair of invariants.

$$(I_0(x) + I_1(x), J_0(y) + J_1(y)), \quad (I_0(x)I_1(x), J_0(y)J_1(y))$$

What are invariants good for?

Functional equation:

$$K(x,y)Q(x,y) = xy - tyQ(0,y) - txQ(x,0).$$

What are invariants good for?

Functional equation:

$$K(x,y)Q(x,y) = xy - tyQ(0,y) - txQ(x,0).$$

1/ Build a new pair of invariants:

$$\begin{aligned}K(x,y)H_1(x,y) &= F(x) + G(y) - tyQ(0,y) - txQ(x,0) \\ &= I_1(x) - J_1(y)\end{aligned}$$

with

$$I_1(x) = F(x) - txQ(x,0), \quad J_1(y) = -G(y) + tyQ(0,y).$$

What are invariants good for?

Functional equation:

$$K(x,y)Q(x,y) = xy - tyQ(0,y) - txQ(x,0).$$

1/ Build a new pair of invariants:

$$\begin{aligned}K(x,y)H_1(x,y) &= F(x) + G(y) - tyQ(0,y) - txQ(x,0) \\ &= I_1(x) - J_1(y)\end{aligned}$$

with

$$I_1(x) = F(x) - txQ(x,0), \quad J_1(y) = -G(y) + tyQ(0,y).$$

2/ Relate this pair to the basic rational invariants $(I_0(x), J_0(y))$ using:

The invariant lemma

If $K(x,y)H(x,y) = I(x) - J(y)$ and the coefficients of $H(x,y)$ (in t) have no pole at $x = 0$ nor at $y = 0$, then

$$H(x,y) = 0 \quad \text{and} \quad I(x) = J(y) \in \mathbb{Q}((t)).$$

Step 1: decoupling



Functional equation:

$$K(x,y)Q(x,y) = xy - tyQ(0,y) - txQ(x,0).$$

1/ Build a new pair of invariants:

$$\begin{aligned} K(x,y)H_1(x,y) &= F(x) + G(y) - tyQ(0,y) - txQ(x,0) \\ &= I_1(x) - J_1(y) \end{aligned}$$

Step 1: decoupling



Functional equation:

$$K(x,y)Q(x,y) = xy - tyQ(0,y) - txQ(x,0).$$

1/ Build a new pair of invariants:

$$\begin{aligned} K(x,y)H_1(x,y) &= F(x) + G(y) - tyQ(0,y) - txQ(x,0) \\ &= I_1(x) - J_1(y) \end{aligned}$$

Easy! The term xy decouples modulo $K(x,y)$:

$$-\frac{1}{txy}K(x,y) = xy + \bar{x} + \bar{y} - 1/t.$$

So we have a new pair of invariants:

$$I_1(x) = txQ(x,0) + \bar{x} - 1/(2t), \quad J_1(y) = -I_1(y).$$

Step 2: relate (I_1, J_1) to the basic invariants (I_0, J_0)

We have two pairs of invariants:

$$\begin{aligned} I_0(x) &= \bar{x}^2 - \bar{x}/t - x, & J_0(y) &= I_0(y), \\ I_1(x) &= \bar{x} - 1/(2t) + txQ(x,0), & J_1(y) &= -I_1(y). \end{aligned}$$

The series $I_1(x)$ has valuation -1 in x , while $I_0(x)$ has valuation -2 .

Step 2: relate (I_1, J_1) to the basic invariants (I_0, J_0)

We have two pairs of invariants:

$$\begin{aligned}I_0(x) &= \bar{x}^2 - \bar{x}/t - x, & J_0(y) &= I_0(y), \\I_1(x) &= \bar{x} - 1/(2t) + txQ(x,0), & J_1(y) &= -I_1(y).\end{aligned}$$

The series $I_1(x)$ has valuation -1 in x , while $I_0(x)$ has valuation -2 .

Observation: $I_1(x)^2 - I_0(x)$ has no pole! Define

$$I(x) := I_1(x)^2 - I_0(x) \quad J(y) := J_1(y)^2 - J_0(y).$$

Then $(I(x), J(y))$ is a pair of invariants,

Step 2: relate (I_1, J_1) to the basic invariants (I_0, J_0)

We have two pairs of invariants:

$$\begin{aligned} I_0(x) &= \bar{x}^2 - \bar{x}/t - x, & J_0(y) &= I_0(y), \\ I_1(x) &= \bar{x} - 1/(2t) + txQ(x,0), & J_1(y) &= -I_1(y). \end{aligned}$$

The series $I_1(x)$ has valuation -1 in x , while $I_0(x)$ has valuation -2 .

Observation: $I_1(x)^2 - I_0(x)$ has no pole! Define

$$I(x) := I_1(x)^2 - I_0(x) \quad J(y) := J_1(y)^2 - J_0(y).$$

Then $(I(x), J(y))$ is a pair of invariants,

$$-K(x, y)H(x, y) = I(x) - J(y),$$

where

$$H(x, y) = \frac{x-y}{tx^2y^2} + \left(Q(x, y) + \frac{1}{xyt} \right) (I_1(x) + J_1(y))$$

has no pole at $x = 0$ nor $y = 0$.

Applying the invariant lemma



Hence

$$I(x) = I_1(x)^2 - I_0(x) = (txQ(x,0) - 1/(2t))^2 + 2tQ(x,0) + x$$

is a series of $\mathbb{Q}((t))$, equal to its value at $x = 0$:

Applying the invariant lemma



Hence

$$I(x) = I_1(x)^2 - I_0(x) = (txQ(x,0) - 1/(2t))^2 + 2tQ(x,0) + x$$

is a series of $\mathbb{Q}((t))$, equal to its value at $x = 0$:

$$(txQ(x,0) - 1/(2t))^2 + 2tQ(x,0) + x = 1/(2t)^2 + 2tQ(0,0).$$

We have obtained an equation in *one* catalytic variable

Applying the invariant lemma



Hence

$$I(x) = I_1(x)^2 - I_0(x) = (txQ(x,0) - 1/(2t))^2 + 2tQ(x,0) + x$$

is a series of $\mathbb{Q}((t))$, equal to its value at $x = 0$:

$$(txQ(x,0) - 1/(2t))^2 + 2tQ(x,0) + x = 1/(2t)^2 + 2tQ(0,0).$$

We have obtained an equation in *one* catalytic variable
 \Rightarrow systematic solution (via Brown's quadratic method, or
[mbm-Jehanne 06])

Applying the invariant lemma



Hence

$$I(x) = I_1(x)^2 - I_0(x) = (txQ(x,0) - 1/(2t))^2 + 2tQ(x,0) + x$$

is a series of $\mathbb{Q}((t))$, equal to its value at $x = 0$:

$$(txQ(x,0) - 1/(2t))^2 + 2tQ(x,0) + x = 1/(2t)^2 + 2tQ(0,0).$$

We have obtained an equation in *one* catalytic variable
 \Rightarrow systematic solution (via Brown's quadratic method, or
[mbm-Jehanne 06])

GF of Kreweras' walks in the quadrant

Let $Z \equiv Z(t)$ be the only series in t such that $Z = t(2 + Z^3)$. Then

$$Q(x,0) = \frac{1}{tx} \left(\frac{1}{2t} - \frac{1}{x} - \left(\frac{1}{Z} - \frac{1}{x} \right) \sqrt{1 - xZ^2} \right).$$

[Kreweras 65], [Gessel 86], [mbm 05]...

Other quadrant models with small steps

Two ingredients:

- A pair $(I_0(x), J_0(y))$ of rational invariants \Rightarrow 23 models with a finite group
- A decoupling identity of the form

$$xy = F(x) + G(y) + K(x,y)H(x,y)$$

\Rightarrow 4 of the 23 models

The use of invariants solves these four models, proving algebraicity.



Kreweras



Reverse
Kreweras



Double
Kreweras



Gessel



- Basic rational invariants

$$\mathcal{H}(x, y) \frac{x-y}{txy} = \mathcal{I}_0(x) - \mathcal{I}_0(y)$$

with

$$\mathcal{I}_0(x) = \bar{x} + x/t - x^2, \quad \mathcal{I}_0(y) = \mathcal{I}_0(y).$$



- Basic rational invariants

$$\mathcal{K}(x, y) \frac{x-y}{txy} = \mathcal{I}_0(x) - \mathcal{I}_0(y)$$

with

$$\mathcal{I}_0(x) = \bar{x} + x/t - x^2, \quad \mathcal{I}_0(y) = \mathcal{I}_0(y).$$

- Invariants related to $\mathcal{Q}(x, y)$:

$$-\frac{1}{ty} \mathcal{K}(x, y) (1 + ty\mathcal{Q}(x, y)) = \mathcal{I}_1(x) - \mathcal{I}_1(y)$$

with

$$\mathcal{I}_1(x) = t\mathcal{Q}(x, 0) - x/t + x^2, \quad \mathcal{I}_1(y) = -t\mathcal{Q}(0, y) - \bar{y} + t\mathcal{Q}(0, 0)$$



- Basic rational invariants

$$\mathcal{K}(x, y) \frac{x-y}{txy} = \mathcal{I}_0(x) - \mathcal{I}_0(y)$$

with

$$\mathcal{I}_0(x) = \bar{x} + x/t - x^2, \quad \mathcal{I}_0(y) = \mathcal{I}_0(y).$$

- Invariants related to $\mathcal{Q}(x, y)$:

$$-\frac{1}{ty} \mathcal{K}(x, y) (1 + ty\mathcal{Q}(x, y)) = \mathcal{I}_1(x) - \mathcal{I}_1(y)$$


with

$$\mathcal{I}_1(x) = t\mathcal{Q}(x, 0) - x/t + x^2, \quad \mathcal{I}_1(y) = -t\mathcal{Q}(0, y) - \bar{y} + t\mathcal{Q}(0, 0)$$

GF of reverse Kreweras walks [Mishna 09, mbm-Mishna 10]

$$\mathcal{Q}(x, 0) = \frac{Z(4-Z^3)}{16t} - \frac{t-x^2+tx^3}{2xt^2} - \frac{(2x^2-xZ^2-Z)\sqrt{1-xZ(Z^3+4)/4+x^2Z^2/4}}{2txZ}.$$

II. Kreweras' walks in three quadrants



$$C(x,y) \equiv C(x,y;t) = \sum_{i,j,n \geq 0} c_{i,j}(n) x^i y^j t^n$$

where $c_{i,j}(n)$ is the number of walks of length n from $(0,0)$ to (i,j) avoiding the negative quadrant.

First functional equation




Functional equation (with $\bar{x} = 1/x, \bar{y} = 1/y$):

$$C(x, y) = 1 + t(xy + \bar{x} + \bar{y})C(x, y) - t\bar{x}C_-(\bar{y}) - t\bar{y}C_-(\bar{x}),$$

where

$$C_-(\bar{x}) = \sum_{i < 0, n \geq 0} c_{-i, 0}(n) x^i t^n$$

counts walks ending on the negative x-axis.



First functional equation



Functional equation (with $\bar{x} = 1/x$, $\bar{y} = 1/y$):

$$C(x, y) = 1 + t(xy + \bar{x} + \bar{y})C(x, y) - t\bar{x}C_-(\bar{y}) - t\bar{y}C_-(\bar{x}),$$

where

$$C_-(\bar{x}) = \sum_{i < 0, n \geq 0} c_{-i, 0}(n) x^i t^n$$

counts walks ending on the negative x -axis.

Equivalently,

$$xy(1 - t(xy + \bar{x} + \bar{y}))C(x, y) = xy - tyC_-(\bar{y}) - txC_-(\bar{x}).$$

- The coefficients of $C(x, y)$ have poles of arbitrarily large order at $x = 0$ and $y = 0$.
- It suffices to determine $C_-(\bar{x})$, the GF of walks ending on the negative x -axis.


Alternative: Split the cone in two convex cones



- À la Raschel-Trotignon (2019):

$$C(x, y) = \bar{x}U(\bar{x}, xy) + D(xy) + \bar{y}U(\bar{y}, xy),$$

where $U(x, y) \in \mathbb{Q}[x, y][[t]]$ and $D(x) \in \mathbb{Q}[x][[t]]$.



Alternative: Split the cone in two convex cones




- À la Raschel-Trotignon (2019):

$$C(x, y) = \bar{x}U(\bar{x}, xy) + D(xy) + \bar{y}U(\bar{y}, xy),$$

where $U(x, y) \in \mathbb{Q}[x, y][[t]]$ and $D(x) \in \mathbb{Q}[x][[t]]$.

- Step-by-step construction:

$$\begin{aligned} \bar{x}U(\bar{x}, xy) &= t(xy + \bar{x} + \bar{y})\bar{x}U(\bar{x}, xy) \\ &\quad + t\bar{x}D(xy) \\ &\quad - t\bar{y}(\bar{x}U(\bar{x}, 0) + \bar{x}U(0, xy) - \bar{x}U(0, 0)). \end{aligned}$$



Alternative: Split the cone in two convex cones



- À la Raschel-Trotignon (2019):

$$C(x, y) = \bar{x}U(\bar{x}, xy) + D(xy) + \bar{y}U(\bar{y}, xy),$$


where $U(x, y) \in \mathbb{Q}[x, y][[t]]$ and $D(x) \in \mathbb{Q}[x][[t]]$.

- Step-by-step construction:

$$\begin{aligned} \bar{x}U(\bar{x}, xy) &= t(xy + \bar{x} + \bar{y})\bar{x}U(\bar{x}, xy) \\ &\quad + t\bar{x}D(xy) \\ &\quad - t\bar{y}(\bar{x}U(\bar{x}, 0) + \bar{x}U(0, xy) - \bar{x}U(0, 0)). \end{aligned}$$

Analogously,

$$D(xy) = 1 + txyD(xy) + 2t\bar{y}(\bar{x}U(0, xy) - \bar{x}U(0, 0)).$$



Split the cone in two convex cones

- Two equations:

$$\begin{aligned}\bar{x}(1 - t(xy + \bar{x} + \bar{y}))U(\bar{x}, xy) &= -t\bar{y}\bar{x}U(\bar{x}, 0) + t\bar{x}D(xy) \\ &\quad - t\bar{y}\bar{x}U(0, xy) + t\bar{y}\bar{x}U(0, 0), \\ (1 - txy)D(xy) &= 1 + 2t\bar{y}\bar{x}U(0, xy) - 2t\bar{y}\bar{x}U(0, 0).\end{aligned}$$

Split the cone in two convex cones

- Two equations:

$$\begin{aligned}\bar{x}(1 - t(xy + \bar{x} + \bar{y}))U(\bar{x}, xy) &= -t\bar{y}\bar{x}U(\bar{x}, 0) + t\bar{x}D(xy) \\ &\quad - t\bar{y}\bar{x}U(0, xy) + t\bar{y}\bar{x}U(0, 0), \\ (1 - txy)D(xy) &= 1 + 2t\bar{y}\bar{x}U(0, xy) - 2t\bar{y}\bar{x}U(0, 0).\end{aligned}$$

- Straightening: $(\bar{x}, xy) \mapsto (x, y)$

$$\begin{aligned}x(1 - t(y + x + \bar{x}\bar{y}))U(x, y) &= -t\bar{y}U(x, 0) + txD(y) \\ &\quad - t\bar{y}U(0, y) + t\bar{y}U(0, 0), \\ (1 - ty)D(y) &= 1 + 2t\bar{y}U(0, y) - 2t\bar{y}U(0, 0).\end{aligned}$$

Split the cone in two convex cones

- Two equations:

$$\begin{aligned} \bar{x}(1 - t(xy + \bar{x} + \bar{y}))U(\bar{x}, xy) &= -t\bar{y}\bar{x}U(\bar{x}, 0) + t\bar{x}D(xy) \\ &\quad - t\bar{y}\bar{x}U(0, xy) + t\bar{y}\bar{x}U(0, 0), \\ (1 - txy)D(xy) &= 1 + 2t\bar{y}\bar{x}U(0, xy) - 2t\bar{y}\bar{x}U(0, 0). \end{aligned}$$

- Straightening: $(\bar{x}, xy) \mapsto (x, y)$

$$\begin{aligned} x(1 - t(y + x + \bar{x}\bar{y}))U(x, y) &= -t\bar{y}U(x, 0) + txD(y) \\ &\quad - t\bar{y}U(0, y) + t\bar{y}U(0, 0), \\ (1 - ty)D(y) &= 1 + 2t\bar{y}U(0, y) - 2t\bar{y}U(0, 0). \end{aligned}$$

- Linear combination:

$$2\mathcal{K}(x, y)U(x, y) = y + (ty + 2tx - 1)yD(y) - 2tU(x, 0),$$

where now $\mathcal{K}(x, y) = xy(1 - t(x + y + \bar{x}\bar{y}))$ is the kernel of **reverse Kreweras'** walks.

We want to construct \mathcal{K} -invariants...

$$2\mathcal{K}(x,y)U(x,y) = y + (ty + 2tx - 1)yD(y) - 2tU(x,0)$$

Two new difficulties:

- one needs to “decouple” y , in some sense,
- the term $(ty + 2tx - 1)$ depends on x (and y).

We want to construct \mathcal{K} -invariants...

$$2\mathcal{K}(x,y)U(x,y) = y + (ty + 2tx - 1)yD(y) - 2tU(x,0)$$

Two new difficulties:

- one needs to “decouple” y , in some sense,
- the term $(ty + 2tx - 1)$ depends on x (and y).

Two good news:

- Decoupling:

$$y = (ty + 2tx - 1)G(y) - F(x)$$

with $G(y) = 1/t$ and $F(x) = 2x - 1/t$.

We want to construct \mathcal{K} -invariants...

$$2\mathcal{K}(x,y)U(x,y) = y + (ty + 2tx - 1)yD(y) - 2tU(x,0)$$

Two new difficulties:

- one needs to “decouple” y , in some sense,
- the term $(ty + 2tx - 1)$ depends on x (and y).

Two good news:

- Decoupling:

$$y = (ty + 2tx - 1)G(y) - F(x)$$

with $G(y) = 1/t$ and $F(x) = 2x - 1/t$.

- The square of $(ty + 2tx - 1)$ is a function of y (modulo $\mathcal{K}(x,y)$):

$$(ty + 2tx - 1)^2 = \Delta(y) - 4t\bar{y}\mathcal{K}(x,y)$$

where $\Delta(y) = (1 - ty)^2 - 4t\bar{y}$.

A new pair of \mathcal{K} -invariants

$$2\mathcal{K}(x,y)U(x,y) = y + (ty + 2tx - 1)yD(y) - 2tU(x,0)$$

- Decoupling: since $y = (ty + 2tx - 1)/t - 2x + 1/t$,

$$2\mathcal{K}(x,y)U(x,y) = (ty + 2tx - 1)(yD(y) + 1/t) - (2tU(x,0) + 2x - 1/t)$$

A new pair of \mathcal{K} -invariants

$$2\mathcal{K}(x,y)U(x,y) = y + (ty + 2tx - 1)yD(y) - 2tU(x,0)$$

- Decoupling: since $y = (ty + 2tx - 1)/t - 2x + 1/t$,

$$2\mathcal{K}(x,y)U(x,y) = (ty + 2tx - 1)(yD(y) + 1/t) - (2tU(x,0) + 2x - 1/t)$$

- Since $(ty + 2tx - 1)^2 = \Delta(y) - 4t\bar{y}\mathcal{K}(x,y)$, multiply out by

$$(ty + 2tx - 1)(yD(y) + 1/t) + (2tU(x,0) + 2x - 1/t).$$

A new pair of \mathcal{K} -invariants

$$2\mathcal{K}(x,y)U(x,y) = y + (ty + 2tx - 1)yD(y) - 2tU(x,0)$$

- Decoupling: since $y = (ty + 2tx - 1)/t - 2x + 1/t$,

$$2\mathcal{K}(x,y)U(x,y) = (ty + 2tx - 1)(yD(y) + 1/t) - (2tU(x,0) + 2x - 1/t)$$

- Since $(ty + 2tx - 1)^2 = \Delta(y) - 4t\bar{y}\mathcal{K}(x,y)$, multiply out by

$$(ty + 2tx - 1)(yD(y) + 1/t) + (2tU(x,0) + 2x - 1/t).$$

- We have found a new pair of \mathcal{K} -invariants:

$$\mathcal{I}_2(x) = \left(2tU(x,0) + 2x - \frac{1}{t}\right)^2, \quad \mathcal{I}_2(y) = \Delta(y) \left(yD(y) + \frac{1}{t}\right)^2$$

with $\Delta(y) = (1 - ty)^2 - 4t\bar{y}$.

Two known pairs of \mathcal{K} -invariants (from Part I)

- Recall:

$$\begin{aligned}\mathcal{I}_0(x) &= \bar{x} + x/t - x^2, & \mathcal{I}_0(x) &= \mathcal{I}_0(y), \\ \mathcal{I}_1(x) &= t\mathcal{Q}(x,0) - x/t + x^2, & \mathcal{I}_1(y) &= -\bar{y} - t\mathcal{Q}(0,y) + t\mathcal{Q}(0,0).\end{aligned}$$

Two known pairs of \mathcal{K} -invariants (from Part I)

- Recall:

$$\mathcal{F}_0(x) = \bar{x} + x/t - x^2, \quad \mathcal{F}_0(y) = \mathcal{F}_0(x),$$

$$\mathcal{F}_1(x) = t\mathcal{Q}(x,0) - x/t + x^2, \quad \mathcal{F}_1(y) = -\bar{y} - t\mathcal{Q}(0,y) + t\mathcal{Q}(0,0).$$

- We have just found another pair:

$$\mathcal{F}_2(x) = \left(2tU(x,0) + 2x - \frac{1}{t}\right)^2 = \mathcal{O}(x^0),$$

$$\mathcal{F}_2(y) = \Delta(y) \left(yD(y) + \frac{1}{t}\right)^2 = -4\bar{y} + \mathcal{O}(y^0)$$

Two known pairs of \mathcal{K} -invariants (from Part I)

- Recall:

$$\mathcal{I}_0(x) = \bar{x} + x/t - x^2, \quad \mathcal{I}_0(y) = \mathcal{I}_0(x),$$

$$\mathcal{I}_1(x) = t\mathcal{Q}(x,0) - x/t + x^2, \quad \mathcal{I}_1(y) = -\bar{y} - t\mathcal{Q}(0,y) + t\mathcal{Q}(0,0).$$

- We have just found another pair:

$$\mathcal{I}_2(x) = \left(2tU(x,0) + 2x - \frac{1}{t}\right)^2 = \mathcal{O}(x^0),$$

$$\mathcal{I}_2(y) = \Delta(y) \left(yD(y) + \frac{1}{t}\right)^2 = -4\bar{y} + \mathcal{O}(y^0)$$

Two known pairs of \mathcal{K} -invariants (from Part I)

- Recall:

$$\begin{aligned}\mathcal{F}_0(x) &= \bar{x} + x/t - x^2, & \mathcal{F}_0(y) &= \mathcal{F}_0(y), \\ \mathcal{F}_1(x) &= t\mathcal{Q}(x,0) - x/t + x^2, & \mathcal{F}_1(y) &= -\bar{y} - t\mathcal{Q}(0,y) + t\mathcal{Q}(0,0).\end{aligned}$$

- We have just found another pair:

$$\begin{aligned}\mathcal{F}_2(x) &= \left(2tU(x,0) + 2x - \frac{1}{t}\right)^2 = \mathcal{O}(x^0), \\ \mathcal{F}_2(y) &= \Delta(y) \left(yD(y) + \frac{1}{t}\right)^2 = -4\bar{y} + \mathcal{O}(y^0)\end{aligned}$$

- Define

$$\mathcal{F}(x) = \mathcal{F}_2(x) - 4\mathcal{F}_1(x), \quad \mathcal{F}(y) = \mathcal{F}_2(y) - 4\mathcal{F}_1(y).$$

Then $(\mathcal{F}(x), \mathcal{F}(y))$ is a pair of invariants with no pole at 0.

Two known pairs of \mathcal{K} -invariants (from Part I)

- Recall:

$$\begin{aligned}\mathcal{F}_0(x) &= \bar{x} + x/t - x^2, & \mathcal{F}_0(y) &= \mathcal{F}_0(y), \\ \mathcal{F}_1(x) &= t\mathcal{Q}(x,0) - x/t + x^2, & \mathcal{F}_1(y) &= -\bar{y} - t\mathcal{Q}(0,y) + t\mathcal{Q}(0,0).\end{aligned}$$

- We have just found another pair:

$$\begin{aligned}\mathcal{F}_2(x) &= \left(2tU(x,0) + 2x - \frac{1}{t}\right)^2 = \mathcal{O}(x^0), \\ \mathcal{F}_2(y) &= \Delta(y) \left(yD(y) + \frac{1}{t}\right)^2 = -4\bar{y} + \mathcal{O}(y^0)\end{aligned}$$

- Define

$$\mathcal{F}(x) = \mathcal{F}_2(x) - 4\mathcal{F}_1(x), \quad \mathcal{F}(y) = \mathcal{F}_2(y) - 4\mathcal{F}_1(y).$$

Then $(\mathcal{F}(x), \mathcal{F}(y))$ is a pair of invariants with no pole at 0. Moreover,

$$\mathcal{K}(x,y)\mathcal{H}(x,y) = \mathcal{F}(x) - \mathcal{F}(y)$$

where $\mathcal{H}(x,y)$ has no pole at 0 either

Two known pairs of \mathcal{H} -invariants (from Part I)

- Recall:

$$\begin{aligned}\mathcal{F}_0(x) &= \bar{x} + x/t - x^2, & \mathcal{F}_0(y) &= \mathcal{F}_0(y), \\ \mathcal{F}_1(x) &= t\mathcal{Q}(x,0) - x/t + x^2, & \mathcal{F}_1(y) &= -\bar{y} - t\mathcal{Q}(0,y) + t\mathcal{Q}(0,0).\end{aligned}$$

- We have just found another pair:

$$\begin{aligned}\mathcal{F}_2(x) &= \left(2tU(x,0) + 2x - \frac{1}{t}\right)^2 = \mathcal{O}(x^0), \\ \mathcal{F}_2(y) &= \Delta(y) \left(yD(y) + \frac{1}{t}\right)^2 = -4\bar{y} + \mathcal{O}(y^0)\end{aligned}$$

- Define

$$\mathcal{F}(x) = \mathcal{F}_2(x) - 4\mathcal{F}_1(x), \quad \mathcal{F}(y) = \mathcal{F}_2(y) - 4\mathcal{F}_1(y).$$

Then $(\mathcal{F}(x), \mathcal{F}(y))$ is a pair of invariants with no pole at 0. Moreover,

$$\mathcal{H}(x,y)\mathcal{H}(x,y) = \mathcal{F}(x) - \mathcal{F}(y)$$

where $\mathcal{H}(x,y)$ has no pole at 0 either $\Rightarrow \mathcal{F}(x)$ and $\mathcal{F}(y)$ are trivial.

Trivial invariants for reverse Kreweras' steps

Conclusion:

$$\mathcal{F}_2(x) = \left(2tU(x,0) + 2x - \frac{1}{t}\right)^2 = 4\left(t\mathcal{Q}(x,0) - x/t + x^2\right) + cst,$$

$$\mathcal{F}_2(y) = \Delta(y) \left(yD(y) + \frac{1}{t}\right)^2 = 4(-\bar{y} - t\mathcal{Q}(0,y) + t\mathcal{Q}(0,0)) + cst,$$

with $\Delta(y) = (1 - ty)^2 - 4t\bar{y}$.

The constant can be determined by specializing y to the unique root of $\Delta(y)$ that is a power series in t .

The GF of Kreweras walks in three quadrants [mbm]

- Walks ending on the negative x-axis: series $U(x,0)$, with

$$\frac{1}{2} \left(2tU(x,0) + 2x - \frac{1}{t} \right)^2 = \frac{(1-Z^3)^{3/2}}{Z^2} + (1-xZ)^2 \left(\frac{1}{Z^2} - \frac{1}{x} \right) + \left(\bar{x} + Z - \frac{2x}{Z} \right) \sqrt{1 - Z \frac{4+Z^3}{4} x + \frac{Z^2}{4} x^2}.$$

- Walks ending on the diagonal: series $D(x)$, with

$$\frac{\Delta(x)}{2} \left(xD(x) + \frac{1}{t} \right)^2 = \frac{(1-Z^3)^{3/2}}{Z^2} + (1-xZ)^2 \left(\frac{1}{Z^2} - \frac{1}{x} \right) - \left(\bar{x} + Z - \frac{2x}{Z} \right) \sqrt{1 - Z \frac{4+Z^3}{4} x + \frac{Z^2}{4} x^2}.$$

where $\Delta(x) = (1-tx)^2 - 4t\bar{x}$ and $Z = t(2+Z^3)$.

The GF of Kreweras walks in three quadrants [mbm]

- Walks ending on the negative x-axis: series $U(x,0)$, with

$$\frac{1}{2} \left(2tU(x,0) + 2x - \frac{1}{t} \right)^2 = \frac{(1-Z^3)^{3/2}}{Z^2} + (1-xZ)^2 \left(\frac{1}{Z^2} - \frac{1}{x} \right) + \left(\bar{x} + Z - \frac{2x}{Z} \right) \sqrt{1 - Z \frac{4+Z^3}{4} x + \frac{Z^2}{4} x^2}.$$

- Walks ending on the diagonal: series $D(x)$, with

$$\frac{\Delta(x)}{2} \left(xD(x) + \frac{1}{t} \right)^2 = \frac{(1-Z^3)^{3/2}}{Z^2} + (1-xZ)^2 \left(\frac{1}{Z^2} - \frac{1}{x} \right) - \left(\bar{x} + Z - \frac{2x}{Z} \right) \sqrt{1 - Z \frac{4+Z^3}{4} x + \frac{Z^2}{4} x^2}.$$

where $\Delta(x) = (1-tx)^2 - 4t\bar{x}$ and $Z = t(2+Z^3)$.

- All walks in three quadrants:

$$xy(1 - t(\bar{x} + \bar{y} + xy))C(x,y) = xy - tU(\bar{x},0) - tU(\bar{y},0).$$

(Algebraicity of excursions proved by [Elvey Price, FPSAC 20])



- Number of n -step walks ending at (i, j) in the three quadrant plane:

$$c_{i,j}(n) \sim -\frac{H_{i,j}}{\Gamma(-3/4)} 3^n n^{-7/4} \quad (\text{for } n + i + j \equiv 0 \pmod{3})$$



- Number of n -step walks ending at (i, j) in the three quadrant plane:

$$c_{i,j}(n) \sim -\frac{H_{i,j}}{\Gamma(-3/4)} 3^n n^{-7/4} \quad (\text{for } n+i+j \equiv 0 \pmod{3})$$

- The generating function

$$\mathcal{H}(x, y) := \sum_{j \geq 0, i \leq j} H_{i,j} x^{j-i} y^j,$$

is algebraic of degree 16, given by

$$(1 + xy^2 + x^2y - 3xy) \mathcal{H}(x, y) = \mathcal{H}_-(x) + \frac{1}{2} (2 + xy^2 - 3xy) \mathcal{H}_d(y)$$

where

$$\mathcal{H}_-(x) := \sum_{i > 0} H_{-i,0} x^i$$

$$\mathcal{H}_d(y) := \sum_{i \geq 0} H_{i,i} y^i$$



- Number of n -step walks ending at (i, j) in the three quadrant plane:

$$c_{i,j}(n) \sim -\frac{H_{i,j}}{\Gamma(-3/4)} 3^n n^{-7/4} \quad (\text{for } n+i+j \equiv 0 \pmod{3})$$

- The generating function

$$\mathcal{H}(x, y) := \sum_{j \geq 0, i \leq j} H_{i,j} x^{j-i} y^j,$$

is algebraic of degree 16, given by

$$(1 + xy^2 + x^2y - 3xy) \mathcal{H}(x, y) = \mathcal{H}_-(x) + \frac{1}{2} (2 + xy^2 - 3xy) \mathcal{H}_d(y)$$

where

$$\mathcal{H}_-(x) := \sum_{i > 0} H_{-i,0} x^i = \frac{9x}{2} \sqrt{\frac{1+2x}{1-x}} \sqrt{\frac{4-x}{1-x}} + 2,$$

$$\mathcal{H}_d(y) := \sum_{i \geq 0} H_{i,i} y^i = \frac{9}{(1-y)\sqrt{y(4-y)}} \sqrt{\frac{1+2y}{1-y}} \sqrt{\frac{4-y}{1-y}} - 2.$$



- Number of n -step walks ending at (i, j) in the three quadrant plane:

$$c_{i,j}(n) \sim -\frac{H_{i,j}}{\Gamma(-3/4)} 3^n n^{-7/4} \quad (\text{for } n+i+j \equiv 0 \pmod{3})$$

- The generating function

$$\mathcal{H}(x, y) := \sum_{j \geq 0, i \leq j} H_{i,j} x^{j-i} y^j,$$

is algebraic of degree 16, given by

$$(1 + xy^2 + x^2y - 3xy) \mathcal{H}(x, y) = \mathcal{H}_-(x) + \frac{1}{2} (2 + xy^2 - 3xy) \mathcal{H}_d(y)$$

where

$$\mathcal{H}_-(x) := \sum_{i > 0} H_{-i,0} x^i = \frac{9x}{2} \sqrt{\frac{1+2x}{1-x} \sqrt{\frac{4-x}{1-x}}} + 2, \quad [\text{Trotignon 19(a)}]$$

$$\mathcal{H}_d(y) := \sum_{i \geq 0} H_{i,i} y^i = \frac{9}{(1-y)\sqrt{y(4-y)}} \sqrt{\frac{1+2y}{1-y} \sqrt{\frac{4-y}{1-y}}} - 2.$$

Other three-quadrant models with small steps

Two new ingredients:

- x/y -symmetry of the step set (for the functional equation)
- A decoupling identity of the form

$$y = p(x,y)G(y) + F(x) + K(x,y)H(x,y)$$



The use of invariants solves these 3 models, proving algebraicity:



Kreweras



Reverse
Kreweras



Double
Kreweras

D-finite: [Raschel-Trotignon 19]

Other three-quadrant models with small steps

Two new ingredients:

- x/y -symmetry of the step set (for the functional equation)
- A decoupling identity of the form

$$y = p(x,y)G(y) + F(x) + K(x,y)H(x,y)$$



The use of invariants solves these 3 models, proving algebraicity:



Kreweras



Reverse
Kreweras



Double
Kreweras



Gessel
TO DO

What's next?

- Gessel's step set (algebraicity of excursions [Budd 20])



What's next?

- Gessel's step set (algebraicity of excursions [Budd 20])



- [Beyond finite groups] Another x/y -symmetric model with an infinite group but a "weak" invariant

(D-algebraic [Dreyfus-Trotignon 20(a)])



What's next?

- **Gessel's step set** (algebraicity of excursions [Budd 20])



- **[Beyond finite groups]** Another x/y -symmetric model with an infinite group but a "weak" invariant

(D-algebraic [Dreyfus-Trotignon 20(a)])




- **[Beyond purely algebraic solutions]** Revisit some solved D-finite cases, e.g. NSEW walks, where the crux is to prove the algebraicity of the series $A(x, y)$ defined by:

$$xy(1 - t(x + \bar{x} + y + \bar{y}))A(x, y) = \frac{1}{3}(2xy + \bar{x}y + x\bar{y}) - txA_-(\bar{x}) - tyA_-(\bar{y}).$$



[mbm 16], [mbm-Wallner 21]

Partial classification of three quadrant walks



+ results for excursions [Budd 20, Elvey Price 20]



algebraic [mbm]

DF transc. [mbm 16, Raschel-Trotignon 19, mbm-Wallner 21]

Non-D-finite [Mustapha 19]

D-alg./not D-alg [Dreyfus-Trotignon 20(a)]