

On self-avoiding walks

Mireille Bousquet-Mélou

CNRS, LaBRI, Bordeaux, France

<http://www.labri.fr/~bousquet>

Outline

I. Self-avoiding walks (SAW): Generalities, predictions and results

II. Some exactly solvable models of SAW

II.0 A toy model: Partially directed walks


II.1 Weakly directed walks

II.2 Prudent walks


II.3 Two related models

III. Au dessert : SAW on the honeycomb lattice,
d'après Duminil-Copin et Smirnov

I. Generalities



Self-avoiding walks (SAW)



What is $c(n)$, the number of n -step SAW?

$$\begin{aligned}c(1) &= 4 \\c(2) &= c(1) \times 3 &= 12 \\c(3) &= c(2) \times 3 &= 36 \\c(4) &= c(3) \times 3 - 8 &= 100\end{aligned}$$

Not so easy! $c(n)$ is only known up to $n = 71$ [Jensen 04]

Problem: a highly non-markovian model

Some (old) conjectures/predictions

- The number of n -step SAW behaves asymptotically as follows:

$$c(n) \sim (\kappa) \mu^n n^\gamma$$

Some (old) conjectures/predictions

- The number of n -step SAW behaves asymptotically as follows:

$$c(n) \sim (\kappa) \mu^n n^\gamma$$

where

- $\gamma = 11/32$ for all 2D lattices (square, triangular, honeycomb) [Nienhuis 82]

Some (old) conjectures/predictions

- The number of n -step SAW behaves asymptotically as follows:

$$c(n) \sim (\kappa) \mu^n n^\gamma$$

where

- $\gamma = 11/32$ for all 2D lattices (square, triangular, honeycomb) [Nienhuis 82]
- $\mu = \sqrt{2 + \sqrt{2}}$ on the honeycomb lattice [Nienhuis 82]
(proved this summer [Duminil-Copin & Smirnov])


Some (old) conjectures/predictions

- The number of n -step SAW behaves asymptotically as follows:

$$c(n) \sim (\kappa) \mu^n n^\gamma$$

⇒ The probability that two n -step SAW starting from the same point do not intersect is

$$\frac{c(2n)}{c(n)^2} \sim n^{-\gamma}$$





Some (old) conjectures/predictions

- The end-to-end distance is on average

$$\mathbb{E}(D_n) \sim n^{3/4} \quad (\text{vs. } n^{1/2} \text{ for a simple random walk})$$

[Flory 49, Nienhuis 82]



Some (recent) conjectures/predictions

- **Limit process:** The scaling limit of SAW is $\text{SLE}_{8/3}$.

*(proved if the scaling limit of SAW exists and is conformally invariant
[Lawler, Schramm, Werner 02])*

This would imply

$$c(n) \sim \mu^n n^{11/32} \quad \text{and} \quad \mathbb{E}(D_n) \sim n^{3/4}$$

In 5 dimensions and above

- The critical exponents are those of the simple random walk:

$$c(n) \sim \mu^n n^0, \quad \mathbb{E}(D_n) \sim n^{1/2}.$$

- The scaling limit exists and is the d -dimensional brownian motion

[Hara-Slade 92]

Proof: a mixture of combinatorics (the lace expansion) and analysis

II. Exactly solvable models


⇒ **Design simpler classes of SAW**, that should be **natural**, as general as possible... but still tractable

- solve better and better approximations of real SAW
- develop new techniques in exact enumeration

II.0. A toy model: Partially directed walks

Definition: A walk is **partially directed** if it avoids (at least) one of the 4 steps N, S, E, W.

Example: A NEW-walk is partially directed




"Markovian with memory 1"

The self-avoidance condition is **local**.

Let $a(n)$ be the number of n -step NEW-walks.

A toy model: Partially directed walks

- Recursive description of NEW-walks:




$$a(0) = 1$$

$$a(n) = 2 + a(n-1) + 2 \sum_{k=0}^{n-2} a(k) \quad \text{for } n \geq 1$$

A toy model: Partially directed walks

- Recursive description of NEW-walks:



$$a(0) = 1$$


$$a(n) = 2 + a(n-1) + 2 \sum_{k=0}^{n-2} a(k) \quad \text{for } n \geq 1$$

- Generating function:

$$A(t) := \sum_{n \geq 0} a(n)t^n = 1 + 2 \frac{t}{1-t} + tA(t) + 2A(t) \frac{t^2}{1-t}$$

A toy model: Partially directed walks

- Recursive description of NEW-walks:



$$\begin{aligned}
 a(0) &= 1 \\
 a(n) &= 2 + a(n-1) + 2 \sum_{k=0}^{n-2} a(k) \quad \text{for } n \geq 1
 \end{aligned}$$

- Generating function:

$$A(t) := \sum_{n \geq 0} a(n)t^n = 1 + 2 \frac{t}{1-t} + tA(t) + 2A(t) \frac{t^2}{1-t}$$

$$A(t) = \frac{1+t}{1-2t-t^2} \quad \Rightarrow \quad a(n) \sim (1+\sqrt{2})^n \sim (2.41\dots)^n$$

Generating functions

Let \mathcal{A} be a set of discrete objects equipped with a size:

$$\begin{aligned} \text{size} : \mathcal{A} &\rightarrow \mathbb{N} \\ a &\mapsto |a| \end{aligned}$$

Assume that for all n ,

$$\mathcal{A}_n := \{a \in \mathcal{A} : |a| = n\} \text{ is finite.}$$

Let $a(n) = |\mathcal{A}_n|$.

The **generating function** of the objects of \mathcal{A} , counted by their size, is

$$\begin{aligned} A(t) &:= \sum_{n \geq 0} a(n)t^n \\ &= \sum_{a \in \mathcal{A}} t^{|a|}. \end{aligned}$$


Notation: $[t^n]A(t) := a(n)$

Combinatorial constructions and operations on series: A dictionary

Construction	Numbers	Generating function
Union $\mathcal{A} = \mathcal{B} \cup \mathcal{C}$	$a(n) = b(n) + c(n)$	$A(t) = B(t) + C(t)$
Product $\mathcal{A} = \mathcal{B} \times \mathcal{C}$ $ (\beta, \gamma) = \beta + \gamma $	$a(n) = b(0)c(n) + \dots + b(n)c(0)$	$A(t) = B(t) \cdot C(t)$
Sequence $\mathcal{A} = \mathfrak{S}(\mathcal{B})$ $\mathcal{A} = \{\epsilon\} \cup \mathcal{B} \cup \mathcal{B} \times \mathcal{B} \cup \dots$		$A(t) = \frac{1}{1-B(t)}$

Partially directed walks revisited

$$A(t) = 1 + 2\frac{t}{1-t} + tA(t) + 2A(t)\frac{t^2}{1-t}$$



A hierarchy of formal power series

- The formal power series $A(t)$ is **rational** if it can be written

$$A(t) = \frac{P(t)}{Q(t)}$$

where $P(t)$ and $Q(t)$ are polynomials in t .

- The formal power series $A(t)$ is **algebraic** (over $\mathbb{Q}(t)$) if it satisfies a (non-trivial) polynomial equation:

$$P(t, A(t)) = 0.$$

- The formal power series $A(t)$ is **D-finite** if it satisfies a (non-trivial) linear differential equation with polynomial coefficients:

$$P_0(t)A^{(k)}(t) + P_1(t)A^{(k-1)}(t) + \cdots + P_k(t)A(t) = 0.$$

$$\text{Rat} \subset \text{Alg} \subset \text{D-finite}$$

Some charms of this hierarchy...

- Closure properties ($+$, \times , derivatives, composition...)
- The series are “easy” to handle (partial fraction decomposition, Puiseux expansions, elimination, resultants, Gröbner bases...)
- A differential (resp. algebraic) equation can be guessed from the first coefficients (GFUN)
- The coefficients can be computed in a linear number of operations.
- (Almost) automatic asymptotics of the coefficients.
- May give insight on the (hidden) structure of the objects – or prove that they are inherently complex

What to do with a generating function?

- Extract the n th coefficient $a(n)$ (when nice...)
- The **asymptotic behaviour** of $a(n)$ can often be derived from the behaviour of $A(t)$ (seen as a function of a complex variable) in the neighborhood of its dominant singularities.

Example: $\limsup a(n)^{1/n} = \mu \iff A(t)$ has radius $1/\mu$

Transfer theorems: under certain hypotheses, if $A(t)$ has a unique dominant singularity at $1/\mu$,

$$A(t) \sim \frac{1}{(1 - \mu t)^{1+\alpha}} \implies a(n) \sim \frac{1}{\Gamma(\alpha + 1)} \mu^n n^\alpha$$

Analytic combinatorics [Flajolet-Sedgewick 09]

Example: $A(t) = \frac{1+t}{1-2t-t^2}$ has a simple pole ($\alpha = 0$) at $t_c = \sqrt{2} - 1$
 $\implies a(n) \sim \kappa(\sqrt{2} + 1)^n n^0$

Multivariate generating functions

- Enumeration according to the size (main parameter) and another parameter:

$$A(t, x) = \sum_{a \in \mathcal{A}} t^{|a|} x^{p(a)}$$

- Then

$$[t^n] \frac{\partial^i A}{\partial x^i}(t, x) \Big|_{x=1} = \sum_{a: |a|=n} p(a)(p(a) - 1) \cdots (p(a) - i + 1)$$

$$\Rightarrow \frac{[t^n] \frac{\partial^i A}{\partial x^i} A(t, 1)}{[t^n] A(t, 1)} = \mathbb{E}_n(p(p - 1) \cdots (p - i + 1))$$

is the i th **factorial moment** of $p(a)$, when a is taken uniformly among objects of size n .

- Asymptotic behaviour of the coefficients \Rightarrow Asymptotic behaviour of the moments

Analytic combinatorics [Flajolet-Sedgewick 09]

Multivariate generating functions

- Enumeration according to the size (main parameter) and another parameter:

$$A(t, x) = \sum_{a \in \mathcal{A}} t^{|a|} x^{p(a)}$$

- Then

$$\frac{[t^n]A(t, x)}{[t^n]A(t, 1)} = \frac{\sum_{a:|a|=n} x^{p(a)}}{\sum_{a:|a|=n} 1}$$

is the probability generating function of $p(a)$, when a is taken uniformly among objects of size n .

- Combined with:
 - asymptotic behaviour of $[t^n]A(t, x)$ (for $x \in \mathbb{R}$)
 - continuity theorems for (Laplace or Fourier) transforms, this often yields a limit law for the additional parameter p .

Analytic combinatorics [Flajolet-Sedgewick 09]

Example: the number of North steps in a partially directed walk

- The bivariate generating function is

$$A(t, x) = \sum_{\omega} t^{|\omega|} x^{N(\omega)} = \frac{1+t}{1-t-tx(1+t)}$$

- For x in a neighborhood of 1, there holds uniformly

$$\frac{[t^n]A(t, x)}{[t^n]A(t, 1)} \sim \kappa(x) \left(\frac{1+x+\sqrt{1+6x+x^2}}{2(1+\sqrt{2})} \right)^n (1+O(1/n))$$

- By the **quasi-power Theorem**, the number N_n of North steps satisfies

$$\mathbb{E}(N_n) \sim m n, \quad \mathbb{V}(N_n) \sim s^2 n,$$

for some $m, s^2 > 0$, and the normalized random variable $\frac{N_n - m n}{s \sqrt{n}}$ converges in law to a standard normal distribution.


+ Similar result for the abscissa of the endpoint (with mean 0).

Random partially directed walks


- Asymptotic properties: coordinates of the endpoint

$$\mathbb{E}(X_n) = 0, \quad \mathbb{E}(X_n^2) \sim n, \quad \mathbb{E}(Y_n) \sim n$$

- Random NEW-walks:



Scaled by n (– and |)



Scaled by \sqrt{n} (–) and n (|)


II.1. Weakly directed walks

(joint work with Axel Bacher)

Irreducible bridges

Def. A bridge is **irreducible** if it is not the concatenation of two bridges.


Observation: A bridge is a sequence of irreducible bridges



Weakly directed bridges

Definition: a bridge is **weakly directed** if each of its irreducible bridges avoids at least one of the steps N, S, E, W.

This means that each irreducible bridge is a NES- or a NWS-walk.



⇒ Count NES- (irreducible) bridges

Enumeration of NES-bridges


Proposition

- The generating function of NES-bridges of height $k+1$ is

$$B^{(k+1)}(t) = \sum_n b_n^{(k+1)} t^n = \frac{t^{k+1}}{G_k(t)},$$

where $G_{-1} = 1$, $G_0 = 1 - t$, and for $k \geq 0$,

$$G_{k+1} = (1 - t + t^2 + t^3)G_k - t^2G_{k-1}.$$



Enumeration of NES-bridges

Proposition

- The generating function of NES-bridges of height $k+1$ is

$$B^{(k+1)}(t) = \sum_n b_n^{(k+1)} t^n = \frac{t^{k+1}}{G_k(t)},$$


where $G_{-1} = 1$, $G_0 = 1 - t$, and for $k \geq 0$,

$$G_{k+1} = (1 - t + t^2 + t^3)G_k - t^2G_{k-1}.$$


- The generating function of NES-excursions of height at most k is

$$E^{(k)}(t) = \frac{1}{t} \left(\frac{G_{k-1}}{G_k} - 1 \right).$$

Excursion: $y_0 = 0 = y_n$ and $y_i \geq 0$ for $1 \leq i \leq n$.



Enumeration of NES-bridges



- Bridges of height $k + 1$:

$$B^{(k+1)} = tB^{(k)} + E^{(k)}t^2B^{(k)}$$

- Excursions of height at most k

$$E^{(k)} = 1 + tE^{(k)} + t^2(E^{(k-1)} - 1) + t^3(E^{(k-1)} - 1)E^{(k)}$$

- Initial conditions: $E^{(-1)} = 1$, $B^{(1)} = t/(1 - t)$.

Enumeration of NES-bridges

Proposition

- The generating function of NES-bridges of height $k+1$ is

$$B^{(k+1)}(t) = \sum_n b_n^{(k+1)} t^n = \frac{t^{k+1}}{G_k(t)},$$


where $G_{-1} = 1$, $G_0 = 1 - t$, and for $k \geq 0$,

$$G_{k+1} = (1 - t + t^2 + t^3)G_k - t^2G_{k-1}.$$

- The generating function of NES-excursions of height at most k is

$$E^{(k)}(t) = \frac{1}{t} \left(\frac{G_{k-1}}{G_k} - 1 \right).$$

Excursion: $y_0 = 0 = y_n$ and $y_i \geq 0$ for $1 \leq i \leq n$.



Enumeration of weakly directed bridges

- GF of NES-bridges:

$$B(t) = \sum_{k \geq 0} \frac{t^{k+1}}{G_k}$$

Enumeration of weakly directed bridges

- GF of NES-bridges:

$$B(t) = \sum_{k \geq 0} \frac{t^{k+1}}{G_k}$$

- GF of irreducible NES-bridges:

$$B(t) = \frac{I(t)}{1 - I(t)} \Rightarrow I(t) = \frac{B(t)}{1 + B(t)}$$

Enumeration of weakly directed bridges

- GF of NES-bridges:

$$B(t) = \sum_{k \geq 0} \frac{t^{k+1}}{G_k}$$

- GF of irreducible NES-bridges:

$$B(t) = \frac{I(t)}{1 - I(t)} \Rightarrow I(t) = \frac{B(t)}{1 + B(t)}$$

- GF of weakly directed bridges (sequences of irreducible NES- or NWS-bridges):

$$W(t) = \frac{1}{1 - (2I(t) - t)} = \frac{1}{1 - \left(\frac{2B(t)}{1+B(t)} - t\right)}$$

with $G_{-1} = 1$, $G_0 = 1 - t$, and for $k \geq 0$,

$$G_{k+1} = (1 - t + t^2 + t^3)G_k - t^2G_{k-1}.$$


Asymptotic results and nature of the generating functions

$$B(t) = \sum_{k \geq 0} \frac{t^{k+1}}{G_k}, \quad W(t) = \frac{1}{1 - \left(\frac{2B(t)}{1+B(t)} - t \right)}$$

with $G_{-1} = 1$, $G_0 = 1 - t$, and for $k \geq 0$,

$$G_{k+1} = (1 - t + t^2 + t^3)G_k - t^2G_{k-1}.$$

The zeroes of G_k (here, $k = 20$):




Asymptotic results and nature of the generating functions

$$B(t) = \sum_{k \geq 0} \frac{t^{k+1}}{G_k}, \quad W(t) = \frac{1}{1 - \left(\frac{2B(t)}{1+B(t)} - t \right)}$$

- The series $B(t)$ and $W(t)$ are meromorphic in $\mathbb{C} \setminus \mathcal{E}$, where \mathcal{E} consists of the two real intervals $[-\sqrt{2} - 1, -1]$ and $[\sqrt{2} - 1, 1]$, and of the curve

$$\mathcal{E}_0 = \left\{ x + iy : x \geq 0, y^2 = \frac{1 - x^2 - 2x^3}{1 + 2x} \right\}.$$

This curve is a natural boundary of B and W . These series thus have infinitely many singularities.



Asymptotic results and nature of the generating function

$$B(t) = \sum_{k \geq 0} \frac{t^{k+1}}{G_k}, \quad W(t) = \frac{1}{1 - \left(\frac{2B(t)}{1+B(t)} - t \right)}$$

- The series $B(t)$ and $W(t)$ are meromorphic in $\mathbb{C} \setminus \mathcal{E}$ where \mathcal{E} consists of the two real intervals $[-\sqrt{2} - 1, -1]$ and $[\sqrt{2} - 1, 1]$, and of the curve

$$\mathcal{E}_0 = \left\{ x + iy : x \geq 0, y^2 = \frac{1 - x^2 - 2x^3}{1 + 2x} \right\}.$$

This curve is a natural boundary of B and W . These series thus have infinitely many singularities.

- The series $B(t)$ has radius $\sqrt{2} - 1$, while $W(t)$ has a simple pole ρ of smaller modulus (for which $1 = \frac{2B(\rho)}{1+B(\rho)} - \rho$).

Asymptotic results and nature of the generating function

$$B(t) = \sum_{k \geq 0} \frac{t^{k+1}}{G_k}, \quad W(t) = \frac{1}{1 - \left(\frac{2B(t)}{1+B(t)} - t \right)}$$

- The series $B(t)$ and $W(t)$ are meromorphic in $\mathbb{C} \setminus \mathcal{E}$ where \mathcal{E} consists of the two real intervals $[-\sqrt{2} - 1, -1]$ and $[\sqrt{2} - 1, 1]$, and of the curve

$$\mathcal{E}_0 = \left\{ x + iy : x \geq 0, y^2 = \frac{1 - x^2 - 2x^3}{1 + 2x} \right\}.$$

This curve is a natural boundary of B and W . These series thus have infinitely many singularities.

- The series $B(t)$ has radius $\sqrt{2} - 1$, while $W(t)$ has a simple pole ρ of smaller modulus (for which $1 = \frac{2B(\rho)}{1+B(\rho)} - \rho$).
- The number $w(n)$ of weakly directed bridges of length n satisfies

$$w(n) \sim \mu^n,$$

with $\mu \simeq 2.54$ (the current record).

The number of irreducible bridges

- The generating function of weakly directed bridges, counted by the length and the number of irreducible bridges, is

$$W(t, x) = \frac{1}{1 - x \left(\frac{2B(t)}{1+B(t)} - t \right)}$$

- Let N_n denote the number N_n of irreducible bridges in a random weakly directed bridge of length n . Then


$$\mathbb{E}(N_n) \sim m n, \quad \mathbb{V}(N_n) \sim s^2 n,$$

where

$$m \simeq 0.318 \quad \text{and} \quad s^2 \simeq 0.7,$$

and the random variable $\frac{N_n - m n}{s \sqrt{n}}$ converges in law to a standard normal distribution. In particular, the average end-to-end distance, being bounded from below by $\mathbb{E}(N_n)$, grows linearly with n .

A random weakly directed bridge



II. 2. Prudent self-avoiding walks

Self-directed walks [Turban-Debierre 86]

Exterior walks [**Préa 97**]

Outwardly directed SAW [Santra-Seitz-Klein 01]

Prudent walks [**Duchi 05**], [**Dethridge, Guttmann, Jensen 07**], [**mbm 08**]


Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.



Prudent self-avoiding walks


A step never points towards a vertex that has been visited before.



not prudent!


Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Prudent self-avoiding walks

A step never points towards a vertex that has been visited before.




Remark: Partially directed walks **are** prudent



Some properties of prudent walks






Some properties of prudent walks



The **box** of a prudent walk

Some properties of prudent walks


The endpoint of a prudent walk is always on the border of the box.



Each new step either *inflates* the box or walks (prudently) *along the border*.

Some properties of prudent walks

Each new step either **inflates** the box or walks (prudently) **along the border**.




- After an inflating step, **3** possible extensions
- Otherwise, only **2**.

⇒ Count prudent walks by looking for inflating steps

Some properties of prudent walks

Each new step either inflates the box or walks (prudently) along the border.




- After an inflating step, 3 possible extensions
- Otherwise, only 2.

⇒ Count prudent walks by looking for inflating steps

When do we create an inflating step?

Some properties of prudent walks

Each new step either **inflates** the box or walks (prudently) **along the border**.




- After an inflating step, **3** possible extensions
- Otherwise, only **2**.

⇒ Count prudent walks by looking for inflating steps

When do we create an inflating step?

Some properties of prudent walks

Each new step either *inflates* the box or walks (prudently) *along the border*.




- After an inflating step, **3** possible extensions
- Otherwise, only **2**.

⇒ Count prudent walks by looking for inflating steps

When do we create an inflating step?

More parameters



If one knows:


- the direction of the last step,
- whether it is inflating or not,
- the distances i , j and h ,

then one can decide which steps can be appended to the walk, and the new values of these parameters.

⇒ Count prudent walks by looking for inflating steps, keeping track of the distances i, j, h

Recursive construction of prudent walks

- Three more parameters
(*catalytic* parameters)



- Generating function of prudent walks ending on the top of their box:


$$\left(1 - \frac{uvwt(1-t^2)}{(u-tv)(v-tu)}\right) T(t; u, v, w) = 1 + \mathcal{T}(t; w, u) + \mathcal{T}(t; w, v) - tv \frac{\mathcal{T}(t; v, w)}{u-tv} - tu \frac{\mathcal{T}(t; u, w)}{v-tu}$$

with $\mathcal{T}(t; u, v) = tvT(t; u, tu, v)$.


- Generating function of all prudent walks, counted by the length and the half-perimeter of the box:

$$P(t; u) = 1 + 4T(t; u, u, u) - 4T(t; 0, u, u)$$


Simpler families of prudent walks [Préa 97]



3-sided



2-sided



1-sided

- The endpoint of a **3-sided walk** lies always on the top, right or left side of the box
- The endpoint of a **2-sided walk** lies always on the top or right side of the box
- The endpoint of a **1-sided walk** lies always on the top side of the box (= partially directed!)

Functional equations for prudent walks:

The more general the class, the more additional variables

(Walks ending on the top of the box)

- General prudent walks: **three** catalytic variables

$$\left(1 - \frac{uvwt(1-t^2)}{(u-tv)(v-tu)}\right) T(t; u, v, w) = 1 + \mathcal{T}(w, u) + \mathcal{T}(w, v) - tv \frac{\mathcal{T}(v, w)}{u-tv} - tu \frac{\mathcal{T}(u, w)}{v-tu}$$

with $\mathcal{T}(u, v) = tvT(t; u, tu, v)$.


- Three-sided walks: **two** catalytic variables

$$\left(1 - \frac{uvt(1-t^2)}{(u-tv)(v-tu)}\right) T(t; u, v) = 1 + \dots - \frac{t^2v}{u-tv} T(t; tv, v) - \frac{t^2u}{v-tu} T(t; u, tu)$$

- Two-sided walks: **one** catalytic variable

$$\left(1 - \frac{tu(1-t^2)}{(1-tu)(u-t)}\right) T(t; u) = \frac{1}{1-tu} + t \frac{u-2t}{u-t} T(t; t)$$

Where do these equations come from? Two-sided walks



Generating function:

$$P(t; u) = \sum_{n; i} p(n; i) t^n u^i = \sum_i P_i(t) u^i \equiv P(u)$$

- For walks ending on the top of their box: series $T(t; u) \equiv T(u)$

Two-sided walks

Take a walk ending on the top of its box: where is the last step that has moved the top or right side of the box?

- Either there was no such step:

$$\frac{1}{1 - tu}$$




- ... or an East step moved the right side:




$$t \sum_i T_i(t) t^i = tT(t; t)$$

Two-sided walks

... or a North step moved the top side:



(at least one West step)



$$\frac{t^2 u}{1 - tu} T(t; u) + t \sum_{i \geq 0} T_i(t) \sum_{k=0}^i t^k u^{i-k} = \frac{t^2 u}{1 - tu} T(t; u) + \frac{t}{u - t} (u T(t; u) - t T(t; t)).$$

Two-sided walks

- The generating function $T(t; u) \equiv T(u)$ of two-sided walks ending on the top of their box satisfies:

$$\left(1 - \frac{tu(1-t^2)}{(1-tu)(u-t)}\right) T(u) = \frac{1}{1-tu} + t \frac{u-2t}{u-t} T(t)$$

- The length generating function $P(t)$ of all two-sided walks is

$$P(t) = 2T(t; 1) - T(t; 0)$$

(inclusion-exclusion)

Two-sided walks: the kernel method

$$\left((1 - tu)(u - t) - tu(1 - t^2) \right) T(t; u) = u - t + t(u - 2t)(1 - tu)T(t; t).$$

- If $u = U(t)$ cancels $(1 - tu)(u - t) - tu(1 - t^2)$, then

$$0 = U(t) - t + t(U(t) - 2t)(1 - tU(t))T(t; t),$$

that is,

$$T(t; t) = \frac{t - U(t)}{t(U(t) - 2t)(1 - tU(t))}$$

- We know such a series $U(t)$:

$$U(t) = \frac{1 - t + t^2 + t^3 - \sqrt{(1 - t^4)(1 - 2t - t^2)}}{2t}$$

[Knuth 72], [mbm-Petkovšek 2000]

Two-sided walks

- The length generating function of 2-sided walks is


$$P(t) = \frac{1}{1 - 2t - 2t^2 + 2t^3} \left(1 + t - t^3 + t(1 - t) \sqrt{\frac{1 - t^4}{1 - 2t - t^2}} \right)$$

- Dominant singularity: a simple pole for $1 - 2t - 2t^2 + 2t^3 = 0$, that is, $t_c = 0.40303\dots$. Asymptotically,

$$p(n) \sim \kappa (2.48\dots)^n$$

Compare with 2.41... for partially directed walks.

- Another approach: factorization of walks [Duchi 05]



Functional equation for 3-sided walks

(Walks ending on the top of the box)

- Three-sided walks: **two** catalytic variables

$$\left(1 - \frac{uvt(1-t^2)}{(u-tv)(v-tu)}\right) T(t; u, v) = 1 + \dots - \frac{t^2v}{u-tv} T(t; tv, v) - \frac{t^2u}{v-tu} T(t; u, tu)$$

Three-sided walks: two catalytic variables

- Functional equation for $T(t; u, v) \equiv T(u, v)$:

$$K(u, v)T(u, v) = A(u, v) + B(u, v)\Phi(u) + B(v, u)\Phi(v)$$

for polynomials $K(u, v)$, $A(u, v)$, $B(u, v)$, with $\Phi(u) = T(u, tu)$.

Three-sided walks: two catalytic variables

- Functional equation for $T(t; u, v) \equiv T(u, v)$:

$$K(u, v)T(u, v) = A(u, v) + B(u, v)\Phi(u) + B(v, u)\Phi(v)$$

for polynomials $K(u, v)$, $A(u, v)$, $B(u, v)$, with $\Phi(u) = T(u, tu)$.

- Cancellation of the kernel: $K(u, V(u)) = 0$ for a series $V(u) \equiv V(t; u)$

$$\Phi(u) = -\frac{A(u, V(u))}{B(u, V(u))} - \frac{B(V(u), u)}{B(u, V(u))} \Phi(V(u))$$

Three-sided walks: two catalytic variables

- Functional equation for $T(t; u, v) \equiv T(u, v)$:

$$K(u, v)T(u, v) = A(u, v) + B(u, v)\Phi(u) + B(v, u)\Phi(v)$$

for polynomials $K(u, v), A(u, v), B(u, v)$, with $\Phi(u) = T(u, tu)$.

- Cancellation of the kernel: $K(u, V(u)) = 0$ for a series $V(u) \equiv V(t; u)$

$$\Phi(u) = -\frac{A(u, V(u))}{B(u, V(u))} - \frac{B(V(u), u)}{B(u, V(u))} \Phi(V(u))$$

- If it is possible to iterate (...), denote $V^{(k)} = V(V(V(\dots(u))))$ (k iterations):

$$\Phi(u) = \sum_{k \geq 0} (-1)^{k-1} \frac{B(V^{(1)}, u)B(V^{(2)}, V^{(1)}) \dots B(V^{(k)}, V^{(k-1)})A(V^{(k)}, V^{(k+1)})}{B(u, V^{(1)})B(V^{(1)}, V^{(2)}) \dots B(V^{(k-1)}, V^{(k)})B(V^{(k)}, V^{(k+1)})}$$

Sum of algebraic series — iteration of algebraic functions

Three-sided prudent walks

- Let

$$U(w) = \frac{1 - tw + t^2 + t^3w - \sqrt{(1 - t^2)(1 + t - tw + t^2w)(1 - t - tw - t^2w)}}{2t},$$

and

$$q = U(1) = \frac{1 - t + t^2 + t^3 - \sqrt{(1 - t^4)(1 - 2t - t^2)}}{2t}.$$

- The length generating function of three-sided prudent walks is:

$$P(t; 1) = \frac{1}{1 - 2t - t^2} \left(\frac{1 + 3t + tq(1 - 3t - 2t^2)}{1 - tq} + 2t^2q T(t; 1, t) \right)$$

where

$$T(t; 1, t) = \sum_{k \geq 0} (-1)^k \frac{\prod_{i=0}^{k-1} \left(\frac{t}{1-tq} - U(q^{i+1}) \right)}{\prod_{i=0}^k \left(\frac{tq}{q-t} - U(q^i) \right)} \left(1 + \frac{U(q^k) - t}{t(1 - tU(q^k))} + \frac{U(q^{k+1}) - t}{t(1 - tU(q^{k+1}))} \right)$$

Three-sided prudent walks

The generating function of three-sided prudent walks is:

$$P(t; 1) = \frac{1}{1 - 2t - t^2} \left(\frac{1 + 3t + tq(1 - 3t - 2t^2)}{1 - tq} + 2t^2 q T(t; 1, t) \right)$$

$$T(t; 1, t) = \sum_{k \geq 0} (-1)^k \frac{\prod_{i=0}^{k-1} \left(\frac{t}{1-tq} - U(q^{i+1}) \right)}{\prod_{i=0}^k \left(\frac{tq}{q-t} - U(q^i) \right)} \left(1 + \frac{U(q^k) - t}{t(1 - tU(q^k))} + \frac{U(q^{k+1}) - t}{t(1 - tU(q^{k+1}))} \right)$$

- The series $P(t; 1)$ has infinitely many poles, satisfying $\frac{tq}{q-t} = U(q^i)$ for some $i \geq 0$. Hence it is neither algebraic, nor even D-finite.
- Dominant singularity: (again) a simple pole for $1 - 2t - 2t^2 + 2t^3 = 0$. Asymptotically,

$$p(n) \sim \kappa (2.48\dots)^n$$

Two- and three-sided walks: asymptotic enumeration

- The numbers of 2-sided and 3-sided n -step prudent walks satisfy

$$p_2(n) \sim \kappa_2 \mu^n, \quad p_3(n) \sim \kappa_3 \mu^n$$

where $\mu \simeq 2.48\dots$ is such that

$$\mu^3 - 2\mu^2 - 2\mu + 2 = 0.$$

Compare with 2.41... for partially directed walks, 2.54... for weakly directed bridges, but 2.64... for general SAW.

- **Conjecture:** for general prudent walks

$$p_4(n) \sim \kappa_4 \mu^n$$


with the same value of μ as above [Dethridge, Guttmann, Jensen 07].

Two-sided walks: properties of large random walks (uniform distribution)


- The random variables X_n , Y_n and δ_n satisfy

$$\mathbb{E}(X_n) = \mathbb{E}(Y_n) \sim n \quad \mathbb{E}((X_n - Y_n)^2) \sim n, \quad \mathbb{E}(\delta_n) \sim 4.15\dots$$


(+ gaussian limit law after normalization)




Two-sided walks: random generation (uniform distribution)




500 steps



780 steps



1354 steps







3148 steps

- Recursive step-by-step construction *à la Wilf* \Rightarrow 500 steps (precomputation of $O(n^2)$ large numbers)
- Boltzmann sampling via a context-free grammar [Duchon-Flajolet-Louchard-Schaeffer 02]

$$\mathbb{E}(X_n) = \mathbb{E}(Y_n) \sim n \quad \mathbb{E}((X_n - Y_n)^2) \sim n, \quad \mathbb{E}(\delta_n) \sim 4.15 \dots$$

Three-sided prudent walks: random generation and asymptotic properties

- **Asymptotic properties:** The average width of the box is $\sim \kappa n$ (variance $\sim n$, gaussian limit law after normalization)
- **Random generation:** Recursive method à la Wilf \Rightarrow 400 steps (pre-computation of $O(n^3)$ numbers)



Four-sided (i.e. general) prudent walks

- An equation with 3 catalytic variables:

$$\left(1 - \frac{uvwt(1-t^2)}{(u-tv)(v-tu)}\right) T(u, v, w) = 1 + \mathcal{T}(w, u) + \mathcal{T}(w, v) - tv \frac{\mathcal{T}(v, w)}{u-tv} - tu \frac{\mathcal{T}(u, w)}{v-tu}$$





with $\mathcal{T}(u, v) = tvT(u, tu, v)$.

- Conjecture:

$$p_4(n) \sim \kappa_4 \mu^n$$


where $\mu \simeq 2.48$ satisfies $\mu^3 - 2\mu^2 - 2\mu + 2 = 0$.

- Random prudent walks: recursive generation, 195 steps (sic! $O(n^4)$ numbers)




Another distribution: Kinetic prudent walks


- **Kinetic model:** recursive generation with no precomputation




500 steps



1000 steps



10000 steps



20000 steps


- **Theorem:** The walk chooses uniformly one quadrant, say the NE one, and then its scaling limit is given by

$$Z(u) = \int_0^{3u/7} \left(1_{W(s) \geq 0} e_1 + 1_{W(s) < 0} e_2 \right) ds$$

where e_1, e_2 form the canonical basis of \mathbb{R}^2 and $W(s)$ is a brownian motion.
[Beffara, Friedli, Velenik 10]

A kinetic, continuous space version: The rancher's walk

At time n , the walk takes a uniform unit step in \mathbb{R}^2 , conditioned so that the new step does not intersect the convex hull of the walk.



Theorem: the end-to-end distance is linear. More precisely, there exists a constant $a > 0$ such that

$$\liminf \frac{\|\omega_n\|}{n} \geq a.$$


[Angel, Benjamini, Virág 03], [Zerner 05]

Conjectures

- Linear speed: There exists $a > 0$ such that $\frac{\|\omega_n\|}{n} \rightarrow a$ a.s.
- Angular convergence: $\frac{\omega_n}{\|\omega_n\|}$ converges a.s.

What's next? Exactly solvable models


- **Exact enumeration:** General prudent walks on the square lattice – Growth constant?
- **Uniform random generation:** better algorithms (maximal length 200 for general prudent walks...)



- **A mixture of both models:** walks formed of a sequence of prudent irreducible bridges?

III. Self-avoiding walks on the honeycomb lattice

d'après H. Duminil-Copin and S. Smirnov, ArXiv 1007.0575





Simple bounds on $c(n)$

Let $c(n)$ be the number of n -step SAW. We have

$$F_{n+1} \leq c(n) \leq 3 \cdot 2^{n-1}$$

with $F_0 = F_1 = 1$ and $F_{n+1} = F_n + F_{n-1}$ (Fibonacci numbers).


- Upper bound: at most two possible extensions at each step
- Lower bound: SAW that never move in the South direction (partially directed).



The connective constant

Clearly,

$$c(m+n) \leq c(m)c(n)$$



$\Rightarrow \lim c(n)^{1/n}$ exists and

$$\mu := \lim c(n)^{1/n} = \inf c(n)^{1/n}$$

The connective constant

$$\mu := \lim c(n)^{1/n} = \inf c(n)^{1/n}$$

Given the bounds

$$F_{n+1} \leq c(n) \leq 3 \cdot 2^{n-1},$$

there holds

$$\frac{1 + \sqrt{5}}{2} \simeq 1.62 \leq \mu \leq 2$$



Theorem [Duminil-Copin & Smirnov 10]

$$\mu = \sqrt{2 + \sqrt{2}} \simeq 1.85$$

Conjectured by Nienhuis in 1982

Sub-families of SAW

Self-avoiding polygons and self-avoiding bridges



- Same growth constant:

$$p(n)^{1/n} \rightarrow \mu \quad \text{and} \quad b(n)^{1/n} \rightarrow \mu$$

[Hammersley 61]

Generating functions

For self-avoiding walks, polygons and bridges:

$$C(t) = \sum c(n)t^n, \quad P(t) = \sum p(n)t^n, \quad B(t) = \sum b(n)t^n$$

Theorem [Duminil-Copin & Smirnov 10]

The radius of convergence of these series is

$$t_c = \frac{1}{\sqrt{2 + \sqrt{2}}}.$$


Moreover:

- $C(t)$ diverges at t_c ,
- $P(t)$ converges at t_c .

Conjectures:


$$C(t) \sim \frac{1}{(1 - t_c)^{43/32}}, \quad P(t)_{sing} \sim (1 - t_c)^{3/2}, \quad B(t) \sim \frac{1}{(1 - t_c)^{9/16}}$$

Walks starting and ending at mid-edges



A key identity

Consider the following domain $D_{h,\ell}$.




A key identity

Consider the following domain $D_{h,\ell}$.

Let $L_{h,\ell}(t)$ (resp. $R_{h,\ell}(t)$, $M_{h,\ell}(t)$) be the generating function of walks that start from a and end on the left (resp. right, middle) border of the domain $D_{h,\ell}$. These series are polynomials in t .

Then for all h and ℓ , at $t = t_c = 1/\sqrt{2 + \sqrt{2}}$,

$$\frac{\sqrt{2 - \sqrt{2}}}{2} L_{h,\ell}(t_c) + R_{h,\ell}(t_c) + \frac{1}{\sqrt{2}} M_{h,\ell}(t_c) = 1$$



An upper bound on μ

$$\frac{\sqrt{2 - \sqrt{2}}}{2} L_{h,\ell}(t_c) + R_{h,\ell}(t_c) + \frac{1}{\sqrt{2}} M_{h,\ell}(t_c) = 1$$

Consider walks in $D_{h,\ell}$ that go from a to a^+ (\simeq self-avoiding polygons). Their generating function $\tilde{P}_{h,\ell}(t)$ satisfies


$$\tilde{P}_{h,\ell}(t_c) \leq L_{h,\ell}(t_c) \leq \frac{2}{\sqrt{2 - \sqrt{2}}}$$

But

$$\tilde{P}_{h,\ell}(t_c) \rightarrow \tilde{P}(t_c) \geq t_c^2 P(t_c)$$

as $h, \ell \rightarrow \infty$. This implies that $P(t_c)$ is finite.

Hence $\mu \leq 1/t_c = \sqrt{2 + \sqrt{2}}$.



A lower bound on μ

$$\frac{\sqrt{2 - \sqrt{2}}}{2} L_{h,\ell}(t_c) + R_{h,\ell}(t_c) + \frac{1}{\sqrt{2}} M_{h,\ell}(t_c) = 1$$

As $h \rightarrow \infty$, $L_{h,\ell}(t_c)$ and $R_{h,\ell}(t_c)$ increase to limit values $L_\ell(t_c)$ and $R_\ell(t_c)$. Hence $M_{h,\ell}(t_c)$ decreases to a limit value $M_\ell(t_c)$.


- If $M_\ell(t_c) > 0$ for some ℓ , the series

$$\sum_h M_{h,\ell}(t_c)$$

diverges. As it counts certain SAW, this implies that $\mu \geq 1/t_c$.

- If $M_\ell(t_c) = 0$ for all ℓ , another argument shows that $\sum_\ell R_\ell(t_c)$ (the generating function of bridges) diverges.

In both cases, $\mu \geq 1/t_c = \sqrt{2 + \sqrt{2}}$.



A lower bound on μ (continued)

$$\alpha L_{h,\ell}(t_c) + R_{h,\ell}(t_c) + \frac{1}{\sqrt{2}} M_{h,\ell}(t_c) = 1$$

- If $M_\ell(t_c) = 0$ for all ℓ , then

$$\alpha L_\ell(t_c) + R_\ell(t_c) = 1$$

$$\Rightarrow \alpha (L_{\ell+1}(t_c) - L_\ell(t_c)) = R_\ell(t_c) - R_{\ell+1}(t_c)$$


$$\Rightarrow 2\alpha t_c R_\ell R_{\ell+1} \geq R_\ell(t_c) - R_{\ell+1}(t_c)$$

$$\Rightarrow 2\alpha t_c \geq \frac{1}{R_{\ell+1}(t_c)} - \frac{1}{R_\ell(t_c)}$$

$$\Rightarrow 2\alpha t_c + \frac{1}{R_1} \geq \frac{1}{R_\ell}$$

$$\Rightarrow R_\ell \geq \frac{1}{2\alpha t_c + \beta},$$

so that $\sum_\ell R_\ell(t_c)$ (the generating function of bridges) diverges.



A key identity

$$\frac{\sqrt{2 - \sqrt{2}}}{2} L_{h,\ell}(t_c) + R_{h,\ell}(t_c) + \frac{1}{\sqrt{2}} M_{h,\ell}(t_c) = 1$$

Where does it come from?


From a local identity that is re-summed over all vertices of the domain.

A local identity


Let D be a finite, simply connected domain, and a a point on the boundary of D . For p a point in the domain, let

$$F(p) \equiv F(t, \alpha; p) = \sum_{\omega: a \rightsquigarrow p} t^{|\omega|} e^{i\alpha W(\omega)},$$


where $|\omega|$ is the length of ω , and $W(\omega)$ its **winding number**.



$$W(\omega) = 0$$



$$W(\omega) = -2\pi$$




$$W(\omega) = -\pi$$

A local identity


Let D be a finite, simply connected domain, and a a point on the boundary of D . For p a point in the domain, let

$$F(p) \equiv F(t, \alpha; p) = \sum_{\omega: a \rightsquigarrow p} t^{|\omega|} e^{i\alpha W(\omega)},$$


where $|\omega|$ is the length of ω , and $W(\omega)$ its **winding number**.



$$W(\omega) = 0$$



$$W(\omega) = -2\pi$$



$$W(\omega) = -\pi$$

Rem. W is additive:

$$W(\omega) = \frac{\pi}{3} (\text{left turns} - \text{right turns}).$$



A local identity

Let


$$F(p) \equiv F(t, \alpha; p) = \sum_{\omega: a \rightsquigarrow p \text{ in } D} t^{|\omega|} e^{i\alpha W(\omega)},$$

If p , q and r are the 3 mid-edges around a vertex v of the honeycomb lattice, taken in counterclockwise order, then, for $t = t_c$ and $\alpha = -5/8$,

$$F(p) + jF(q) + j^2F(r) = 0,$$

where $j = e^{2i\pi/3}$, or, more symmetrically,


$$(p - v)F(p) + (q - v)F(q) + (r - v)F(r) = 0.$$




A local identity

Proof: Group walks three by three or two by two as follows

- Walks that avoid v + walks in which v is the last visited vertex:




- Walks that visit v before their last vertex:



The contribution of all walks in a group is zero.

A local identity


- Walks that avoid v + walks in which v is the last visited vertex:



$$1 + jt_c e^{-i\alpha\pi/3} + j^2 t_c e^{i\alpha\pi/3} = 0$$


A local identity

- Walks that avoid v + walks in which v is the last visited vertex:



$$1 + jt_c e^{-i\alpha\pi/3} + j^2 t_c e^{i\alpha\pi/3} = 0$$

- Walks that visit v before their last vertex:



$$e^{-i\alpha\pi/3} e^{5i\alpha\pi/3} + j^2 e^{i\alpha\pi/3} e^{-5i\alpha\pi/3} = 2j \cos(\alpha\pi/3 - 2\pi/3) = 0$$


Proof of the key identity

Sum the local identity

$$(p - v)F(p) + (q - v)F(q) + (r - v)F(r) = 0$$

over all vertices v of the domain $D_{h,\ell}$.

- The inner mid-edges do not contribute.
- The winding number of walks ending on the boundary is known.



Proof of the key identity

Sum the local identity

$$(p - v)F(p) + (q - v)F(q) + (r - v)F(r) = 0$$

over all vertices v of the domain $D_{h,\ell}$.

- The inner mid-edges do not contribute.
- The domain has an up-down symmetry.
- The winding number of walks ending on the boundary is known.

This gives:

$$\frac{\sqrt{2 - \sqrt{2}}}{2} L_{h,\ell}(tc) + R_{h,\ell}(tc) + \frac{1}{\sqrt{2}} M_{h,\ell}(tc) = 1,$$

where $L_{h,\ell}(t)$ (resp. $R_{h,\ell}(t)$, $M_{h,\ell}(t)$) are the generating function of walks that start from a and end on the left (resp. right, middle) border of the domain $D_{h,\ell}$.

