

# The expected number of inversions after $n$ adjacent transpositions

Mireille Bousquet-Mélou  
CNRS, LaBRI, Bordeaux, France

ArXiv 0909:0103


Aurora  
Borealis  
Combinatorics



## A Markov chain

- Consider the symmetric group  $\mathfrak{S}_{d+1}$  on the elements  $\{0, 1, \dots, d\}$ .
- Start from the identity permutation  $\pi^{(0)} = 012 \dots d$ .
- Apply an **adjacent transposition**, taken uniformly at random (probability  $1/d$  for each).
- Repeat.

**Example:**  $d = 2$




$$\pi^{(0)} = 012$$

## A Markov chain

- Consider the symmetric group  $\mathfrak{S}_{d+1}$  on the elements  $\{0, 1, \dots, d\}$ .
- Start from the identity permutation  $\pi^{(0)} = 012 \dots d$ .
- Apply an **adjacent transposition**, taken uniformly at random (probability  $1/d$  for each).
- Repeat.

**Example:**  $d = 2$




$$\pi^{(1)} = 102$$

## A Markov chain

- Consider the symmetric group  $\mathfrak{S}_{d+1}$  on the elements  $\{0, 1, \dots, d\}$ .
- Start from the identity permutation  $\pi^{(0)} = 012 \dots d$ .
- Apply an **adjacent transposition**, taken uniformly at random (probability  $1/d$  for each).
- Repeat.

**Example:**  $d = 2$




$$\pi^{(2)} = 120$$

## A Markov chain

- Consider the symmetric group  $\mathfrak{S}_{d+1}$  on the elements  $\{0, 1, \dots, d\}$ .
- Start from the identity permutation  $\pi^{(0)} = 012 \dots d$ .
- Apply an **adjacent transposition**, taken uniformly at random (probability  $1/d$  for each).
- Repeat.

Example:  $d = 2$




$$\pi^{(3)} = 102$$

## A Markov chain

- Consider the symmetric group  $\mathfrak{S}_{d+1}$  on the elements  $\{0, 1, \dots, d\}$ .
- Start from the identity permutation  $\pi^{(0)} = 012 \dots d$ .
- Apply an **adjacent transposition**, taken uniformly at random (probability  $1/d$  for each).
- Repeat.

**Example:**  $d = 2$




$$\pi^{(4)} = 012$$

## A Markov chain

- Consider the symmetric group  $\mathfrak{S}_{d+1}$  on the elements  $\{0, 1, \dots, d\}$ .
- Start from the identity permutation  $\pi^{(0)} = 012 \dots d$ .
- Apply an **adjacent transposition**, taken uniformly at random (probability  $1/d$  for each).
- Repeat.

**Example:**  $d = 2$




$$\pi^{(5)} = 012$$




## Periodicity

This chain,  $\pi^{(0)}, \pi^{(1)}, \pi^{(2)}, \dots$  is **periodic** of period 2: it takes an even number of steps to return to a point.



## Aperiodic variants


This chain,  $\pi^{(0)}, \pi^{(1)}, \pi^{(2)}, \dots$  is **periodic** of period 2: it takes an even number of steps to return to a point.



- Either: do nothing with probability  $1/(d+1)$ , and otherwise apply an adjacent transposition chosen uniformly

## Aperiodic variants


This chain,  $\pi^{(0)}, \pi^{(1)}, \pi^{(2)}, \dots$  is **periodic** of period 2: it takes an even number of steps to return to a point.



- Either: do nothing with probability  $1/(d+1)$ , and otherwise apply an adjacent transposition chosen uniformly
- Or: consider the chains  $(\pi^{(2n)})_n$  and  $(\pi^{(2n+1)})_n$

## Aperiodic variants

This chain,  $\pi^{(0)}, \pi^{(1)}, \pi^{(2)}, \dots$  is **periodic** of period 2: it takes an even number of steps to return to a point.



- Either: do nothing with probability  $1/(d+1)$ , and otherwise apply an adjacent transposition chosen uniformly
- Or: consider the chains  $(\pi^{(2n)})_n$  and  $(\pi^{(2n+1)})_n$

These three chains are aperiodic, irreducible and symmetric, and thus converge to the uniform distribution on their respective state spaces.

## Motivations

- 1980 → 2010: Random walks in finite groups (Aldous, Diaconis, Letac, Saloff-Coste, Wilson...)

Tools: coupling techniques, representation theory...

## Motivations

- 1980 → 2010: Random walks in finite groups (Aldous, Diaconis, Letac, Saloff-Coste, Wilson...)

Tools: coupling techniques, representation theory...

- More recently: Computational biology (N. Beresticky, Durrett, Eriksen, Hultman, H. Eriksson, K. Eriksson, Sjöstrand...)


A transposition: a gene mutation

## What do we ask? What do we expect?


- **Mixing time:** How much time does it take to “reach” the uniform distribution?

The **total variation distance** between the distribution at time  $n$  and the uniform distribution on  $\mathfrak{S}_{d+1}$ :


$d = 3$



$d = 4$



$d$  large




## What do we ask? What do we expect?


- **Mixing time:** How much time does it take to “reach” the uniform distribution?

The **total variation distance** between the distribution at time  $n$  and the uniform distribution on  $\mathfrak{S}_{d+1}$ :


$d = 3$



$d = 4$



$d$  large





[Aldous 83, Diaconis & Saloff-Coste 93, Wilson 04]



## What do we ask? What do we expect?

- **Focus on observables**, for instance the inversion number  $\mathcal{I}_{d,n} = \text{inv}(\pi^{(n)})$ .

The expected value of the inversion number,  $I_{d,n} := \mathbb{E}(\mathcal{I}_{d,n})$ :




$\implies$  Estimate the number of transpositions (mutations) that have occurred, and hence the **evolutionary distance** between species.

Of particular interest: what happens **before mixing**.

## What do we ask? What do we expect?

Let  $Q = (Q_{\sigma,\tau})$  be the transition matrix of the chain:


$$Q = \begin{pmatrix} 0 & 1/2 & 1/2 & 0 & 0 & 0 \\ 1/2 & 0 & 0 & 0 & 1/2 & 0 \\ 1/2 & 0 & 0 & 1/2 & 0 & 0 \\ 0 & 0 & 1/2 & 0 & 0 & 1/2 \\ 0 & 0 & 0 & 1/2 & 1/2 & 0 \end{pmatrix}$$



## What do we ask? What do we expect?

Let  $Q = (Q_{\sigma,\tau})$  be the transition matrix of the chain:

$$Q = \begin{pmatrix} 0 & 1/2 & 1/2 & 0 & 0 & 0 \\ 1/2 & 0 & 0 & 0 & 1/2 & 0 \\ 1/2 & 0 & 0 & 1/2 & 0 & 0 \\ 0 & 0 & 1/2 & 0 & 0 & 1/2 \\ 0 & 0 & 0 & 1/2 & 1/2 & 0 \end{pmatrix}$$




- For all  $\sigma \in \mathfrak{S}_{d+1}$ , let

$$G_\sigma(t) := \sum_{n \geq 0} \mathbb{P}(\pi^{(n)} = \sigma) t^n$$

## What do we ask? What do we expect?

Let  $Q = (Q_{\sigma,\tau})$  be the transition matrix of the chain:

$$Q = \begin{pmatrix} 0 & 1/2 & 1/2 & 0 & 0 & 0 \\ 1/2 & 0 & 0 & 0 & 1/2 & 0 \\ 1/2 & 0 & 0 & 1/2 & 0 & 0 \\ 0 & 0 & 1/2 & 0 & 0 & 1/2 \\ 0 & 0 & 0 & 1/2 & 1/2 & 0 \end{pmatrix}$$




- For all  $\sigma \in \mathfrak{S}_{d+1}$ ,

$$G_{\sigma}(t) := \sum_{n \geq 0} \mathbb{P}(\pi^{(n)} = \sigma) t^n = \sum_{n \geq 0} Q_{\text{id},\sigma}^n t^n$$

## What do we ask? What do we expect?

Let  $Q = (Q_{\sigma,\tau})$  be the transition matrix of the chain:

$$Q = \begin{pmatrix} 0 & 1/2 & 1/2 & 0 & 0 & 0 \\ 1/2 & 0 & 0 & 0 & 1/2 & 0 \\ 1/2 & 0 & 0 & 1/2 & 0 & 0 \\ 0 & 0 & 1/2 & 0 & 0 & 1/2 \\ 0 & 0 & 0 & 1/2 & 1/2 & 0 \end{pmatrix}$$



- For all  $\sigma \in \mathfrak{S}_{d+1}$ ,


$$G_{\sigma}(t) := \sum_{n \geq 0} \mathbb{P}(\pi^{(n)} = \sigma) t^n = \sum_{n \geq 0} Q_{\text{id},\sigma}^n t^n = \left( (1 - tQ)^{-1} \right)_{\text{id},\sigma}$$

is a **rational series** in  $t$ .

## What do we ask? What do we expect?

Let  $Q = (Q_{\sigma,\tau})$  be the transition matrix of the chain:

$$Q = \begin{pmatrix} 0 & 1/2 & 1/2 & 0 & 0 & 0 \\ 1/2 & 0 & 0 & 0 & 1/2 & 0 \\ 1/2 & 0 & 0 & 1/2 & 0 & 0 \\ 0 & 0 & 1/2 & 0 & 0 & 1/2 \\ 0 & 0 & 0 & 1/2 & 1/2 & 0 \end{pmatrix}$$



- For all  $\sigma \in \mathfrak{S}_{d+1}$ ,

$$G_{\sigma}(t) := \sum_{n \geq 0} \mathbb{P}(\pi^{(n)} = \sigma) t^n = \sum_{n \geq 0} Q_{\text{id},\sigma}^n t^n = \left( (1 - tQ)^{-1} \right)_{\text{id},\sigma}$$

is a **rational series** in  $t$ .

- The GF of the expected inversion number

$$\sum_{n \geq 0} I_{d,n} t^n = \sum_{n \geq 0} \left( \sum_{\sigma \in \mathfrak{S}_{d+1}} \text{inv}(\sigma) \mathbb{P}(\pi^{(n)} = \sigma) \right) t^n = \sum_{\sigma \in \mathfrak{S}_{d+1}} \text{inv}(\sigma) G_{\sigma}(t)$$

is rational as well.

## A formula for the expected inversion number

**Theorem:** The expected number of inversions after  $n$  adjacent transpositions in  $\mathfrak{S}_{d+1}$  is

$$I_{d,n} = \frac{d(d+1)}{4} - \frac{1}{8(d+1)^2} \sum_{k,j=0}^d \frac{(c_j + c_k)^2}{s_j^2 s_k^2} x_{jk}^n,$$

where

$$c_k = \cos \alpha_k, \quad s_k = \sin \alpha_k, \quad \alpha_k = \frac{(2k+1)\pi}{2d+2},$$

and

$$x_{jk} = 1 - \frac{4}{d}(1 - c_j c_k).$$

### Remarks

- The series  $\sum_n I_{d,n} t^n$  is rational.
- For  $d$  large enough ( $d \geq 8$ ),  $I_{d,n}$  increases, as  $n$  grows, to  $\frac{d(d+1)}{4}$ , which is the average inversion number of a random permutation in  $\mathfrak{S}_{d+1}$ .

## Another formula for the expected inversion number [Eriksen 05]

$$I_{d,n} = \sum_{r=1}^n \frac{1}{d^r} \binom{n}{r} \sum_{s=1}^r \binom{r-1}{s-1} (-4)^{r-s} g_{s,d} h_{s,d},$$

with

$$g_{s,d} = \sum_{\ell=0}^d \sum_{k \geq 0} (-1)^k (p - 2\ell) \binom{2\lceil s/2 \rceil - 1}{\lceil s/2 \rceil + \ell + k(d+1)}$$

and

$$h_{s,d} = \sum_{j \in \mathbb{Z}} (-1)^j \binom{2\lfloor s/2 \rfloor}{\lfloor s/2 \rfloor + j(d+1)}.$$

Based on [Eriksson & Eriksson & Sjöstrand 00]

Beresticky & Durrett 08: “it is far from obvious how to extract useful asymptotic from this formula”.

Combinatorialists could not throw in the sponge!



## A formula for the expected inversion number

**Theorem:** The expected number of inversions after  $n$  adjacent transpositions in  $\mathfrak{S}_{d+1}$  is

$$I_{d,n} = \frac{d(d+1)}{4} - \frac{1}{8(d+1)^2} \sum_{k,j=0}^d \frac{(c_j + c_k)^2}{s_j^2 s_k^2} x_{jk}^n,$$

where

$$c_k = \cos \alpha_k, \quad s_k = \sin \alpha_k, \quad \alpha_k = \frac{(2k+1)\pi}{2d+2},$$

and

$$x_{jk} = 1 - \frac{4}{d}(1 - c_j c_k).$$

**Remark:** For  $d$  large enough ( $d \geq 8$ ),  $I_{d,n}$  increases, as  $n$  grows, to  $\frac{d(d+1)}{4}$ , which is the average inversion number of a random permutation in  $\mathfrak{S}_{d+1}$ .

## The expected inversion number: asymptotics

Three regimes, as  $d$  and  $n$  tend to  $\infty$

- When  $n$  is “small”, each step of the chain increases the inversion number with high probability. For example,

$$\text{inv}(\pi^{(1)}) = 1, \quad \mathbb{P}(\text{inv}(\pi^{(2)}) = 2) = 1 - \frac{1}{d}, \quad \mathbb{P}(\text{inv}(\pi^{(n)}) = n) = 1 - O\left(\frac{1}{d}\right).$$

## The expected inversion number: asymptotics

Three regimes, as  $d$  and  $n$  tend to  $\infty$

- When  $n$  is “small”, each step of the chain increases the inversion number with high probability. For example,

$$\text{inv}(\pi^{(1)}) = 1, \quad \mathbb{P}(\text{inv}(\pi^{(2)}) = 2) = 1 - \frac{1}{d}, \quad \mathbb{P}(\text{inv}(\pi^{(n)}) = n) = 1 - O\left(\frac{1}{d}\right).$$

- When  $n$  is “large”, the expected inversion number must approach its limit value  $d(d+1)/4 \sim d^2/4$ .

## The expected inversion number: asymptotics

Three regimes, as  $d$  and  $n$  tend to  $\infty$

- When  $n$  is “small”, each step of the chain increases the inversion number with high probability. For example,

$$\text{inv}(\pi^{(1)}) = 1, \quad \mathbb{P}(\text{inv}(\pi^{(2)}) = 2) = 1 - \frac{1}{d}, \quad \mathbb{P}(\text{inv}(\pi^{(n)}) = n) = 1 - O\left(\frac{1}{d}\right).$$

- When  $n$  is “large”, the expected inversion number must approach its limit value  $d(d+1)/4 \sim d^2/4$ .
- An intermediate regime?

## Small times: linear and before

- Sub-linear regime. If  $n = o(d)$ ,

$$\frac{l_{d,n}}{n} = 1 + O(n/d).$$

## Small times: linear and before

- **Sub-linear regime.** If  $n = o(d)$ ,

$$\frac{l_{d,n}}{n} = 1 + O(n/d).$$

- **Linear regime.** If  $n \sim \kappa d$ ,

$$\frac{l_{d,n}}{n} = f(\kappa) + O(1/d)$$

where

$$\begin{aligned} f(\kappa) &= \frac{1}{2\pi\kappa} \int_0^\infty \frac{1 - \exp(-8\kappa t^2/(1+t^2))}{t^2(1+t^2)} dt \\ &= \sum_{j \geq 0} (-1)^j \frac{(2j)!}{j!(j+1)!^2} (2\kappa)^j. \end{aligned}$$

The function  $f(\kappa)$  decreases from  $f(0) = 1$  to  $f(\infty) = 0$ .

## Large times: cubic and beyond

- Super-cubic regime. If  $n \gg d^3$ ,

$$\frac{l_{d,n}}{d^2} \rightarrow \frac{1}{4}.$$

## Large times: cubic and beyond

- Super-cubic regime. If  $n \gg d^3$ ,

$$\frac{I_{d,n}}{d^2} \rightarrow \frac{1}{4}.$$

- Cubic regime. If  $n \sim \kappa d^3$ ,

$$\frac{I_{d,n}}{d^2} \sim g(\kappa)$$

where

$$g(\kappa) = \frac{1}{4} - \frac{16}{\pi^4} \left( \sum_{j \geq 0} \frac{e^{-\kappa \pi^2 (2j+1)^2 / 2}}{(2j+1)^2} \right)^2.$$

The function  $g(\kappa)$  increases from  $g(0) = 0$  to  $g(\infty) = 1/4$ .



## Around the mixing time (super-cubic regime)

Assume  $n \sim \kappa d^3 \log d$ .

- If  $\kappa < 1/\pi^2$ , there exists  $\gamma > 0$  such that

$$l_{d,n} \leq \frac{d(d+1)}{4} - \Theta(d^{1+\gamma}),$$

- If  $\kappa > 1/\pi^2$ , there exists  $\gamma > 0$  such that

$$l_{d,n} = \frac{d(d+1)}{4} - O(d^{1-\gamma}).$$

- For the critical value  $\kappa = 1/\pi^2$ , the following refined estimate holds: if  $n \sim 1/\pi^2 d^3 \log d + \alpha d^3 + o(d^3)$ , then

$$l_{d,n} = \frac{d(d+1)}{4} - \frac{16d}{\pi^4} e^{-\alpha\pi^2} (1 + o(1)).$$

## The intermediate regime

- If  $d \ll n \ll d^3$ ,

$$\frac{I_{d,n}}{\sqrt{dn}} \rightarrow \sqrt{\frac{2}{\pi}}.$$

**Remark.** For a related continuous time chain, the normalized inversion number  $\mathcal{I}_{d,n}/\sqrt{dn}$  converges in probability to  $\sqrt{2/\pi}$  [Beresticky & Durrett 08]

## A formula for the expected inversion number

**Theorem:** The expected number of inversions after  $n$  adjacent transpositions in  $\mathfrak{S}_{d+1}$  is

$$I_{d,n} = \frac{d(d+1)}{4} - \frac{1}{8(d+1)^2} \sum_{k,j=0}^d \frac{(c_j + c_k)^2}{s_j^2 s_k^2} x_{jk}^n,$$

where

$$c_k = \cos \alpha_k, \quad s_k = \sin \alpha_k, \quad \alpha_k = \frac{(2k+1)\pi}{2d+2},$$

and

$$x_{jk} = 1 - \frac{4}{d}(1 - c_j c_k).$$

## Where are the inversions? [Eriksson *et al.* 00]

For  $i \leq j$ , let  $p_{i,j}^{(n)}$  be the probability that there is an inversion at time  $n$  in the positions  $i$  and  $j + 1$ :

$$p_{i,j}^{(n)} = \mathbb{P}(\pi_i^{(n)} > \pi_{j+1}^{(n)}).$$

- The expected number of inversions at time  $n$  is

$$I_{d,n} = \sum_{0 \leq i \leq j < d} p_{i,j}^{(n)}.$$

## Where are the inversions? [Eriksson *et al.* 00]

For  $i \leq j$ , let  $p_{i,j}^{(n)}$  be the probability that there is an inversion at time  $n$  in the positions  $i$  and  $j + 1$ :

$$p_{i,j}^{(n)} = \mathbb{P}(\pi_i^{(n)} > \pi_{j+1}^{(n)}).$$

- The numbers  $p_{i,j}^{(n)}$  can be described recursively by examining where were the values  $\pi_i^{(n)}$  and  $\pi_{j+1}^{(n)}$  at time  $n - 1$ .

## Where are the inversions? [Eriksson *et al.* 00]

For  $i \leq j$ , let  $p_{i,j}^{(n)}$  be the probability that there is an inversion at time  $n$  in the positions  $i$  and  $j + 1$ :

$$p_{i,j}^{(n)} = \mathbb{P}(\pi_i^{(n)} > \pi_{j+1}^{(n)}).$$

• The numbers  $p_{i,j}^{(n)}$  can be described recursively by examining where were the values  $\pi_i^{(n)}$  and  $\pi_{j+1}^{(n)}$  at time  $n - 1$ . For instance:

–  $i = j$  and the  $n$ th transposition has switched the  $i$ th and  $i + 1$ st values:

$$\left(1 - p_{i,j}^{(n-1)}\right) \frac{1}{d}$$

–  $0 \neq i = j \neq d$  and the  $n$ th transposition has not changed the values at positions  $i$  and  $i + 1$ ,

$$p_{i,j}^{(n-1)} \left(1 - \frac{3}{d}\right).$$

– etc.

## A recursion for the inversion probabilities


**Lemma.** The inversion probabilities  $p_{i,j}^{(n)}$  are characterized by:

$$p_{i,j}^{(0)} = 0 \quad \text{for} \quad 0 \leq i \leq j < d,$$

and for  $n \geq 0$ ,

$$p_{i,j}^{(n+1)} = p_{i,j}^{(n)} + \frac{1}{d} \sum_{(k,\ell) \leftrightarrow (i,j)} \left( p_{k,\ell}^{(n)} - p_{i,j}^{(n)} \right) + \frac{\delta_{i,j}}{d} \left( 1 - 2p_{i,j}^{(n)} \right),$$

where  $\delta_{i,j} = 1$  if  $i = j$  and 0 otherwise, and the neighbour relations  $\leftrightarrow$  are those of the following graph:



$\rightsquigarrow$  A (weighted) walk in a triangle.

## A functional equation for the GF of the inversion probabilities

Let  $P(t; u, v)$  be the generating function of the numbers  $p_{i,j}^{(n)}$ :

$$P(t; u, v) \equiv P(u, v) := \sum_{n \geq 0} t^n \sum_{0 \leq i \leq j < d} p_{i,j}^{(n)} u^i v^j.$$



## A functional equation for the GF of the inversion probabilities


Let  $P(t; u, v)$  be the generating function of the numbers  $p_{i,j}^{(n)}$ :

$$P(t; u, v) \equiv P(u, v) := \sum_{n \geq 0} t^n \sum_{0 \leq i \leq j < d} p_{i,j}^{(n)} u^i v^j.$$

The above recursion translates as

$$\left( 1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v}) \right) P(u, v) = \frac{t}{d} \left( \frac{1 - u^d v^d}{(1 - uv)(1 - t)} - (\bar{u} - 1)P_\ell(v) - (v - 1)v^{d-1}P_t(u) - (u + \bar{v})P_\delta(uv) \right),$$

where  $\bar{u} = 1/u$ ,  $\bar{v} = 1/v$ , and the series  $P_\ell$ ,  $P_t$  and  $P_\delta$  describe the numbers  $p_{i,j}^{(n)}$  on the boundaries of the graph:



## Back to the inversion number

We are interested in

$$l_d(t) = \sum_{n \geq 0} l_{d,n} t^n = P(t; 1, 1),$$

which, according to the functional equation, may be rewritten

$$l_d(t) = \frac{t}{(1-t)^2} - \frac{2tP_\delta(1)}{d(1-t)}.$$

◁ ◁ ◊ ▷ ▷

$$P(t; u, v) \equiv P(u, v) := \sum_{n \geq 0} t^n \sum_{0 \leq i \leq j < d} p_{i,j}^{(n)} u^i v^j.$$

$$\left( 1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v}) \right) P(u, v) = \frac{t}{d} \left( \frac{1 - u^d v^d}{(1 - uv)(1 - t)} - (\bar{u} - 1)P_\ell(v) - (v - 1)v^{d-1}P_t(u) - (u + \bar{v})P_\delta(uv) \right),$$

$$\left(1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v})\right) P(u, v) =$$

$$\frac{t}{d} \left( \frac{1 - u^d v^d}{(1 - uv)(1 - t)} - (\bar{u} - 1) P_\ell(v) - (v - 1) v^{d-1} P_t(u) - (u + \bar{v}) P_\delta(uv) \right)$$



## The ingredients of the solution

$$\left(1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v})\right) P(u, v) = \frac{t}{d} \left( \frac{1 - u^d v^d}{(1 - uv)(1 - t)} - (\bar{u} - 1) P_\ell(v) - (v - 1) v^{d-1} P_t(u) - (u + \bar{v}) P_\delta(uv) \right)$$

- Cancel the kernel  $\left(1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v})\right)$  by coupling  $u$  and  $v$

## The ingredients of the solution

$$\left(1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v})\right) P(u, v) = \frac{t}{d} \left( \frac{1 - u^d v^d}{(1 - uv)(1 - t)} - (\bar{u} - 1) P_\ell(v) - (v - 1) v^{d-1} P_t(u) - (u + \bar{v}) P_\delta(uv) \right)$$

- Cancel the kernel  $\left(1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v})\right)$  by coupling  $u$  and  $v$
- Exploit the symmetries of this kernel, which is invariant by  $(u, v) \mapsto (\bar{u}, v)$   
 $(u, v) \mapsto (u, \bar{v})$ ,  $(u, v) \mapsto (\bar{u}, \bar{v})$  (the reflection principle)

## The ingredients of the solution

$$\left(1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v})\right) P(u, v) = \frac{t}{d} \left( \frac{1 - u^d v^d}{(1 - uv)(1 - t)} - (\bar{u} - 1) P_\ell(v) - (v - 1) v^{d-1} P_t(u) - (u + \bar{v}) P_\delta(uv) \right)$$

- Cancel the kernel  $\left(1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v})\right)$  by coupling  $u$  and  $v$
- Exploit the symmetries of this kernel, which is invariant by  $(u, v) \mapsto (\bar{u}, v)$   
 $(u, v) \mapsto (u, \bar{v})$ ,  $(u, v) \mapsto (\bar{u}, \bar{v})$  (the reflection principle)
- Plus one more coupling between  $u$  and  $v$ .

## The ingredients of the solution

$$\left(1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v})\right) P(u, v) = \frac{t}{d} \left( \frac{1 - u^d v^d}{(1 - uv)(1 - t)} - (\bar{u} - 1) P_\ell(v) - (v - 1) v^{d-1} P_t(u) - (u + \bar{v}) P_\delta(uv) \right)$$

- Cancel the kernel  $\left(1 - t + \frac{t}{d}(4 - u - \bar{u} - v - \bar{v})\right)$  by coupling  $u$  and  $v$
- Exploit the symmetries of this kernel, which is invariant by  $(u, v) \mapsto (\bar{u}, v)$ ,  $(u, v) \mapsto (u, \bar{v})$ ,  $(u, v) \mapsto (\bar{u}, \bar{v})$  (the reflection principle)
- Plus one more coupling between  $u$  and  $v$ .

One obtains an **explicit expression of  $P_\delta(q)$**  at every  $q \neq -1$  such that  $q^{d+1} = -1$ , and this is enough to reconstruct the whole polynomial  $P_\delta(u)$  (and in particular,  $P_\delta(1)$ ) by interpolation.



## The final result

The generating function  $I_d(t) = \sum_{n \geq 0} I_{d,n} t^n$  is

$$I_d(t) = \frac{d(d+1)}{4(1-t)} - \frac{1}{8(d+1)^2} \sum_{k,j=0}^d \frac{(c_j + c_k)^2}{s_j^2 s_k^2} \frac{1}{1 - tx_{jk}}$$

with

$$c_k = \cos \alpha_k, \quad s_k = \sin \alpha_k, \quad \alpha_k = \frac{(2k+1)\pi}{2d+2},$$

and

$$x_{jk} = 1 - \frac{4}{d}(1 - c_j c_k).$$

## Perspectives

- Other generators (ex: all transpositions [Sjöstrand 10], transpositions  $(0, i)$ , block transpositions...)

## Perspectives

- **Other generators** (ex: all transpositions [Sjöstrand 10], transpositions  $(0, i)$ , block transpositions...)
- **Other statistics:** inversion number  $\mapsto$  measure of the “distance” between the identity and a permutation (ex: [Eriksen & Hultman 04], expected transposition distance after  $n$  transpositions)

## Perspectives

- **Other generators** (ex: all transpositions [Sjöstrand 10], transpositions  $(0, i)$ , block transpositions...)
- **Other statistics**: inversion number  $\mapsto$  measure of the “distance” between the identity and a permutation (ex: [Eriksen & Hultman 04], expected transposition distance after  $n$  transpositions)
- **Other groups**: mostly, finite irreducible Coxeter groups, with the length as the distance statistics ([Troili 02]: the case of  $I_2(d)$ ). When the generators are all reflections, see [Sjöstrand 10]!