



D-finite power series and diagonals in enumerative combinatorics



Mireille Bousquet-Mélou
CNRS, LaBRI, Université de Bordeaux, France

In this talk

I. Enumerative combinatorics and classes of generating functions

Main question:

How does one prove that a generating function
is D-finite/algebraic/rational?

II. Proving D-finiteness, algebraicity, etc. : direct solutions

[interlude] “Positive” functional equations

III. Proving D-finiteness, algebraicity, etc. : undirect solutions

IV. Disproving D-finiteness, algebraicity, etc.

I. Enumerative combinatorics and generating functions

Enumerative combinatorics

- Let \mathcal{A} be a set of discrete objects, equipped with an integer size:

$$|\cdot|: \mathcal{A} \rightarrow \mathbb{N}$$

$$o \rightarrow |o|$$

and assume that the number of objects of size n is **finite** for any n .
Denote this number by **$a(n)$** .

Enumerative combinatorics

- Let \mathcal{A} be a set of discrete objects, equipped with an integer size:

$$|\cdot|: \mathcal{A} \rightarrow \mathbb{N}$$

$$o \rightarrow |o|$$

and assume that the number of objects of size n is **finite** for any n . Denote this number by **$a(n)$** .

Def. The **ordinary generating function (GF)** of the objects of \mathcal{A} , counted by their size, is

$$A := \sum_{n \geq 0} a(n)t^n = \sum_{o \in \mathcal{A}} t^{|o|}$$

It is a formal power series **with integer coefficients**.

Enumerative combinatorics

- Let \mathcal{A} be a set of discrete objects, equipped with an integer size:

$$|\cdot|: \mathcal{A} \rightarrow \mathbb{N}$$

$$o \rightarrow |o|$$

and assume that the number of objects of size n is **finite** for any n . Denote this number by **$a(n)$** .

Def. The **ordinary generating function (GF)** of the objects of \mathcal{A} , counted by their size, is

$$A := \sum_{n \geq 0} a(n)t^n = \sum_{o \in \mathcal{A}} t^{|o|}$$

It is a formal power series **with integer coefficients**.



Multivariate enumeration: record additional (integer) statistics with additional variables x_1, \dots, x_d .

Examples

- A permutation of $\{1, 2, \dots, n\}$ is a **permutation of size n** .

Then $a(n)=n!$ and the associated generating function is:

$$A = \sum_{n \geq 0} n! t^n$$



Examples


- A permutation of $\{1, 2, \dots, n\}$ is a **permutation of size n** .

Then $a(n)=n!$ and the associated generating function is:

$$A = \sum_{n \geq 0} n! t^n$$

- **Walks** of length (size) n on the square lattice, starting at $(0,0)$:

$$A = \sum_{n \geq 0} 4^n t^n = \frac{1}{1 - 4t}$$



Examples

- A permutation of $\{1, 2, \dots, n\}$ is a **permutation of size n** .

Then $a(n)=n!$ and the associated generating function is:

$$A = \sum_{n \geq 0} n! t^n$$

- **Walks** of length (size) n on the square lattice, starting at $(0,0)$:

$$A = \sum_{n \geq 0} 4^n t^n = \frac{1}{1 - 4t}$$

Refinement: record the coordinates (i,j) of the endpoint


$$A = \sum_{\text{walk}} x^i y^j t^{\text{length}}$$

$$= \sum_{n \geq 0} \left(x + \frac{1}{x} + y + \frac{1}{y} \right)^n t^n$$

$$= \frac{1}{1 - t(x + \bar{x} + y + \bar{y})}$$

with $\bar{x}=1/x$ and $\bar{y}=1/y$.

D-finite



A hierarchy of formal power series

- Rational

$$A(t) = \frac{1-t}{1-t-t^2}$$

- Algebraic

$$1 - A(t) + tA(t)^2 = 0$$

- D-finite

$$t(1 - 16t)A''(t) + (1 - 32t)A'(t) - 4A(t) = 0$$

- D-algebraic

$$(2t + 5A(t) - 3tA'(t))A''(t) = 48t$$



Several variables: one DE per variable

A hierarchy of formal power series

- Rational

$$A(t) = \frac{1-t}{1-t-t^2}$$

- Algebraic

$$1 - A(t) + tA(t)^2 = 0$$

- D-finite

$$t(1 - 16t)A''(t) + (1 - 32t)A'(t) - 4A(t) = 0$$

- D-algebraic

$$(2t + 5A(t) - 3tA'(t))A''(t) = 48t$$



Several variables: one DE per variable



Inside the D-finite class (in one variable)


- **G-functions**: D-finite GFs with a **positive radius** of convergence
- **Diagonals (of rational series)**: for R a rational series in d variables,

$$R = \sum_{n_1, \dots, n_d \geq 0} a(n_1, \dots, n_d) x_1^{n_1} \cdots x_d^{n_d} \in \mathbb{Q}(x_1, \dots, x_d)$$

the **diagonal** of R is

$$\Delta R := \sum_{n \geq 0} a(n, n, \dots, n) t^n.$$

Recall: our GFs have integer coefficients.



Main question

How to prove that a GF is D-finite/algebraic/rational?

Main question

How to prove that a GF is D-finite/algebraic/rational?

It takes some **combinatorial understanding** of the objects, e.g.:

- a **recursive description** of the objects,
- that translates into a **recurrence relation** for the numbers $a(n)$.

Main question

How to prove that a GF is D-finite/algebraic/rational?

It takes some **combinatorial understanding** of the objects, e.g.:

- a **recursive description** of the objects,
- that translates into a **recurrence relation** for the numbers $a(n)$.

Two types of solutions:

- direct: the recurrence relation translates into a **differential/algebraic/rational** equation for the GF
- undirect: the recurrence relation translates into **another type** of functional equation for the GF

Main question

How to prove that a GF is D-finite/algebraic/rational?

It takes some **combinatorial understanding** of the objects, e.g.:

- a **recursive description** of the objects,
- that translates into a **recurrence relation** for the numbers $a(n)$.

Two types of solutions:


- direct: the recurrence relation translates into a **differential/algebraic/rational** equation for the GF
- undirect: the recurrence relation translates into **another type** of functional equation for the GF

When a GF is a G-function, is it a diagonal?

II. Direct solutions


A basic rational example

Number of **tilings** of a 2 by n board with dominoes (here, $n=12$):




A basic rational example

Number of **tilings** of a 2 by n board with dominoes (here, $n=12$):




Recursive description:




A basic rational example

Number of **tilings** of a 2 by n board with dominoes (here, n=12):



Recursive description:




Hence for $n > 0$,

$$a(n) = a(n-1) + a(n-2)$$



with $a(-1) = 0$ and $a(0) = 1$.

A basic rational example

Number of **tilings** of a 2 by n board with dominoes (here, n=12):



Recursive description:



Hence for $n > 0$,

$$a(n) = a(n-1) + a(n-2)$$


with $a(-1) = 0$ and $a(0) = 1$.

Equivalently,


$$A := \sum_{n \geq 0} a(n)t^n = 1 + tA + t^2A = \frac{1}{1 - t - t^2}.$$

A basic rational example

Number of **tilings** of a 2 by n board with dominoes (here, n=12):



Recursive description:



Hence for $n > 0$,

$$a(n) = a(n-1) + a(n-2)$$

with $a(-1) = 0$ and $a(0) = 1$.

Equivalently,



$$A := \sum_{n \geq 0} a(n)t^n = 1 + tA + t^2A = \frac{1}{1 - t - t^2}.$$

*Objects with
a linear structure*

A basic algebraic example

Binary trees with n internal nodes (here $n=6$)



Recursive description:



A basic algebraic example

Binary trees with n internal nodes (here $n=6$)

Recursive description:



Hence for $n > 0$,



$$a(n) = a(0) a(n-1) + a(1) a(n-2) + \dots + a(n-1) a(0)$$

with $a(0) = 1$.

A basic algebraic example

Binary trees with n internal nodes (here $n=6$)

Recursive description:



Hence for $n > 0$,

$$a(n) = a(0)a(n-1) + a(1)a(n-2) + \dots + a(n-1)a(0)$$

with $a(0) = 1$.



Equivalently,

$$A = \sum_{n \geq 0} a(n)t^n = 1 + \sum_{n \geq 1} t^n \left(\sum_{k=0}^{n-1} a(k)a(n-k-1) \right) = 1 + tA^2$$

A basic algebraic example

Binary trees with n internal nodes (here $n=6$)

Recursive description:



Objects with a branching structure

Hence for $n > 0$,

$$a(n) = a(0) a(n-1) + a(1) a(n-2) + \dots + a(n-1) a(0)$$


with $a(0) = 1$.

Equivalently,

$$A = \sum_{n \geq 0} a(n) t^n = 1 + \sum_{n \geq 1} t^n \left(\sum_{k=0}^{n-1} a(k) a(n-k-1) \right) = 1 + tA^2$$

A basic D-finite example

Involutions on $\{1, 2, \dots, n\}$ (here $n=7$)



A basic D-finite example

Involutions on $\{1, 2, \dots, n\}$ (here $n=7$)


Recursive description: for $n > 0$,

$$a(n) = a(n-1) + (n-1) a(n-2)$$

with $a(-1)=0$ and $a(0)=1$.

n is a fixed point

n is sent to some i in $\{1, \dots, n-1\}$



A basic D-finite example

Involutions on $\{1, 2, \dots, n\}$ (here $n=7$)

Recursive description: for $n > 0$,

$$a(n) = a(n-1) + (n-1)a(n-2)$$


with $a(-1)=0$ and $a(0)=1$.

n is a fixed point

n is sent to some i in $\{1, \dots, n-1\}$

Equivalently,

$$A = \sum_{n \geq 0} a(n)t^n = 1 + tA + t^2A'$$



A basic D-finite example

Involutions on $\{1, 2, \dots, n\}$ (here $n=7$)

Recursive description: for $n > 0$,

$$a(n) = a(n-1) + (n-1) a(n-2)$$

with $a(-1)=0$ and $a(0)=1$.

n is a fixed point


n is sent to some i in $\{1, \dots, n-1\}$

Equivalently,

$$A = \sum_{n \geq 0} a(n) t^n = 1 + tA + t^2 A'$$

Rem. This series is **NOT a G-function**, because $a(n)$ grows too

$$a(n) \sim \frac{1}{\sqrt{2}} \exp\left(-\frac{n}{2} + \sqrt{n} - \frac{1}{4}\right) n^{n/2}$$



Another basic D-finite example

All walks on the square lattice, counted by length (var. t) and final coordinates (vars. x, y):


$$R(t; x, y) = \sum_{\text{walk}} x^i y^j t^{\text{length}} = \frac{1}{1 - t(x + \bar{x} + y + \bar{y})}$$

Another basic D-finite example

All walks on the square lattice, counted by length (var. t) and final coordinates (vars. x, y):

$$R(t; x, y) = \sum_{\text{walk}} x^i y^j t^{\text{length}} = \frac{1}{1 - t(x + \bar{x} + y + \bar{y})}$$

Closed walks (i.e., ending at $(i, j) = (0, 0)$):




Another basic D-finite example

All walks on the square lattice, counted by length (var. t) and final coordinates (vars. x, y):

$$R(t; x, y) = \sum_{\text{walk}} x^i y^j t^{\text{length}} = \frac{1}{1 - t(x + \bar{x} + y + \bar{y})}$$

Closed walks (i.e., ending at $(i, j) = (0, 0)$):

$$\begin{aligned} A &= [x^0 y^0] R(t; x, y) \\ &= \Delta R(txy; x, y) \\ &= \Delta \frac{1}{1 - t(x^2 y + y + xy^2 + x)} \end{aligned}$$




Another basic D-finite example

All walks on the square lattice, counted by length (var. t) and final coordinates (vars. x, y):

$$R(t; x, y) = \sum_{\text{walk}} x^i y^j t^{\text{length}} = \frac{1}{1 - t(x + \bar{x} + y + \bar{y})}$$

Closed walks (i.e., ending at $(i, j) = (0, 0)$):

$$\begin{aligned} A &= [x^0 y^0] R(t; x, y) \\ &= \Delta R(txy; x, y) \\ &= \Delta \frac{1}{1 - t(x^2 y + y + xy^2 + x)} \end{aligned}$$



Here A is the **diagonal of a rational series**, hence a G -function, hence D-finite.

Another basic D-finite example


All walks on the square lattice, counted by length (var. t) and final coordinates (vars. x, y):

$$R(t; x, y) = \sum_{\text{walk}} x^i y^j t^{\text{length}} = \frac{1}{1 - t(x + \bar{x} + y + \bar{y})}$$

Closed walks (i.e., ending at $(i, j) = (0, 0)$):

$$\begin{aligned} A &= [x^0 y^0] R(t; x, y) \\ &= \Delta R(txy; x, y) \\ &= \Delta \frac{1}{1 - t(x^2 y + y + xy^2 + x)} \end{aligned}$$

Intuition?



Here A is the **diagonal of a rational series**, hence a G -function, hence D-finite.

Symmetric functions and D-finiteness

Closure property: the (partial) diagonals of a multivariate D-finite series are always D-finite.

Application: some classes of objects can be counted by extracting coefficients in series $A(x_1, x_2, \dots)$ that are D-finite in the power sums

$$p_k = \sum_i x_i^k$$

Examples:

- d-regular graphs on vertices $1, 2, \dots, n$, for d fixed
- square matrices of size n with non-negative coefficients summing to d on every row/column
- permutations of $\{1, 2, \dots, n\}$ with no ascending sub-sequence of length d

[Gessel 90, Goulden & Jackson ~90 ; Chyzak, Mishna & Salvy 05]

Symmetric functions and D-finiteness: example

Let $a(n)$ be the number of **3-regular graphs** on vertices $1, 2, \dots, n$.

Then

$$a(n) = [x_1^3 x_2^3 \cdots x_n^3] \prod_{1 \leq i < j} (1 + x_i x_j) = [x_1^3 x_2^3 \cdots x_n^3] G$$

where

$$G = \prod_{1 \leq i < j} (1 + x_i x_j) = \exp \left(\sum_{k \geq 1} (-1)^{k+1} \frac{p_k^2 - p_{2k}}{2k} \right)$$


is D-finite is the p_k 's, with

$$p_k = \sum_i x_i^k$$

The D-finiteness of

$$A = \sum_{n \geq 0} a(n) t^n$$

follows from closure properties.



[Interlude]

**Some rational/algebraic/
D-finite series**

...

**are more combinatorial
than others**

Reverse engineering

So far: a class of objects gives a **generating function**.

Here: some formal power series have a **natural combinatorial interpretation** as generating functions of certain objects.

Two “historical” families:

- **\mathbb{N} -rational series**
- **\mathbb{N} -algebraic series**

More recently:

- “natural” combinatorial families of **D-finite series**

\mathbb{N} -algebraic series

Example:

$$\begin{aligned} A_1 &= 1 + 2t + 4tA_1 + (12t + t^2)A_2 + t^2A_1A_2^2 \\ A_2 &= 14tA_1 + 8tA_2 + 3tA_1^3 \end{aligned}$$

Definition: a series A_1 is \mathbb{N} -algebraic if it is the first component of a k -tuple (A_1, \dots, A_k) satisfying a polynomial system of the form

$$\mathcal{A} = \begin{pmatrix} A_1 \\ \vdots \\ A_k \end{pmatrix} = \Phi(\mathcal{A}) = \begin{pmatrix} P_1(t, A_1, \dots, A_k) \\ \vdots \\ P_k(t, A_1, \dots, A_k) \end{pmatrix}$$

where

- $P_j(t, x_1, \dots, x_k)$ is a polynomial with coefficients in \mathbb{N} , for each j
- the map Φ^m is a contraction for some m .

\mathbb{N} -rational series

Example:

$$\begin{aligned} A_1 &= 1 + 2t + 4tA_1 + (12t + t^2)A_2 \\ A_2 &= 14tA_1 + 8t^2A_2 \end{aligned}$$

Definition: a series A_1 is \mathbb{N} -rational if it is the first component A_1 of a k -tuple (A_1, \dots, A_k) satisfying a polynomial system of the form

$$\mathcal{A} = \begin{pmatrix} A_1 \\ \vdots \\ A_k \end{pmatrix} = \Phi(\mathcal{A}) = \begin{pmatrix} P_1(t, A_1, \dots, A_k) \\ \vdots \\ P_k(t, A_1, \dots, A_k) \end{pmatrix}$$

where

- $P_j(t, x_1, \dots, x_k)$ is a polynomial with coefficients in \mathbb{N} , for each j , of degree at most 1 in each x_i
- the map Φ^m is a contraction for some m .

Combinatorial interpretations: \mathbb{N} -rational series

Let A_i be an \mathbb{N} -rational series. Then $A_i - a_i(0)$ counts walks on a graph.

Each series A_i will count walks from vertex 0 to vertex i on a directed finite graph.

Combinatorial interpretations: \mathbb{N} -rational series


Let A_i be an \mathbb{N} -rational series. Then $A_i - a_i(0)$ counts walks on a graph.

Each series A_i will count walks from vertex 0 to vertex i on a directed finite graph.

Example:

$$A_1 = 2tA_2 + t^3 A_1$$

$$A_2 = \dots$$



Combinatorial interpretations: \mathbb{N} -rational series


Let A_i be an \mathbb{N} -rational series. Then $A_i - a_i(0)$ counts walks on a graph.

Each series A_i will count walks from vertex 0 to vertex i on a directed finite graph.

Example:

$$A_1 = 2tA_2 + t^3 A_1$$

$$A_2 = \dots$$



Walks on finite graphs have a **linear structure**.

Combinatorial interpretations: \mathbb{N} -rational series


Let A_i be an \mathbb{N} -rational series. Then $A_i - a_i(0)$ counts walks on a graph.

Each series A_i will count walks from vertex 0 to vertex i on a directed finite graph.

Example:

$$A_1 = 2tA_2 + t^3 A_1$$

$$A_2 = \dots$$



Walks on finite graphs have a **linear structure**.

In fact, all objects known to have a rational GF have an \mathbb{N} -rational GF.

Combinatorial interpretations: \mathbb{N} -algebraic series


Let A_1 be an \mathbb{N} -algebraic series. Then $A_1 - a_1(0)$ counts by the number of nodes certain coloured trees.

Combinatorial interpretations: \mathbb{N} -algebraic series

Let A_1 be an \mathbb{N} -algebraic series. Then $A_1 - a_1(0)$ counts by the number of nodes certain coloured trees.

Example:

$$A_1 = 2t + t^2 A_1 A_2 + t A_2^3$$




Combinatorial interpretations: \mathbb{N} -algebraic series

Let A_1 be an \mathbb{N} -algebraic series. Then $A_1 - a_1(0)$ counts by the number of nodes certain **coloured trees**.

Example:

$$A_1 = 2t + t^2 A_1 A_2 + t A_2^3$$




Trees of all kinds have a **branching structure**.

Combinatorial interpretations: \mathbb{N} -algebraic series

Let A_1 be an \mathbb{N} -algebraic series. Then $A_1 - a_1(0)$ counts by the number of nodes certain **coloured trees**.

Example:

$$A_1 = 2t + t^2 A_1 A_2 + t A_2^3$$



Trees of all kinds have a **branching structure**.

However, many objects have an algebraic, but NOT \mathbb{N} -algebraic, GF.

Some criteria

Thm. Let A be a rational series with coefficients in \mathbb{N} . It is \mathbb{N} -rational if and only if all its poles of minimal modulus are of the form $\rho e^{i\theta}$, where $e^{i\theta}$ is a root of unity. [Berstel 71, Soittola 76]

Some criteria

Thm. Let A be a rational series with coefficients in \mathbb{N} . It is \mathbb{N} -rational if and only if all its poles of minimal modulus are of the form $\rho e^{i\theta}$, where $e^{i\theta}$ is a root of unity. [Berstel 71, Soittola 76]

No counterpart for \mathbb{N} -algebraic series...

Some criteria

Thm. Let A be a rational series with coefficients in \mathbb{N} . It is \mathbb{N} -rational if and only if all its poles of minimal modulus are of the form $pe^{i\theta}$, where $e^{i\theta}$ is a root of unity. [Berstel 71, Soittola 76]

No counterpart for \mathbb{N} -algebraic series...

A necessary asymptotic condition for \mathbb{N} -algebraicity: if

$$a(n) \sim \kappa \mu^n n^{\gamma-1}$$

then μ is algebraic and

$$\gamma \in \left\{ -\frac{1}{2^k}, k \geq 1 \right\} \cup \left\{ \frac{m}{2^k}, m \geq 1, k \geq 0 \right\}$$

NB. The algebraicity itself only implies that $\gamma \in \mathbb{Q} \setminus \{0, -1, -2, \dots\}$.

[Banderier-Drmotu 15]

\mathbb{N} -algebraic series vs. \mathbb{N} -rational series

Remark. \mathbb{N} -rationality and \mathbb{N} -algebraicity can be defined (via positive systems) for **multivariate series**.

Example:

$$\begin{aligned} A_1 &= 1 + 2x + 4yA_1 + (12x + y^2)A_2 + y^2A_1A_2^2 \\ A_2 &= 14yA_1 + 8xA_2 + 3xA_1^3 \end{aligned}$$

\mathbb{N} -algebraic series vs. \mathbb{N} -rational series

Remark. \mathbb{N} -rationality and \mathbb{N} -algebraicity can be defined (via positive systems) for **multivariate series**.

Known: algebraic series in one variable are the diagonals of bivariate rational series [Furstenberg 67].

\mathbb{N} -algebraic series vs. \mathbb{N} -rational series

Remark. \mathbb{N} -rationality and \mathbb{N} -algebraicity can be defined (via positive systems) for **multivariate series**.

Known: algebraic series in one variable are the diagonals of bivariate rational series [Furstenberg 67].

Questions:

- If $R(x,y)$ is \mathbb{N} -rational, is its diagonal \mathbb{N} -algebraic?

\mathbb{N} -algebraic series vs. \mathbb{N} -rational series

Remark. \mathbb{N} -rationality and \mathbb{N} -algebraicity can be defined (via positive systems) for **multivariate series**.

Known: algebraic series in one variable are the diagonals of bivariate rational series [Furstenberg 67].

Questions:

- If $R(x,y)$ is \mathbb{N} -rational, is its diagonal \mathbb{N} -algebraic?
- If $R(x_1, \dots, x_k)$ is \mathbb{N} -rational with an algebraic diagonal, is this diagonal \mathbb{N} -algebraic?

\mathbb{N} -algebraic series vs. \mathbb{N} -rational series

Remark. \mathbb{N} -rationality and \mathbb{N} -algebraicity can be defined (via positive systems) for **multivariate series**.

Known: algebraic series in one variable are the diagonals of bivariate rational series [Furstenberg 67].

Questions:

- If $R(x,y)$ is \mathbb{N} -rational, is its diagonal \mathbb{N} -algebraic?
- If $R(x_1, \dots, x_k)$ is \mathbb{N} -rational with an algebraic diagonal, is this diagonal \mathbb{N} -algebraic?
- If A is \mathbb{N} -algebraic in one variable, is it the diagonal of an \mathbb{N} -rational series?

\mathbb{N} -algebraic series vs. \mathbb{N} -rational series

Remark. \mathbb{N} -rationality and \mathbb{N} -algebraicity can be defined (via positive systems) for **multivariate series**.

Known: algebraic series in one variable are the diagonals of bivariate rational series [Furstenberg 67].

Questions:

- If $R(x,y)$ is \mathbb{N} -rational, is its diagonal \mathbb{N} -algebraic?
- If $R(x_1, \dots, x_k)$ is \mathbb{N} -rational with an algebraic diagonal, is this diagonal \mathbb{N} -algebraic?
- If A is \mathbb{N} -algebraic in one variable, is it the diagonal of an \mathbb{N} -rational series?

Conjecture [Garrabrant-Pak 14(a)] Not true for $A=1+tA^2$.

Combinatorial D-finite series?

Several attempts:

- [Kotek-Makowsky 12]: an interpretation in terms of square lattice paths for some D-finite series
- [Garrabrant-Pak 14(a)]: a combinatorial interpretation for all diagonals of \mathbb{N} -rational series $R(x_1, \dots, x_k)$ in terms of tilings of a strip with tiles of irrational sizes
- [Koechlin et al. 20+]: a combinatorial interpretation of the same class in terms of constrained walks on a graph
- [Koechlin et al. 20+]: a related combinatorial interpretation for diagonals of \mathbb{N} -algebraic series

III. Undirect solutions

A recursive description of the objects yields a recurrence relation for $a(n)$, and a functional equation for the series A , that is **not of rational/polynomial/differential type**.

Additional variables

Instead of $A=A(t)$, one considers a multivariate series $A(t; x_1, \dots, x_d)$, and proves that it satisfies some functional equation.

Examples:

- Integer partitions: a **q-equation** (or t-equation...)

$$A(t; x) = tx + \frac{1}{1 - tx} A(t; xt)$$

- Set partitions (one variable only!): a **difference equation**

$$A(t) = 1 + \frac{t}{1 - t} A\left(\frac{t}{1 - t}\right)$$

- Walks on a half-line: a **discrete differential equation**

$$A(t; x) = 1 + txA(t; x) + t \frac{A(t; x) - A(t; 0)}{x}$$

The variable x is “**catalytic**”.

A basic example: walks on a half-line

Let $a(n;k)$ be the number of n -step walks with steps $+1$ and -1 , that start at 0 , **remain non-negative**, and end at height k . Then


$$a(n;k) = a(n-1;k-1) + a(n-1;k+1)$$

with the boundary condition $a(n;k)=0$ for $k<0$. This translates into

$$A(t;x) = 1 + txA(t;x) + t \frac{A(t;x) - A(t;0)}{x},$$

for the bi-variate generating function

$$A(t;x) = \sum_{n,k \geq 0} a(n;k)t^n x^k.$$



A basic example: walks on a half-line

Let $a(n;k)$ be the number of n -step walks with steps $+1$ and -1 , that start at 0 , **remain non-negative**, and end at height k . Then

$$a(n;k) = a(n-1;k-1) + a(n-1;k+1)$$

with the boundary condition $a(n;k)=0$ for $k<0$. This translates into


$$A(t;x) = 1 + txA(t;x) + t \frac{A(t;x) - A(t;0)}{x},$$

for the bi-variate generating function

$$A(t;x) = \sum_{n,k \geq 0} a(n;k) t^n x^k.$$

In short,

$$A(x) = 1 + txA(x) + t \frac{A(x) - A(0)}{x}.$$



A basic example: walks on a half-line

Remark. In the equation

$$A(t; x) = 1 + txA(t; x) + t \frac{A(t; x) - A(t; 0)}{x},$$

we can replace the series $A(t; 0)$ by a series $B(t)$:

$$A(t; x) = 1 + txA(t; x) + t \frac{A(t; x) - B(t)}{x},$$

Taking the limit $x \rightarrow 0$ shows that $B(t)$ must be $A(t; 0)$ as soon as $A(t; x)$ belongs to $\mathbb{C}[[t, x]]$.

A basic example: walks on a half-line

Remark. In the equation

$$A(t; x) = 1 + txA(t; x) + t \frac{A(t; x) - A(t; 0)}{x},$$

we can replace the series $A(t; 0)$ by a series $B(t)$:

$$A(t; x) = 1 + txA(t; x) + t \frac{A(t; x) - B(t)}{x},$$

Taking the limit $x \rightarrow 0$ shows that $B(t)$ must be $A(t; 0)$ as soon as $A(t; x)$ belongs to $\mathbb{C}[[t, x]]$.

A single equation may define several formal power series.



More examples in one catalytic variable

- **Walks with steps +1 and -2** on a half-line:

$$A(x) = 1 + txA(x) + t \frac{A(x) - A(0) - xA'(0)}{x^2}$$

- **Planar maps** [Tutte 68]: non-linear, one catalytic variable

$$A(x) = 1 + tx^2A(x)^2 + tx \frac{xA(x) - A(1)}{x-1}$$



One catalytic variable: algebraicity

Theorem [MBM-Jehanne 06]

Let $P(A(x), B_1, B_2, \dots, B_k, t, x)$ be a polynomial equation in one catalytic variable x . Under natural (combinatorial) assumptions, the series $A(x) = A(t; x)$ and the $B_i = B_i(t)$'s are **algebraic**.

Example: planar maps

$$A(x) = 1 + tx^2 A(x)^2 + tx \frac{x A(x) - A(1)}{x - 1}$$



Algebraic series



One catalytic variable: algebraicity

Theorem [MBM-Jehanne 06]

Let $P(A(x), B_1, B_2, \dots, B_k, t, x)$ be a polynomial equation in one catalytic variable x . Under natural (combinatorial) assumptions, the series $A(x) = A(t; x)$ and the $B_i = B_i(t)$'s are **algebraic**.

Extension [Notarantonio-Yurkevich 22(a)]

The same holds for **systems** of such equations, defining series $A_1(x), \dots, A_n(x)$ and B_1, B_2, \dots, B_k .

Example: planar maps

$$A(x) = 1 + tx^2 A(x)^2 + tx \frac{x A(x) - A(1)}{x - 1}$$



Algebraic series



One catalytic variable: algebraicity

Theorem [MBM-Jehanne 06]

Let $P(A(x), B_1, B_2, \dots, B_k, t, x)$ be a polynomial equation in one catalytic variable x . Under natural (combinatorial) assumptions, the series $A(x) = A(t; x)$ and the $B_i = B_i(t)$'s are **algebraic**.

[Popescu 85, Swan 98]

Extension [Notarantonio-Yurkevich 22(a)]

The same holds for **systems** of such equations, defining series $A_1(x), \dots, A_n(x)$ and B_1, B_2, \dots, B_k .

Example: planar maps

$$A(x) = 1 + tx^2 A(x)^2 + tx \frac{x A(x) - A(1)}{x - 1}$$



Algebraic series



One catalytic variable: algebraicity

Theorem [MBM-Jehanne 06]

Let $P(A(x), B_1, B_2, \dots, B_k, t, x)$ be a polynomial equation in one catalytic variable x . Under natural (combinatorial) assumptions, the series $A(x) = A(t; x)$ and the $B_i = B_i(t)$'s are **algebraic**.

[Popescu 85, Swan 98]

Extension [Notarantonio-Yurkevich 22(a)]

The same holds for **systems** of such equations, defining series $A_1(x), \dots, A_n(x)$ and B_1, B_2, \dots, B_k .

effective

Example: planar maps

$$A(x) = 1 + tx^2 A(x)^2 + tx \frac{x A(x) - A(1)}{x - 1}$$



Algebraic series



One catalytic variable: algebraicity

Theorem [MBM-Jehanne 06]

Let $P(A(x), B_1, B_2, \dots, B_k, t, x)$ be a polynomial equation in one catalytic variable x . Under natural (combinatorial) assumptions, the series $A(x) = A(t; x)$ and the $B_i = B_i(t)$'s are **algebraic**.

[Popescu 85, Swan 98]

Extension [Notarantonio-Yurkevich 22(a)]

The same holds for **systems** of such equations, defining series $A_1(x), \dots, A_n(x)$ and B_1, B_2, \dots, B_k .

effective

Example: planar maps

$$A(x) = 1 + tx^2 A(x)^2 + tx \frac{x A(x) - A(1)}{x - 1}$$

$$\Rightarrow 7A(1)^2 t^2 + (1 - 18t) A(1) + 16t - 1 = 0$$



Algebraic series



Popescu's theorem (in $d+1$ variables)

Here $d=2$, hence 3 variables t, x, y .

Thm. If a system of equations of the form

$$\text{Pol}(A_1, \dots, A_n, B_1, \dots, B_m, C_1, \dots, C_k, t, x, y) = 0$$

has a **unique solution** such that the following **diagonal conditions** hold

$$A_i \in \mathbb{C}[[t, x, y]], \quad B_i \in \mathbb{C}[[t, x]], \quad C_i \in \mathbb{C}[[t]],$$

then each of these series is **algebraic**.

Example:

$$A(t; x, y), \quad A(t; x, 0), \quad A(t; 0, 0) \quad \cancel{A(t; 0, y)}$$

Popescu's theorem (in $d+1$ variables)

Here $d=2$, hence 3 variables t, x, y .

Thm. If a system of equations of the form

$$\text{Pol}(A_1, \dots, A_n, B_1, \dots, B_m, C_1, \dots, C_k, t, x, y) = 0$$

has a **unique solution** such that the following **diagonal conditions** hold

$$A_i \in \mathbb{C}[[t, x, y]], \quad B_i \in \mathbb{C}[[t, x]], \quad C_i \in \mathbb{C}[[t]],$$

then each of these series is **algebraic**.

Example:

$$A(t; x, y), \quad A(t; x, 0), \quad A(t; 0, 0) \quad \cancel{A(t; 0, y)}$$

Alas... such equations do not occur much in combinatorics!



Two discrete derivatives: some examples

- Square lattice walks confined to a quadrant: linear equation

$$Q(x, y) = 1 + t(x + y)Q(x, y) + t \frac{Q(x, y) - Q(x, 0)}{y} + t \frac{Q(x, y) - Q(0, y)}{x}$$

- q-Coloured planar triangulations [Tutte, 1973-1984]: non-linear

$$T(x, y) = x(q - 1) + txyT(1, y)T(x, y) + tx \frac{T(x, y) - T(x, 0)}{y} - tx^2y \frac{T(x, y) - T(1, y)}{x - 1}.$$



Two discrete derivatives: some examples

- Square lattice walks confined to a quadrant: linear equation

$$Q(x, y) = 1 + t(x + y)Q(x, y) + t \frac{Q(x, y) - Q(x, 0)}{y} + t \frac{Q(x, y) - Q(0, y)}{x}$$

- q-Coloured planar triangulations [Tutte, 1973-1984]: non-linear

$$T(x, y) = x(q - 1) + txyT(1, y)T(x, y) + tx \frac{T(x, y) - T(x, 0)}{y} - tx^2y \frac{T(x, y) - T(1, y)}{x - 1}.$$

- Three-stack sortable permutations [Defant, Elvey Price, Guttman 21]

$$P(x, y) = t(x + 1)^2(y + 1)^2 + ty(1 + x)P(x, y) + t(1 + x) \frac{P(x, y) - P(x, 0)}{y} \left((1 + y)^2 + y \frac{P(x, y) - P(0, y)}{x} \right)$$

A natural question

Solutions of discrete differential equations (with derivatives with respect to **one variable** only) are **algebraic**.

$$A(t; x) = 1 + txA(t; x) + t \frac{A(t; x) - A(t; 0)}{x}$$

Is it always true for solutions of discrete PDEs (with derivatives with respect to **several variables**)?

$$Q(x, y) = 1 + t(x + y)Q(x, y) + t \frac{Q(x, y) - Q(x, 0)}{y} + t \frac{Q(x, y) - Q(0, y)}{x}$$

A natural question

Solutions of discrete differential equations (with derivatives with respect to **one variable** only) are **algebraic**.

$$A(t; x) = 1 + txA(t; x) + t \frac{A(t; x) - A(t; 0)}{x}$$

Is it always true for solutions of discrete PDEs (with derivatives with respect to **several variables**)?

$$Q(x, y) = 1 + t(x + y)Q(x, y) + t \frac{Q(x, y) - Q(x, 0)}{y} + t \frac{Q(x, y) - Q(0, y)}{x}$$

no

A natural question

Solutions of discrete differential equations (with derivatives with respect to **one variable** only) are **algebraic**.

$$A(t; x) = 1 + txA(t; x) + t \frac{A(t; x) - A(t; 0)}{x}$$

Is it always true for solutions of discrete PDEs (with derivatives with respect to **several variables**)?

$$Q(x, y) = 1 + t(x + y)Q(x, y) + t \frac{Q(x, y) - Q(x, 0)}{y} + t \frac{Q(x, y) - Q(0, y)}{x}$$

no



Well, sometimes

The story of walks confined to the quadrant

About twenty years ago...

- Systematic study of quadrant walks

Set of steps ("model") in



- Some models are trivial, or equivalent to a half plane problem



⇒ 79 really interesting and distinct small step models [mbm-Mishna10]

The story of walks confined to the quadrant

About twenty years ago...

- Systematic study of quadrant walks


Set of steps ("model") in



- Some models are trivial, or equivalent to a half plane problem

⇒ 79 really interesting and distinct small step models [mbm-Mishna10]

- Systematic approach via a functional equation, e.g.





$$Q(x, y) = 1 + t(x + y)Q(x, y) + t \frac{Q(x, y) - Q(x, 0)}{y} + t \frac{Q(x, y) - Q(0, y)}{x}$$

The story of walks confined to the quadrant

About twenty years ago...

- Systematic study of quadrant walks


Set of steps ("model") in



- Some models are trivial, or equivalent to a half plane problem

⇒ 79 really interesting and distinct small step models [mbm-Mishna10]

- Systematic approach via a functional equation, e.g.




$$Q(x, y) = 1 + t(x + y)Q(x, y) + t \frac{Q(x, y) - Q(x, 0)}{y} + t \frac{Q(x, y) - Q(0, y)}{x}$$

Equivalently,

$$(1 - t(x + \bar{x} + y + \bar{y}))xyQ(x, y) = xy - txQ(x, 0) - tyQ(0, y)$$


Four examples



Algebraic

[Kreweras 65, Gessel 86]


$$(1 - t(\bar{x} + \bar{y} + xy))xyQ(x, y) = xy - tyQ(0, y) - txQ(x, 0)$$



D-finite transcendental


[Gessel 90]

$$(1 - t(y + \bar{x} + x\bar{y}))xyQ(x, y) = xy - tyQ(0, y) - tx^2Q(x, 0)$$



D-algebraic, but not D-finite [Bernardi, mbm, Raschel 17(a)]

$$(1 - t(x + \bar{x} + y + x\bar{y}))xyQ(x, y) = xy - tyQ(0, y) - tx^2Q(x, 0)$$




Not D-algebraic

[Dreyfus, Hardouin, Roques, Singer 17]


$$(1 - t(x\bar{y} + \bar{x} + \bar{y} + y))xyQ(x, y) = xy - tyQ(0, y) - tx(1 + x)Q(x, 0)$$

Twenty years later: classification of quadrant walks




Bernardi, Bostan, mbm, Budd, Chyzak, Dreyfus, Elvey Price, Gessel, Hardouin, Kauers, Koutschan, Kurkova, Melczer, Mishna, Pech, Raschel, Rechnitzer, Roques, Salvy, Singer, van Hoeij, Zeilberger...

Twenty years later: classification of quadrant walks



Bernardi, Bostan, mbm, Budd, Chyzak, Dreyfus, Elvey Price, Gessel, Hardouin, Kauers, Koutschan, Kurkova, Melczer, Mishna, Pech, Raschel, Rechnitzer, Roques, Salvy, Singer, van Hoeij, Zeilberger...

Twenty years later: classification of quadrant walks




Bernardi, Bostan, mbm, Budd, Chyzak, Dreyfus, Elvey Price, Gessel, Hardouin, Kauers, Koutschan, Kurkova, Melczer, Mishna, Pech, Raschel, Rechnitzer, Roques, Salvy, Singer, van Hoeij, Zeilberger...

Linear discrete PDEs and diagonals


Walks on the square lattice confined to the first quadrant:

$$(1 - t(x + \bar{x} + y + \bar{y}))xyQ(x, y) = xy - txQ(x, 0) - tyQ(0, y).$$



Linear discrete PDEs and diagonals

Walks on the square lattice confined to the first quadrant:




$$(1 - t(x + \bar{x} + y + \bar{y}))xyQ(x, y) = xy - txQ(x, 0) - tyQ(0, y).$$

By playing with the symmetries $x \rightarrow 1/x$ and $y \rightarrow 1/y$, one obtains

$$xyQ(x, y) - \bar{x}yQ(\bar{x}, y) + \bar{x}\bar{y}Q(\bar{x}, \bar{y}) - x\bar{y}Q(x, \bar{y}) = \frac{xy - \bar{x}y + \bar{x}\bar{y} - x\bar{y}}{1 - t(x + \bar{x} + y + \bar{y})}$$

Linear discrete PDEs and diagonals

Walks on the square lattice confined to the first quadrant:



$$(1 - t(x + \bar{x} + y + \bar{y}))xyQ(x, y) = xy - txQ(x, 0) - tyQ(0, y).$$

By playing with the symmetries $x \rightarrow 1/x$ and $y \rightarrow 1/y$, one obtains


$$xyQ(x, y) - \bar{x}yQ(\bar{x}, y) + \bar{x}\bar{y}Q(\bar{x}, \bar{y}) - x\bar{y}Q(x, \bar{y}) = \frac{xy - \bar{x}y + \bar{x}\bar{y} - x\bar{y}}{1 - t(x + \bar{x} + y + \bar{y})}$$

so that

$$xyQ(x, y) = [x^{>0}y^{>0}] \frac{xy - x\bar{y} + \bar{x}\bar{y} - \bar{x}y}{1 - t(x + \bar{x} + y + \bar{y})}.$$

Linear discrete PDEs and diagonals

Walks on the square lattice confined to the first quadrant:



$$(1 - t(x + \bar{x} + y + \bar{y}))xyQ(x, y) = xy - txQ(x, 0) - tyQ(0, y).$$

By playing with the symmetries $x \rightarrow 1/x$ and $y \rightarrow 1/y$, one obtains

$$xyQ(x, y) - \bar{x}yQ(\bar{x}, y) + \bar{x}\bar{y}Q(\bar{x}, \bar{y}) - x\bar{y}Q(x, \bar{y}) = \frac{xy - \bar{x}y + \bar{x}\bar{y} - x\bar{y}}{1 - t(x + \bar{x} + y + \bar{y})}$$

so that


$$xyQ(x, y) = [x^{>0}y^{>0}] \frac{xy - x\bar{y} + \bar{x}\bar{y} - \bar{x}y}{1 - t(x + \bar{x} + y + \bar{y})}.$$

Equivalently,

$$Q(u, v) = [x^0y^0] \frac{1}{(1 - u\bar{x})(1 - v\bar{y})} \frac{1 - \bar{y}^2 + \bar{x}^2\bar{y}^2 - \bar{x}^2}{1 - t(x + \bar{x} + y + \bar{y})},$$

and $Q(u, v)$ can also be written as a **diagonal**.

Twenty years later: classification of quadrant walks



Bernardi, Bostan, mbm, Budd, Chyzak, Dreyfus, Elvey Price, Gessel, Hardouin, Kauers, Koutschan, Kurkova, Melczer, Mishna, Pech, Raschel, Rechnitzer, Roques, Salvy, Singer, van Hoeij, Zeilberger...

Linear discrete PDEs and diagonals

In fact, all linear equations* in two or more catalytic variables whose solution has been proved D-finite (transcendental) have been solved in terms of **diagonals of rational functions**.

Linear discrete PDEs and diagonals

In fact, all linear equations* in two or more catalytic variables whose solution has been proved D-finite (transcendental) have been solved in terms of **diagonals of rational functions**.

- **2D walks** confined to the non-negative quadrant, with steps in $\{(1,0), (-1,0), (0,-1), (-2,1)\}$: [Bostan, mbm, Melczer 21]

$$Q(x, y) = [x^{\geq 0} y^{\geq 0}] \frac{(x^2 + 1)(x + y)(y - x)(x^2 y - 2x - y)(x^3 - x - 2y)}{x^7 y^3 (1 - t(x + \bar{x} + \bar{x}^2 y + \bar{y}))}$$

- **3D walks** confined to the non-negative orthant, with steps in $\{(-1,-1,-1), (-1,-1,1), (-1,1,0), (1,0,0)\}$: [Bostan, mbm, Kauers, Melczer 16⁷]

$$O(x, y, z) = [x^{\geq 0} y^{\geq 0} z^{\geq 0}] \frac{(x - \bar{x}y - \bar{x}\bar{y}z - \bar{x}\bar{y}\bar{z})(y - \bar{y}z - \bar{y}\bar{z})(z - \bar{z})}{x^2 y^2 z^2 (1 - t(\bar{x}\bar{y}\bar{z} + \bar{x}\bar{y}z + \bar{x}y + x))}$$

Linear discrete PDEs and diagonals

In fact, all linear equations* in two or more catalytic variables whose solution has been proved D-finite (transcendental) have been solved in terms of **diagonals of rational functions**.

- For $m=2\ell+1$, the GF of **involutions with no ascending sub-sequence** of length $m+1$ is:

$$I_m(t) = [x_1 x_2^2 \cdots x_\ell^\ell] \frac{\det(x_j^i - \bar{x}_j^i)_{1 \leq i, j \leq \ell}}{1 - t(1 + x_1 + \cdots + x_\ell + \bar{x}_1 + \cdots + \bar{x}_\ell)}$$

- The GF of **permutations with no ascending sub-sequence** of length $m+1$ is:

$$[x_1^0 \cdots x_m^0] \frac{\det(z_j^{i-j})_{1 \leq i, j \leq m}}{1 - t(\bar{z}_1 + \cdots + \bar{z}_m)} \sum_{i=0}^m \prod_{j=1}^i \frac{x_j}{1 - x_j}$$

with $z_j = x_j - x_{j-1}$.

[mbm 11]

Linear discrete PDEs and diagonals

In fact, all linear equations* in two or more catalytic variables whose solution has been proved D-finite (transcendental) have been solved in terms of **diagonals of rational functions**.

This must be taken with a grain of salt!

- Quadrant walks with steps in $\{(-1,-1), (-1,1), (-1,0), (1,0), (1,1)\}$ are D-finite transcendental, but this has only been proved by a **guess & check** computer approach [Bostan, mbm, Kauers, Melczer 16].

$$\begin{aligned} (1 - t(1 + y) (\bar{x} + \bar{x}\bar{y} + x))xyQ(x, y) \\ = xy - tQ(x, 0) - t(1 + y)^2Q(0, y) + tQ(0, 0) \end{aligned}$$

Linear discrete PDEs and diagonals

In fact, all linear equations* in two or more catalytic variables whose solution has been proved D-finite (transcendental) have been solved in terms of **diagonals of rational functions**.

This must be taken with a grain of salt!

- Quadrant walks with steps in $\{(-1,-1), (-1,1), (-1,0), (1,0), (1,1)\}$ are D-finite transcendental, but this has only been proved by a **guess & check** computer approach [Bostan, mbm, Kauers, Melczer 16].

$$\begin{aligned}(1 - t(1 + y) (\bar{x} + \bar{x}\bar{y} + x))xyQ(x, y) \\ = xy - tQ(x, 0) - t(1 + y)^2Q(0, y) + tQ(0, 0)\end{aligned}$$

- Some quadrant walks are **conjectured** D-finite, with no proof at all. E.g. with steps in $\{(1,0), (0,1), (-2,-1), (-1,-2)\}$ [Bostan, mbm, Melczer 21]

**IV. Disproving
D-finiteness,
algebraicity, etc.**

Not really a combinatorial question!

Some tools

Asymptotics. Assume that the (integer) sequence $a(n)$ satisfies

$$a(n) \sim \kappa \mu^n n^\gamma, \quad n \rightarrow \infty.$$

If the associated series $A(t)$ is **rational/algebraic/D-finite** then

- μ is algebraic
- γ belongs to \mathbb{N} , $\mathbb{Q} \setminus \{-1, -2, \dots\}$, \mathbb{Q} .

G-functions

[Katz 70, Chudnovsky² 85, André 89... Bostan, Raschel, Salvy 14]

Some tools

Asymptotics. Assume that the (integer) sequence $a(n)$ satisfies

$$a(n) \sim \kappa \mu^n n^\gamma, \quad n \rightarrow \infty.$$

If the associated series $A(t)$ is **rational/algebraic/D-finite** then

- μ is algebraic
- γ belongs to \mathbb{N} , $\mathbb{Q} \setminus \{-1, -2, \dots\}$, \mathbb{Q} .

G-functions

[Katz 70, Chudnovsky² 85, André 89... Bostan, Raschel, Salvy 14]

Singularities. A D-finite series $A(t)$ of positive radius of convergence has **finitely many singularities**.

Some tools

Asymptotics. Assume that the (integer) sequence $a(n)$ satisfies

$$a(n) \sim \kappa \mu^n n^\gamma, \quad n \rightarrow \infty.$$

If the associated series $A(t)$ is **rational/algebraic/D-finite** then

- μ is algebraic
- γ belongs to \mathbb{N} , $\mathbb{Q} \setminus \{-1, -2, \dots\}$, \mathbb{Q} .

G-functions

[Katz 70, Chudnovsky² 85, André 89... Bostan, Raschel, Salvy 14]

Singularities. A D-finite series $A(t)$ of positive radius of convergence has **finitely many singularities**.

Functional equations. Start from a difference/differential/etc. equation --- **Galois theory**




Quadrant walks [Dreyfus, Hardouin, Roques, Singer 17]

Final comments

- It sometimes takes **a long time** to find a “direct” solution once an “undirect” one has been found.

Planar maps: algebraicity proved in the 60's using a catalytic variable [Tutte]; first direct proof in 81 [Cori-Vauquelin], complete understanding in the 90's [Schaeffer]

- Many algebraicity/D-finiteness results still wait for a **direct proof**






Final comments

- It sometimes takes **a long time** to find a “direct” solution once an “undirect” one has been found.

Planar maps: algebraicity proved in the 60's using a catalytic variable [Tutte]; first direct proof in 81 [Cori-Vauquelin], complete understanding in the 90's [Schaeffer]

- Many algebraicity/D-finiteness results still wait for a **direct proof**






- Interesting questions on **\mathbb{N} -rational and \mathbb{N} -algebraic series**

Final comments

- It sometimes takes **a long time** to find a “direct” solution once an “undirect” one has been found.

Planar maps: algebraicity proved in the 60's using a catalytic variable [Tutte]; first direct proof in 81 [Cori-Vauquelin], complete understanding in the 90's [Schaeffer]

- Many algebraicity/D-finiteness results still wait for a **direct proof**



- Interesting questions on **\mathbb{N} -rational and \mathbb{N} -algebraic series**
- Combinatorialists need specialists of functional equations

