

# Rational and algebraic series in enumerative combinatorics

Mireille Bousquet-Mélou, CNRS, Bordeaux, France

*<http://www.labri.fr/~bousquet>*

## Enumeration and generating functions

Let  $\mathcal{A}$  be a set of discrete objects equipped with a size:

$$\begin{aligned} \text{size} : \mathcal{A} &\rightarrow \mathbb{N} \\ a &\mapsto |a| \end{aligned}$$

Assume that for all  $n$ ,

$$\mathcal{A}_n := \{a \in \mathcal{A} : |a| = n\} \text{ is finite.}$$

Let  $a(n) = |\mathcal{A}_n|$ .

The **generating function** of the objects of  $\mathcal{A}$ , counted by their size, is

$$\begin{aligned} A(t) &:= \sum_{n \geq 0} a(n)t^n \\ &= \sum_{a \in \mathcal{A}} t^{|a|}. \end{aligned}$$

**Applications:** probability, algebra, computer science (analysis of algorithms), statistical physics... and curiosity

## Rational and algebraic formal power series

- The formal power series  $A(t)$  is **algebraic** (over  $\mathbb{Q}(t)$ ) if it satisfies a (non-trivial) polynomial equation:

$$P(t, A(t)) = 0.$$

- The formal power series  $A(t)$  is **rational** if it can be written

$$A(t) = \frac{P(t)}{Q(t)}$$

where  $P(t)$  and  $Q(t)$  are polynomials in  $t$ .

## Rational and algebraic formal power series

- The formal power series  $A(t)$  is **algebraic** (over  $\mathbb{Q}(t)$ ) if it satisfies a (non-trivial) polynomial equation:

$$P(t, A(t)) = 0.$$

- The formal power series  $A(t)$  is **rational** if it can be written

$$A(t) = \frac{P(t)}{Q(t)}$$

where  $P(t)$  and  $Q(t)$  are polynomials in  $t$ .

- A class of objects having a rational [algebraic] generating function will be said to be **rational [algebraic]**.

## Some charms of rational and algebraic series

- Closure properties (+, ×, /, derivatives, composition...)
- “Easy” to handle (partial fraction decomposition, Puiseux expansions, elimination, resultants, Gröbner bases...)
- Algebraicity can be guessed from the first coefficients (GFUN)
- The coefficients can be computed in a linear number of operations.
- (Almost) automatic asymptotics of the coefficients: in general,

$$a(n) \sim \frac{\kappa}{\Gamma(d+1)} \mu^n n^d,$$

where  $\kappa$  and  $\rho$  are algebraic over  $\mathbb{Q}$  and  $d \in \mathbb{Q} \setminus \{-1, -2, \dots\}$ .

**What does it mean for a class of objects to have  
a rational [algebraic] generating function?**

## Combinatorial constructions and operations on series: A dictionary

Construction	Numbers	Generating function
Union $\mathcal{A} = \mathcal{B} \uplus \mathcal{C}$	$a(n) = b(n) + c(n)$	$A(t) = B(t) + C(t)$
Product $\mathcal{A} = \mathcal{B} \times \mathcal{C}$	$a(n) = b(0)c(n) + \dots$ $+b(n-1)c(1) + b(n)c(0)$	$A(t) = B(t) \cdot C(t)$
$ (\beta, \gamma)  =  \beta  +  \gamma $		

“Symbolic Combinatorics” [Flajolet-Sedgewick]

## A rational example: integer compositions


- Let  $n \in \mathbb{N}$ . A **composition** of size  $n$  with  $k$  parts is a  $k$ -tuple  $c = (n_1, \dots, n_k)$  with  $n_i \in \{1, 2, \dots\}$  for all  $i$  and  $n_1 + \dots + n_k = n$ .



(1)




(2)




(1, 1)




(3)



(2, 1)



(1, 2)




(1, 1, 1)

## A rational example: integer compositions

• Let  $n \in \mathbb{N}$ . A **composition** of size  $n$  with  $k$  parts is a  $k$ -tuple  $c = (n_1, \dots, n_k)$  with  $n_i \in \{1, 2, \dots\}$  for all  $i$  and  $n_1 + \dots + n_k = n$ .


• Let  $C(t)$  be the **generating function** of compositions, counted by their **size**. Then

$$C(t) = t + tC(t) + tC(t).$$




$$C(t) = \frac{t}{1 - 2t}.$$

## An algebraic example: binary trees



- Let  $B(t)$  be the **generating function** of binary trees, counted by the number of (inner) nodes. Then

$$B(t) = 1 + tB(t)^2$$




$$B(t) = \frac{1 - \sqrt{1 - 4t}}{2t} = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} t^n$$

Catalan number.

# Rational and algebraic series: a combinatorial intuition

- “Linear” structure of rational objects




- “Tree-like” structure of algebraic objects



# Rational and algebraic series: a combinatorial intuition

- “Linear” structure of rational objects



- “Tree-like” structure of algebraic objects



1. **Formalize this intuition:** tree-like structures,  $\mathbb{N}$ -algebraic series
2. **Are all algebraic objects tree-like?**
3. How does one prove algebraicity?

Similar questions for rational objects and linear structures

# Part 1: Tree-like structures and $\mathbb{N}$ -algebraic series

[Formal languages – Regular and context-free languages]

## Tree-like constructions

- Building blocks: a finite collection of initial objects (atoms)
- Take (disjoint) unions:  $(A, B) \rightarrow A + B$
- Take cartesian products:  $(A, B) \rightarrow A \cdot B$
- The construction may involve  $k$  different classes of objects  $\mathcal{A}_1, \dots, \mathcal{A}_k$

Example:

$$\begin{aligned} A_1 &= 1 + 2t + 4tA_1 + (12t + t^2)A_2 + t^2A_1A_2^2 \\ A_2 &= 14tA_1 + 8tA_2 + 3tA_1^3 \end{aligned}$$

## Tree-like constructions

- Building blocks: a finite collection of initial objects (atoms)
- Take (disjoint) unions:  $(A, B) \rightarrow A + B$
- Take cartesian products:  $(A, B) \rightarrow A \cdot B$
- The construction may involve  $k$  different classes of objects  $\mathcal{A}_1, \dots, \mathcal{A}_k$

**Example:**

$$\begin{aligned} A_1 &= 1 + 2t + 4tA_1 + (12t + t^2)A_2 + t^2A_1A_2^2 \\ A_2 &= 14tA_1 + 8tA_2 + 3tA_1^3 \end{aligned}$$

**Definition:** A series  $A$  is  **$\mathbb{N}$ -algebraic** if there exists a  $k$ -tuple of formal power series  $\mathcal{A} = (A_1, \dots, A_k)$  such that  $A = A_1$  and for  $1 \leq j \leq k$

$$\mathcal{A} = \begin{pmatrix} A_1 \\ \vdots \\ A_k \end{pmatrix} = \Phi(\mathcal{A}) = \begin{pmatrix} P_1(t, A_1, \dots, A_k) \\ \vdots \\ P_k(t, A_1, \dots, A_k) \end{pmatrix}$$

where

- each  $P_j$  is a polynomial **with coefficients in  $\mathbb{N}$**
- the map  $\mathcal{B} \mapsto \Phi^m(\mathcal{B})$  is a contraction for some  $m$ .

## Tree-like constructions

- Building blocks: a finite collection of initial objects (atoms)
- Take (disjoint) unions:  $(A, B) \rightarrow A + B$
- Take cartesian products:  $(A, B) \rightarrow A \cdot B$
- The construction may involve  $k$  different classes of objects  $\mathcal{A}_1, \dots, \mathcal{A}_k$

**Example:**

$$\begin{aligned} A_1 &= 1 + 2t + 4tA_1 + (12t + t^2)A_2 + t^2A_1A_2^2 \\ A_2 &= 14tA_1 + 8tA_2 + 3tA_1^3 \end{aligned}$$

**Definition:** A series  $A$  is  **$\mathbb{N}$ -algebraic** if there exists a  $k$ -tuple of formal power series  $\mathcal{A} = (A_1, \dots, A_k)$  such that  $A = A_1$  and for  $1 \leq j \leq k$

$$\mathcal{A} = \begin{pmatrix} A_1 \\ \vdots \\ A_k \end{pmatrix} = \Phi(\mathcal{A}) = \begin{pmatrix} P_1(t, A_1, \dots, A_k) \\ \vdots \\ P_k(t, A_1, \dots, A_k) \end{pmatrix}$$

where

- each  $P_j$  is a polynomial **with coefficients in  $\mathbb{N}$**
- the map  $\mathcal{B} \mapsto \Phi^m(\mathcal{B})$  is a contraction for some  $m$ .

**Property:** The series  $A_i$  are uniquely defined by this system, have **coefficients in  $\mathbb{N}$**  and are **algebraic**.

## Linear constructions

- Building blocks: a finite collection of initial objects (atoms)
- Take (disjoint) unions:  $(A, B) \rightarrow A + B$
- Take cartesian products **with atoms only**:  $A \rightarrow t^d A$
- The construction may involve  $k$  different types of objects

**Example:**

$$\begin{aligned} A_1 &= 1 + 2t + 4tA_1 + (12t + t^2)A_2 \\ A_2 &= 14tA_1 + 8tA_2 \end{aligned}$$

**Definition:** A series  $A$  is  **$\mathbb{N}$ -rational** if there exists a  $k$ -tuple of formal power series  $\mathcal{A} = (A_1, \dots, A_k)$  such that  $A = A_1$  and for  $1 \leq j \leq k$


$$\mathcal{A} = \begin{pmatrix} A_1 \\ \vdots \\ A_k \end{pmatrix} = \Phi(\mathcal{A}) = \begin{pmatrix} P_1(t, A_1, \dots, A_k) \\ \vdots \\ P_k(t, A_1, \dots, A_k) \end{pmatrix}$$

where

- each  $P_j(t, x_1, \dots, x_k)$  is a polynomial **with coefficients in  $\mathbb{N}$**  of total degree at **most 1** (in the  $x_i$ ),
- the map  $\mathcal{B} \mapsto \Phi^m(\mathcal{B})$  is contractant for some  $m$ .

**Property:** The series  $A_i$  are uniquely defined by this system, have **coefficients in  $\mathbb{N}$**  and are **rational**.

## Combinatorial models for $\mathbb{N}$ -rational and $\mathbb{N}$ -algebraic series



- Let  $A(t)$  be an  $\mathbb{N}$ -rational series. There exists a directed graph  $G$ , and two vertices  $v_1$  and  $v_2$  in  $G$  such that  $A(t) - A(0)$  counts walks going from  $v_1$  to  $v_2$  in the graph (by their length).
- Let  $A(t)$  be an  $\mathbb{N}$ -algebraic series. There exists a family of coloured trees that has generating function  $A(t) - A(0)$ .

**Part 2: Are all algebraic objects tree-like?**

## Are all algebraic objects tree-like?

Let  $\mathcal{A}$  be a class of objects with an algebraic generating function  $A(t)$ .

1.  $\mathbb{N}$ -algebraicity

Is the series  $A(t)$   $\mathbb{N}$ -algebraic?

2. Tree-like structure

Is a tree-like structure visible directly on the objects?

How easy is it to **detect** that  $\mathcal{A}$  has an algebraic generating function?

Similar questions for rational GFs and linear structures.

## Are all algebraic objects tree-like?

Let  $\mathcal{A}$  be a class of objects with a rational [algebraic] generating function  $A(t)$ .

1.  $\mathbb{N}$ -rationality/ $\mathbb{N}$ -algebraicity

Is the series  $A(t)$   $\mathbb{N}$ -rational [ $\mathbb{N}$ -algebraic]?

Rational GF

$\approx$  YES

I have never met a counting problem  
with a rational,  
but not  $\mathbb{N}$ -rational  
solution

Algebraic GF

???

I do not know how to detect  
 $\mathbb{N}$ -algebraicity

## A criterion for $\mathbb{N}$ -rationality [Soittola]

A rational series  $A(t) = \sum_k a(k)t^k$  with coefficients in  $\mathbb{N}$  is  $\mathbb{N}$ -rational iff

- either  $\sum_k a(k)t^k$  has a unique pole of minimal modulus (a *dominant* pole),
- or each of the series  $\sum_k a(2k)t^k$  and  $\sum_k a(2k+1)t^k$  have a unique dominant pole,
- or each of the series  $\sum_k a(3k)t^k$ ,  $\sum_k a(3k+1)t^k$  and  $\sum_k a(3k+2)t^k$  have a unique dominant pole,
- etc.

## A criterion for $\mathbb{N}$ -rationality [Soittola]

A rational series  $A(t) = \sum_k a(k)t^k$  with coefficients in  $\mathbb{N}$  is  $\mathbb{N}$ -rational iff

- either  $\sum_k a(k)t^k$  has a unique pole of minimal modulus (a *dominant* pole),
- or each of the series  $\sum_k a(2k)t^k$  and  $\sum_k a(2k+1)t^k$  have a unique dominant pole,
- or each of the series  $\sum_k a(3k)t^k$ ,  $\sum_k a(3k+1)t^k$  and  $\sum_k a(3k+2)t^k$  have a unique dominant pole,
- etc.

**Application:** The series

$$A(t) = \frac{1 - 2t + 225t^2}{(1 - 25t)(625t^2 + 14t + 1)}$$

has non-negative integer coefficients, but is **NOT**  $\mathbb{N}$ -rational.

## A criterion for $\mathbb{N}$ -algebraicity?

**Question:** Given an algebraic series  $A(t)$ , how can we decide whether it is  $\mathbb{N}$ -algebraic?

At least, can we state a **necessary** condition for an algebraic series to be  $\mathbb{N}$ -algebraic?

**Suggestion:** look at the singularities of  $A(t)$

**What are the possible singular behaviours  
of an  $\mathbb{N}$ -algebraic series?**

## Singular behaviour of algebraic series

### Theorem

Let  $A(t) = \sum_n a(n)t^n$  be an **algebraic** series with coefficients in  $\mathbb{N}$ .

Let  $\rho$  be its radius of convergence. Then  $A(t)$  is singular at  $\rho$ , with a singular expansion of the form

$$A_{\text{sing}}(t) = C(1 - t/\rho)^\gamma + o((1 - t/\rho)^\gamma)$$

where  $\gamma \in \mathbb{Q} \setminus \mathbb{N}$ .

[Newton-Puiseux]

## Singular behaviour of $\mathbb{N}$ -algebraic series

(Future) theorem

Let  $A(t) = \sum_n a(n)t^n$  be an  $\mathbb{N}$ -algebraic series.

Let  $\rho$  be its radius of convergence. Then  $A(t)$  is singular at  $\rho$ , with a singular expansion of the form

$$A_{\text{sing}}(t) = C(1 - t/\rho)^\gamma + o((1 - t/\rho)^\gamma)$$

where


$$\gamma = \frac{k}{2d},$$

for  $k \in \{1\} \cup \{-1, -2, -3, \dots\}$  and  $d \in \{1, 2, 3, \dots\}$

[mbm-Schaeffer, in preparation]

## Rooted planar maps: a candidate for non- $\mathbb{N}$ -algebraicity

Proper embedding of a connected planar graph in the plane.  
One edge is oriented, with the infinite face on its right.



Enumeration of planar maps by edges [Tutte 63]

$$M = A - tA^3, \quad \text{with} \quad A = 1 + 3tA^2.$$

**Remark.**  $A_{\text{sing}}(t) \sim C(1 - 12t)^{3/2} \Rightarrow$  A good candidate for non- $\mathbb{N}$ -algebraicity

## Are all algebraic objects tree-like?

### 2. Tree-like structure

How easy is it to **detect** that  $\mathcal{A}$  has a rational [algebraic] generating function?  
Is a linear [tree-like] structure visible directly on the objects?

Rational GF	Algebraic GF
$\approx$ y e s	NO
<b>Detecting rational GFs and a linear structure is (usually) easy</b>	<b>Many mysterious algebraic objects</b>

## Part 3: A gallery of “hard” algebraic objects

## Algebraic objects with no obvious tree-like structure

### 1. Brute force approach

Describe **any** recursive construction of the objects. Convert it into **any** functional equation for the generating function. Prove that the solution of this equation is algebraic.

### 2. More combinatorics

Describe a recursive construction of the objects based on **concatenation** that gives **directly** a system of algebraic equation(s).


Both approaches may require to describe first a **bijection** with another class of objects.

## Planar maps: the brute force approach

Let

$$F(t, u) \equiv F(u) = \sum_M t^{e(M)} u^{d(M)}$$

where  $e(M)$  is the number of edges of  $M$  and  $d(M)$  is the degree of the infinite face. Then



$$F(u) = 1 + tu^2 F(u)^2 + \frac{tu}{u-1} (uF(u) - F(1))$$

Polynomial equation with one catalytic variable,  $u$ .

[Tutte 68]

# Polynomial equations with one catalytic variable [MBM-Jehanne 05]

- Example

$$F(u) = 1 + tuF(u)^3 + \frac{tu(F(u) - F(1))(2F(u) + F(1))}{u - 1} + \frac{tu(F(u) - F(1) - (u - 1)F'(1))}{(u - 1)^2}.$$

- General framework

Assume a  $(k+1)$ -tuple  $(F(u), F_1, \dots, F_k)$  of formal power series in  $t$  is completely defined by the equation


$$P(F(u), F_1, \dots, F_k, t, u) = 0 \tag{1}$$

Typically,  $F(u) \equiv F(t, u)$  has **polynomial** coefficients in  $u$ , and  $F_i \equiv F_i(t)$  is the coefficient of  $u^{i-1}$  in  $F(t, u)$ .

- Results

1. The solution of every proper equation of this type is **algebraic**.
2. A practical strategy allows to solve specific examples (that is, to derive from (1) an algebraic equation for  $F(u)$ , or  $F_1, \dots, F_k$ ).

## Hard particles on planar maps




- $F(u) = 1 + G(u) + tu^2 F(u)^2 + \frac{tu(uF(u) - F_1)}{u - 1}$
- $G(u) = tuF(u) + tuF(u)G(u) + \frac{tu(G(u) - G_1)}{u - 1}$


$$t^2 F(1) = T^2(1 - 7T + 16T^2 + T - 15T^3 + 4T^4) \quad \text{with} \quad T(1 - 2T)(1 - 3T + 3T^2) = t.$$

## Rooted planar maps: a combinatorial explanation

- **Bijection** between planar maps ( $n$  edges) and *balanced* blossoming trees ( $n$  nodes) [Schaeffer 97]




A planar map (GF  $M$ )



A blossoming tree  $A = 1 + 3tA^2$

- An *unbalanced* blossoming tree is, in essence, a 3-tuple of blossoming subtrees [Bouttier *et al.* 02]:


$$\Rightarrow M = A - tA^3.$$



**Find combinatorial explanations for...**

## Walks on the slit plane

$$S(u, v, t) := \sum_{i, j, n} a(i, j, n) u^i v^j t^n$$



- Why is  $S(u, v; t)$  algebraic? Let  $\bar{u} = 1/u$  and  $\bar{v} = 1/v$ . Then

$$S(u, v; t) = \frac{\left(1 - 2t(1 + \bar{u}) + \sqrt{1 - 4t}\right)^{1/2} \left(1 + 2t(1 - \bar{u}) + \sqrt{1 + 4t}\right)^{1/2}}{1 - t(u + \bar{u} + v + \bar{v})}.$$

- Why is  $S_{i, j}(t) := \sum_n a(i, j; n) t^n$  algebraic for all  $i$  and  $j$ ?

$$a(1, 0; 2n + 1) = C_{2n+1}, \quad a(0, 1; 2n + 1) = 4^n C_n, \quad a(-1, 1; 2n) = C_{2n} \dots$$


[Barcucci et al. 01]

[MBM-Schaeffer 02] [MBM 01]

$$C_n = \frac{1}{n+1} \binom{2n}{n} = \text{nth Catalan}$$

## Kreweras' walks in the quarter plane

$$Q(t) := \sum_n a(n)t^n$$



$n$  steps  
 $a(n)$

- Why is  $Q(t)$  algebraic? Let  $W = t(2 + W^3)$ . Then


$$Q(t) = 2 \frac{(1/W - 1)\sqrt{1 - W^2}}{1 - 3t} - \frac{1}{t}.$$

- Why is  $a(i, 0; n)$ , the number of walks ending at  $(i, 0)$ , nice and simple?

$$a(i, 0; 3n + 2i) = \frac{4^n (2i + 1)}{(n + i + 1)(2n + 2i + 1)} \binom{2i}{i} \binom{3n + 2i}{n}$$

[Kreweras 65], [Niederhausen 83], [Gessel 86],  
[MBM 04], [Bernardi 06]

## Embedded binary trees



The generating function of binary trees of right width  $\leq j$  is:

$$W_{\leq j} = A \frac{(1 - Z^{j+2})(1 - Z^{j+7})}{(1 - Z^{j+4})(1 - Z^{j+5})},$$

where  $A$  counts binary trees and  $Z \equiv Z(t)$  is the unique series in  $t$  such that

$$Z = t \frac{(1 + Z^2)^2}{(1 - Z + Z^2)} \quad \text{and} \quad Z(0) = 0.$$

[mbm 05]

But. . . w h y ?

## Concluding remarks


- Not all generating functions are algebraic!

Analytic techniques - Asymptotics:

[Flajolet 87]

Algebraic techniques - Reduction modulo  $p$ :

[Allouche 97]



- One more step in the hierarchy of generating functions

Type of series	Combinatorial intuition
polynomials	finite sets
rational	linear structures
algebraic	algebraic structures
D-finite	???

D-finite series  $\leftrightarrow$  P-recursive sequences:

$$P_0(n)a(n) + P_1(n)a(n-1) + \cdots + P_k(n)a(n-k) = 0.$$

## Reverse engineering?

For

$$f_n = \frac{(2n)!}{n!^2}, \quad f_n = \frac{(6n)!(n)!}{(3n)!(2n)!^2}, \quad f_n = \frac{(10n)!(n)!}{(5n)!(4n)!(2n)!}, \quad f_n = \frac{(20n)!(n)!}{(10n)!(7n)!(4n)!},$$

the series

$$F(t) = \sum_{n \geq 0} f_n t^n$$

has coefficients in  $\mathbb{N}$  and is algebraic [Rodriguez-Villegas 05].

Does it count something?