

The diagram shows a blue cloud-like shape containing the word "Macsyma". A blue arrow points from this cloud towards a central area. To the left of the arrow is a small diagram of a convex polyomino. To the right is a larger diagram titled "the Catalan garden" which shows various paths and shapes related to combinatorial mathematics.

Fig 1. A convex polyomino.

the Catalan garden

Macsyma

Theorem 1.1. The number p_{2n} of convex polyominoes having a perimeter $2n$ is

$$p_4 = 1, \quad p_6 = 2$$


and for $n \geq 0$,

$$p_{2n+8} = (2n+11)4^n - 4(2n+1)\binom{2n}{n}.$$



The collage includes several photographs of men, some with blue starburst overlays indicating specific individuals. It also features a map of Canada with a red highlighted region, a vintage Apple computer monitor, and a color bar at the bottom.

1989

1984








What are Strahler numbers ?

Ershov 1958


- Defined on **binary** trees



What are Strahler numbers ?


Ershov 1958

- Defined on **binary** trees




What are Strahler numbers ?

- $3 \times (4+5) - (2 \times 7 + 1)$




Strahler – pruning

Vauchaussade Viennot (1985), Zeilberger (1990)
Defined on rooted planar trees



Strahler – pruning


Vauchaussade Viennot (1985), Zeilberger (1990)
Définé on rooted planar maps




Strahler – an other way

Auber&alt 2003


- Node's children ordering
- Computing node value




Strahler – an other way




Strahler – an other way




Strahler – an other way



Strahler – an other way



Strahler – an other way



New Strahler numbers for rooted plane trees

D. Auber, J.P. Domenger, M. Delest, P. Duchon, J.M. Fédou



Proceedings of Third Colloquium on Mathematics and Computer Science
Algorithms, Trees, Combinatorics and Probabilities 2004

Theorem 5.5 *The branching ratio of trees of any family of finite degree D is asymptotically 4.*





Strahler – for graphs

Quadri-Partition gives a pair of integers





Strahler et ARN



Classification by ARN order
Waterman (1978)


Strahler et ARN



Classification complexité ARN
(Viennot & al., 1985)


Drawing ARN

- Sequence : AGGCCUUAGGCCUU
- liens: 1, 14, 1, 4, 3, 7, 5, 10, 6, 9, 8, 11




Drawing ARN


Reduction to one page





Drawing ARN





Drawing ARN



ARN dessin




ARN dessin






J. Graphs & Appl. Auber & Alt 2006

ARN Visualization?





ARN Visualization?





Parallelogram polyominoes having area n



Parallelogram polyominoes having area n


$$\begin{array}{c} \text{A 3D representation of a parallelogram polyomino of area n, composed of blue and red squares.} \\ = \end{array}$$



$$\begin{array}{ccccccc} *_0 & + & *_q & + & *_q & + & *_q \\ \text{Blue triangle} & & \text{Blue triangle} & & \text{Blue triangle} & & \text{Red triangle} \end{array}$$


$$D = xD\bar{x}D + x\bar{t}\bar{x}D + xD\bar{x} + x\bar{t}\bar{x}$$

1988




$$D = x D \bar{x} D + x t \bar{x} D + x D \bar{x} + x t \bar{x}$$

perimeter	Catalan
area	
Area and columns	



perimeter	Catalan
area	q-analog of Catalan
Area and columns	q-analog of Bessel functions

Maylis! Jean-Marc! It's great ! Fantastic !



perimeter	Catalan
area	q -analog of Catalan
Area and columns	q -analog of Bessel functions

Bijection with paths in binary trees (Fedou 89)

Proposition 2.10 — Let $T_{d,n}(x, y)$ be the area generating function for animals on the n -decorated triangular lattice:

$$T_{d,n}(x, y) = \sum_A x^{|T(A)|} y^{|A| - |T(A)|}$$

where $|T(A)|$ is the number of cells of A on the underlying triangular lattice. We have

$$T_{d,n}(x, y) = T_1\left(x \frac{1-y^{n+1}}{1-y}, y^n \frac{1-y}{1-y^{n+1}}, 1, 1, 1\right)$$

where $T_1(t, u, v, w, x)$ is defined by (8).

Théorème 4.6 Les ensembles de permutations à motifs exclus $S_n(1234, 1243, 1423, 4123)$, $S_n(1324, 1342, 1432, 4132)$, $S_n(2134, 2143, 2413, 4213)$, $S_n(2314, 2413, 3142, 3241)$, $S_n(1234, 1324, 2134, 2314)$, $S_n(1234, 2134, 2314, 3124)$, $S_n(1324, 2134, 2314, 3124)$, $S_n(1324, 2314, 3124, 3214)$, $S_n(1324, 2314, 3124, 3214)$, $S_n(1324, 1342, 3124, 3214)$, $S_n(1324, 1342, 3142, 3214)$, $S_n(1324, 1342, 2314, 2341)$, sont en bijection avec les mots de $GD_{\leq 3}$ de longueur $2n - 2$, et sont donc énumérés par le coefficient binomial central

$$\binom{2n-2}{n-1}$$

Proposition 2.10 Let $T_{G,n}(x,y)$ be the area generating function for animals on the n -decorated triangular lattice:



$$T_{G,n}(x,y) = \sum_{A \in \mathcal{A}} x^{\text{area}(A)} y^{|\text{cells}(A)|}$$


where $|\text{cells}(A)|$ is the number of cells of A on the underlying triangular lattice. We have

$$T_{G,n}(x,y) = T_1 \left(\frac{1-y^{n+1}}{1-y^n} \cdot y^{\frac{1-y}{1-y^{n+1}}} \cdot \frac{1-y}{1-y^{n+1}}, 1, 1 \right)$$

where $T_1(t, a, c, v, x)$ is defined by (8).

FIG. 3 – Six opérations d'édition sur deux séquences






Many algorithms split data based on a quality measure

Manconridis (98)



$$MQ(G; C) = \frac{1}{k} \left(\sum_{i=1}^k \frac{e_{ii}}{t_i^2} - \frac{1}{k-1} \sum_{\substack{j=1 \\ j \neq i}}^k \frac{e_{ij}}{t_i t_j} \right)$$

The graph The partition Number of edges between cluster i and cluster j
Number of nodes of cluster j

Usually one algorithm is run recursively on the graph
-> fragmentation tree




Usually one algorithm is run recursively on the graph
 -> fragmentation tree



Bijection with paths in binary trees (Fedou 89)

First : modify MQ


$$MQ(G;C) = \frac{1}{k} \left(\sum_{i=1}^k \frac{e_{ii}}{t_i^2} - \frac{1}{k-1} \sum_{\substack{j=1 \\ j \neq i}}^k \frac{e_{ij}}{t_i t_j} \right)$$



$$MQ(G;C) = \frac{1}{\sum_{i=1}^k t_i} \sum_{i=1}^k \left(\frac{t_i e_{ii}}{t_i^2} \right) - \frac{1}{\sum_{\substack{i=1 \\ j=1 \\ j \neq i}}^k t_i t_j} \sum_{\substack{j=1 \\ j \neq i}}^k \frac{t_i t_j e_{ij}}{t_i t_j}$$


Second : simplify notations

$$MQ(G;C) = \frac{1}{\sum_{i=1}^k t_i} \sum_{i=1}^k \left(\frac{t_i e_{ii}}{t_i^2} \right) - \frac{1}{\sum_{\substack{i=1 \\ j=1 \\ j \neq i}}^k t_i t_j} \sum_{\substack{j=1 \\ j \neq i}}^k \frac{t_i t_j e_{ij}}{t_i t_j}$$



$$MQ(G;C) = \frac{1}{\sum_{i=1}^k x_i} \left(\sum_{i=1}^k \frac{x_i e_{ii}}{\delta_i} - \frac{1}{\sum_{\substack{i=1 \\ i \neq j}}^k x_i} \sum_{\substack{j=1 \\ i \neq j}}^k \frac{x_i e_{ij}}{\delta_{ij}} \right)$$

Second : simplify notations



$$MQ(G; C) = \frac{1}{\sum_{i=1}^k x_i} \left(\sum_{i=1}^k \frac{x_i e_{i,i}}{\delta_i} - \frac{1}{\sum_{i=1}^k x_i} \sum_{\substack{j=1 \\ i \neq j}}^k x_j e_{i,j} \right)$$

$$MQ(G, T; q) = \begin{cases} 0 & \text{if } k=0 \\ \frac{1}{X} \sum_{i=1}^k x_i (\alpha_i - \beta_i)(1 + qMQ(G_i, T_i; q)) & \text{otherwise} \end{cases}$$

- This formula makes sense on small tractable examples
 - The same idea can be applied to others quality measures
 - It can be interpreted as sum of paths in the graph



2010

