

COMBINATORICS OF LABELLED PARALLELOGRAM POLYOMINOES

J.-C. AVAL, F. BERGERON, AND A. GARSIA

ABSTRACT. We obtain explicit formulas for the enumeration of labelled parallelogram polyominoes. These are the polyominoes that are bounded, above and below, by north-east lattice paths going from the origin to a point (k, n) . The numbers from 1 to n (the labels) are bijectively attached to the n north steps of the above-bounding path, with the condition that they appear in increasing values along consecutive north steps. We calculate the Frobenius characteristic of the action of the symmetric group \mathbb{S}_n on these labels. All these enumeration results are refined to take into account the area of these polyominoes. We make a connection between our enumeration results and the theory of operators for which the integral Macdonald polynomials are joint eigenfunctions. We also explain how these same polyominoes can be used to explicitly construct a linear basis of a ring of SL_2 -invariants.

CONTENTS

1. Introduction	1
2. Paths and labelled paths	2
3. Parallelogram polyominoes	5
4. Parallelogram polyominoes, as indexing set of SL_2 -invariants	8
5. Labelled parallelogram polyominoes	10
6. Doubly labelled polyominoes	14
7. Operators and Macdonald polynomials	16
8. Formula for the q -Frobenius characteristic of the labelled polyomino module	19
9. Thanks and future considerations	23
References	24

1. INTRODUCTION

Parallelogram polyominoes have been studied by many authors (see [5, 11, 8] for a nice survey and enumeration results). They correspond to pairs $\pi = (\alpha, \beta)$ of north-east paths going from the origin to a point (k, n) in the combinatorial plane $\mathbb{N} \times \mathbb{N}$, with the path α staying “above” the path β . Our aim here is to study properties, and related nice formulas, of “labelled parallelogram polyominoes”. These are obtained by bijectively labelling each of the n north steps of the path α with the numbers between 1 and

Date: November 26, 2014.

This work has been supported by ANR-France (PSYCO project), by CNRS-France (LIA LIRCO), by NSERC-Canada and by NSF-USA.

n . Our motivation stems from a similarity between this new notion and recent work on labelled intervals in the Tamari lattice, in connection with the study of trivariate diagonal harmonic polynomials for the symmetric group (see [4]).

We calculate explicitly the Frobenius characteristic of the natural action of the symmetric group \mathbb{S}_n on these labelled polyominoes; and study aspects of a weighted version of this Frobenius characteristic with respect to the area of the polyominoes. This connects our study to interesting operators for which adequately normalized Macdonald polynomials are joint eigenfunctions. This is the same theory that appears in the study of the \mathbb{S}_n -module of bi-variate diagonal harmonics (see [13, 14, 10]).

We also extend some of our considerations to parallelogram polyominoes, with added labels on east steps of the below-bounding path; with a corresponding action of the group $\mathbb{S}_k \times \mathbb{S}_n$. Several components of these spaces are naturally related to parking function modules.

2. PATHS AND LABELLED PATHS

Let k and n be two positive integers and set $N = k + n$. A $k \times n$ north-east (lattice) path in $\mathbb{N} \times \mathbb{N}$ is a sequence $\alpha = (p_0, \dots, p_i, \dots, p_N)$ of points $p_i = (x_i, y_i)$ in $\mathbb{N} \times \mathbb{N}$, with $p_0 = (0, 0)$ and $p_N = (k, n)$, and such that

$$(x_{i+1}, y_{i+1}) = \begin{cases} (x_i, y_i) + (1, 0) & \text{an east step, or} \\ (x_i, y_i) + (0, 1) & \text{a north step.} \end{cases}$$

Figure 1 gives an example of a 10×6 north-east lattice path. We denote by $P_{k,n}$ the set of $k \times n$ north-east paths. These paths are often encoded as a word $\omega = w_1 w_2 \cdots w_N$, on the alphabet $\{x, y\}$, with x standing for an east step, and y for a north step; and k is the number of x 's, while n is that of y 's. Thus, the path of Figure 1 is encoded as $yyxxxxyxyxyxxxxyxx$.

Another description of such a path α may be given in terms of the sequence of ‘‘heights’’ of its k horizontal steps (written from left to right), so that we may write

$$\alpha = a_1 a_2 \cdots a_k.$$

More specifically, a_i is equal to the unique y_j such that $x_j = i$ and $x_{j-1} = i - 1$. We say that this is the *height sequence* description of α . Thus, the top path in Figure 1 may be encoded as the height sequence 2224455566. We observe that the height sequence is increasing, *i.e.* $a_i \leq a_{i+1}$; and that any increasing sequence $a_1 a_2 \cdots a_k$, with $0 \leq a_i \leq n$ corresponds to a unique path.

Exchanging the role of the axes, we may also describe α in terms of its *indentation sequence*. This is simply the bottom to top sequence of distance between vertical steps in α , and the y -axis. For example, the indentation sequence of our running example is 003358. Once again, this establishes a bijection between $k \times n$ north-east lattice paths, and length n increasing sequences of values between 0 and k .

There is a classical bijection between north-east paths in $P_{k,n}$ and monomials of degree k in the variables $\mathbf{x} = (x_1, \dots, x_{n+1})$. One simply associates to a height sequence $\alpha = (a_1, a_2, \dots, a_k)$, the monomial

$$\mathbf{x}_\alpha := x_{a_1+1}x_{a_2+1} \cdots x_{a_k+1}.$$

Thus, we get the well known formula for the *Hilbert series*, denoted $R(x)$, of the polynomial ring $R = \mathbb{C}[\mathbf{x}]$:

$$R(x) = \sum_{k=0}^{\infty} \binom{n+k}{k} x^k = \left(\frac{1}{1-x} \right)^{n+1}. \quad (2.1)$$

Recall that the coefficients of this series correspond to the dimension of the homogeneous components of the space considered.

The q -weighted *area* enumeration of $P_{k,n}$, is defined to be

$$P_{k,n}(q) := \sum_{\alpha \in P_{k,n}} q^{\text{area}(\alpha)}, \quad (2.2)$$

with $\text{area}(\alpha)$ equal to the number of cells in $\mathbb{N} \times \mathbb{N}$ lying below the path α . In Figure 1, the area of the path is the shaded region. It is also well known that $P_{k,n}(q)$ is equal to the classical q -analog of the binomial coefficient:

$$P_{k,n}(q) = \left[\begin{matrix} n+k \\ k \end{matrix} \right]_q = h_k(1, q, \dots, q^{n+1}). \quad (2.3)$$




FIGURE 1. A north-east path, and its area.

Labelled paths. We now add labels to north steps of paths in $P_{k,n}$. We require these labels, going from 1 to n , to be increasing whenever they lie on north steps having same horizontal coordinate. The underlying path, of a labelled path ℓ , is said to be its *shape*. We denote by $L_{k,n}$, the set of labelled paths having shapes lying in $P_{k,n}$. It is easy to see that the number of labelled paths is $(k+1)^n$, with q -weighted enumeration given by the straightforward q -analog

$$\sum_{\alpha \in L_{k,n}} q^{\text{area}(\alpha)} = ([k+1]_q)^n. \quad (2.4)$$

Observe¹ that labelled path ℓ may be bijectively turned into a function from $[n] := \{1, 2, \dots, n\}$ to $[k+1] := \{1, 2, \dots, k+1\}$, as follows. We simply set $\ell(i) := j$, whenever i lies along a north step having first coordinate equal to $j-1$. Thus, the fibers of the resulting function correspond to the labels of consecutive north steps. We say that the *set partition* of ℓ , is the partition of $[n]$ into these fibers.




FIGURE 2. A labelled path with below area equal to 41.

The symmetric group \mathbb{S}_n acts by permuting labels, up to reordering labels that lie one above the other. In fact, it may be best to consider this in the context of the indentation sequence encoding of paths. In this point of view, a labelled path is simply a permutation of this sequence. Thus, $k \times n$ labelled paths are just another name for length n sequences of numbers between 0 and k . The shape α (underlying path), of a labelled path ℓ , is simply the increasing ordering of this sequence. Still we will keep using the path terminology, which is better adapted to our upcoming study of parallelogram polyominoes.

It is a classical result that the corresponding Frobenius characteristic is given by the formula²

$$\text{Frob}(\mathbb{L}_{k,n})(\mathbf{z}) = h_n[(k+1)\mathbf{z}] \quad (2.5)$$

$$= \sum_{\mu \vdash n} (k+1)^{\ell(\mu)} \frac{p_\mu(\mathbf{z})}{z_\mu} \quad (2.6)$$

$$= \sum_{\mu \vdash n} s_\mu[k+1] s_\mu(\mathbf{z}). \quad (2.7)$$

Recall that this last equality says that $s_\mu[k+1]$ is the multiplicity, in $\mathbb{L}_{k,n}$, of the irreducible representation indexed by the partition μ . Taking into account the area, we have

¹Our reason for considering functions from $[n]$ to $[k+1]$ as labelled paths will become apparent in the sequel.

²We follow Macdonald's notation (see [17]) for symmetric functions. Complete symmetric and Schur functions are respectively denoted by the letters h and s . We also use plethystic notations. See [3] for more on this.

a direct q -analog of the previous formula

$$\text{Frob}(\mathbb{L}_{k,n})(\mathbf{z}; q) = h_n \left[\mathbf{z} \frac{q^{k+1} - 1}{q - 1} \right] \quad (2.8)$$

$$= \sum_{\mu \vdash n} s_\mu(1, q, \dots, q^k) s_\mu(\mathbf{z}). \quad (2.9)$$

3. PARALLELOGRAM POLYOMINOES

Beside setting up notations, the aim of this section is to recall basic (well-known) facts about parallelogram polyominoes. For two paths α and β in $\mathbb{P}_{k,n}$, we get a polyomino $\pi = (\alpha, \beta)$ if α stays “above” β . One usually thinks of a parallelogram polyomino $\pi = (\alpha, \beta)$ as the region of the plane bounded above by the path α , and below by the path β . Except for endpoints, all points of the path $\beta = (q_0, \dots, q_i, \dots, q_N)$ are required to be strictly below those of $\alpha = (p_0, \dots, p_i, \dots, p_N)$. Hence, writing $q_i = (x'_i, y'_i)$, this is to say that $y'_i < y_j$ whenever $x'_i = x_j$, for $1 \leq i, j \leq N$. We say that n is the *height* of π , and that k is its *width*; and we denote by $\mathbb{P}_{k,n}$ the set of parallelogram polyominoes of height n and width k . The number of 1×1 boxes lying between the two paths is always larger or equal to $k + n - 1$, and equality holds for *ribbon shapes*. Thus, it is natural to subtract this value in the definition of the *area* of the polyomino π , denoted by $\text{area}(\pi)$. Hence ribbon shapes have zero area. The *area-enumerating polynomial* is then defined as

$$\mathbb{P}_{k,n}(q) := \sum_{\pi \in \mathbb{P}_{k,n}} q^{\text{area}(\pi)}. \quad (3.1)$$

Clearly the reflection in the diagonal $x = y$ maps bijectively the set $\mathbb{P}_{k,n}$ on the set $\mathbb{P}_{n,k}$, we have the symmetry $\mathbb{P}_{k,n}(q) = \mathbb{P}_{n,k}(q)$. This is reflected in the *generating function*

$$\begin{aligned} \mathbb{P}(x, y; q) &:= \sum_{(k,n) \in \mathbb{N} \times \mathbb{N}} \mathbb{P}_{k,n}(q) x^k y^n, \\ &= xy + xy^2 + x^2y + xy^3 + (2 + q)x^2y^2 + x^3y \\ &\quad + xy^4 + (3 + 2q + q^2)x^2y^3 + (3 + 2q + q^2)x^3y^2 + x^4y \\ &\quad + xy^5 + (4 + 3q + 2q^2 + q^3)x^2y^4 \\ &\quad + (6 + 6q + 5q^2 + 2q^3 + q^4)x^3y^3 \\ &\quad + (4 + 3q + 2q^2 + q^3)x^4y^2 + x^5y + \dots \end{aligned}$$

which can be expressed in terms of Bessel functions (see [5]). When $q = 1$, we have

$$\mathbb{P}_{k+1,n}(1) = |\mathbb{P}_{k+1,n}| = s_{k,k}[n + 1] \quad (3.2)$$

$$= \frac{1}{k + 1} \binom{n + k}{k} \binom{n + k - 1}{k}. \quad (3.3)$$

Indeed this is a direct consequence of the principal specialisation formula for Schur functions.




FIGURE 3. A Parallelogram polyomino and the corresponding Motzkin path.

Alternate descriptions. We may shed new light on our study of parallelogram polyominoes by exploiting alternate encodings for them. For paths $\alpha, \beta \in \mathbb{L}_{k+1, n}$, respectively³ given as increasing sequences $\alpha = a_0 a_1 \cdots a_k$ and $\beta = b_1 b_2 \cdots b_{k+1}$, the pair $\pi = (\alpha, \beta)$ is a parallelogram polyomino if and only if $a_0 = 0$, $b_{k+1} = n + 1$, and $a_i > b_i$, for $1 \leq i \leq k$. Hence, we may identify parallelogram polyominoes with *semi-standard tableaux* (see Figure 4) of shape k^2 with values in $\{1, 2, \dots, n + 1\}$, thus explaining formula (3.2). Naturally, we may also describe parallelogram polyominoes in terms of indentation sequences, thus getting semi-standard tableaux of shape n^2 , with values in $\{1, 2, \dots, n\}$. Going from the first encoding of a given polyomino to the second one, establishes a classical bijection between the two equally numerous families of semi-standard tableaux.

a_1	a_2	a_3	a_k
b_1	b_2	b_3	b_k

FIGURE 4. Semi-standard tableau of shape k^2 ; $a_i \leq a_{i+1}$ and $a_i > b_i$.

There is also a well-known bijection (see [8]) between parallelogram polyominoes and *Primitive Motzkin Paths*. Recall that these are the paths, in $\mathbb{N} \times \mathbb{N}$, going from $(0, 0)$ to $(N, 0)$, with steps either north-east, east (either red or blue), or south-east. Thus, the consecutive points, along the path, are linked as follows

$$(x_{i+1}, y_{i+1}) = \begin{cases} (x_i, y_i) + (1, 1) & \text{a north-east step, or} \\ (x_i, y_i) + (\mathbf{1}, 0) & \text{a red-east step, or} \\ (x_i, y_i) + (\mathbf{1}, 0) & \text{a blue-east step, or} \\ (x_i, y_i) + (1, -1) & \text{a south-east step.} \end{cases}$$

Primitive paths are those that never return to the horizontal, except at endpoints. Motzkin paths are often simply presented in terms of *Motzkin words*. Recall that these are the words $\omega = w_1 w_2 \cdots w_N$, on the alphabet $\{d, \bar{d}, r, b\}$, such that

$$(1) \quad |\omega_{\leq i}|_d \geq |\omega_{\leq i}|_{\bar{d}}, \text{ for all } 1 \leq i \leq N - 1,$$

³The shift in indices is intentional here.

$$(2) |\omega|_d = |\omega|_{\bar{d}},$$

where $|\omega|_d$ (resp. $|\omega_{\leq i}|_{\bar{d}}$) stands for the number of d 's occurring in ω (resp. \bar{d}), and $\omega_{\leq i}$ denotes the prefix of length i of ω . The word corresponds to a primitive path, precisely when all inequalities are strict in the first condition. Each letter encodes one of the possible steps: d for north-east, r for red-east, b for blue-east, and \bar{d} for south-east. For a polyomino $\pi = (\alpha, \beta)$, with $\alpha = u_1u_2 \cdots u_N$ and $\beta = v_1v_2 \cdots v_N$ given as words in $\{x, y\}$, Viennot's encoding [8] consists in setting

$$w_i := \begin{cases} d & \text{if } (u_i, v_i) = (x, y), \\ r & \text{if } (u_i, v_i) = (y, y), \\ b & \text{if } (u_i, v_i) = (x, x), \\ \bar{d} & \text{if } (u_i, v_i) = (y, x). \end{cases}$$

See Figure 3 for an example.

Lastly, a parallelogram polyomino may be encoded as a word in the ordered alphabet

$$\mathbb{A} = \{\bar{0}, 1, \bar{1}, 2, \bar{2}, 3, \bar{3}, 4, \dots\}.$$

To do this, we start with its primitive Motzkin path encoding π , and successively replace each step in π by a m -sequence of letters in the \mathbb{A} , $0 \leq m \leq 2$, according to the following rules.

- A down step is replaced by an empty sequence;
- an up step, at height j , is replaced by the 2-sequence $\bar{j}(j+1)$;
- a red horizontal step, at height j , is replaced by \bar{j} ;
- a blue horizontal step at height j is replaced by \bar{j} .

Figure 5 illustrates this process.




FIGURE 5. From Motzkin path to word in \mathbb{A} .

Remark 1. The correspondence between parallelogram polyominoes and primitive Motzkin paths, suggests that we should study “sequences” of parallelogram polyominoes which correspond to general Motzkin paths. In particular, this will make apparent more ties with the study of parking functions that correspond to Dyck path touching the diagonal at specific points.




FIGURE 6. A parallelogram polyomino sequence.

4. PARALLELOGRAM POLYOMINOES, AS INDEXING SET OF SL_2 -INVARIANTS

Part of the motivation for our current study of the combinatorics of parallelogram polyominoes is to tie this study to that of parking functions and labelled intervals in the Tamari poset. Recall that more and more evidence shows that this combinatorics is intimately related to the study of the S_n -modules of bivariate and trivariate diagonal harmonics. Along these lines, it is interesting to observe that the number of polyominoes corresponds to the dimension of the space of SL_2 -invariants of weight $k + 1$ of the Grassmannian $\mathbb{G}(2, n + 1)$, see [19, page 238]. We make this more explicit as follows. Let X be the matrix of $2 \times n$ variables

$$X = \begin{pmatrix} x_1 & x_2 & \cdots & x_n \\ y_1 & y_2 & \cdots & y_n \end{pmatrix}.$$

The special group acts on polynomials $f(X)$, in $\mathbb{C}[X]$, by left multiplication of X by matrices in SL_2 . Invariants under this action are precisely the polynomial expressions in the 2×2 minors⁴ of X . To construct an explicit linear basis of the resulting ring, $\mathcal{R}_n := \mathbb{C}[X]^{SL_2}$, one needs only take into account the Plücker relations on these 2×2 minors. More precisely, as described in [18], we have the exact sequence

$$0 \longrightarrow \mathcal{P}_n \longrightarrow \mathbb{C}[Y_{i,j}]_{1 \leq i < j \leq n} \longrightarrow \mathcal{R}_n \longrightarrow 0, \quad (4.1)$$

where the $Y_{i,j}$ are variables, the third arrow sends $Y_{i,j}$ to $X_{i,j}$ (the 2×2 minor corresponding to the choice of columns i and j in X), and \mathcal{P}_n stands for the ideal generated by the Plücker relations

$$Y_{il}Y_{jk} - Y_{ik}Y_{jl} + Y_{ij}Y_{kl} = 0, \quad \text{for } 1 \leq i < j < k < l \leq n. \quad (4.2)$$

Moreover, using a somewhat reformulated result also discussed in [18, Thm 14.6], it follows that a linear basis of $\mathcal{R}_n = \mathbb{C}[X]^{SL_2}$ may be indexed by parallelogram polyominoes. In fact, we do this in a graded fashion⁵, constructing for each $d \geq 0$ the d -homogeneous part of the basis as a collection of product of d minors. The characterizing property

⁴These are clearly SL_2 -invariants.

⁵Considering 2×2 minors to be of degree 1.

of these products is that they must not be divisible by a binomial $X_{ij}X_{k\ell}$, for $\{i, j\}$ incomparable to $\{k, \ell\}$ relative to the following order on pairs of integers. We set

$$\{i, j\} \prec \{k, \ell\}, \quad \text{iff} \quad i < k \text{ and } j < \ell,$$

assuming, without loss of generality, that $i < j$ and $k < \ell$.

Now, given a polyomino $\pi = (\alpha, \beta) \in \mathbb{P}_{d+1, n-1}$, for which the height sequences of α and β are respectively⁶ $a_0 a_1 \cdots a_d$ and $b_1 b_2 \cdots b_{d+1}$, we consider the minor monomial

$$X_\pi := \prod_{i=1}^d X_{(b_i+1, a_i+1)}. \quad (4.3)$$

The fact that the path β remains below the path α is equivalent to the fact that the minor monomial X_π satisfies the charactering property of the previous paragraph. For example, the minor monomial associated to the polyomino $\pi = (2224455566, 0011112223)$ of Figure 7 is

$$X_\pi = X_{13} X_{23}^2 X_{25}^2 X_{36}^3 X_{47}.$$

Proposition 1. *The family $\{X_\pi\}_\pi$, with π varying in the set of parallelogram polyominoes $\mathbb{P}_{d+1, n-1}$, constitute a linear basis for the degree d homogeneous component of the ring \mathcal{R}_n .*

It follows that, for a given n , the Hilbert series of \mathcal{R}_n is given by the formula

$$\mathcal{R}_n(x) = \sum_{d=0}^{\infty} |\mathbb{P}_{k+1, n}| x^k \quad (4.4)$$

$$= \frac{1}{1 - x^{2n-1}} \sum_{k=0}^{n-2} \frac{1}{k+1} \binom{n-2}{k} \binom{n-1}{k} x^k. \quad (4.5)$$

This approach may readily be expanded to cover families of r non-intersecting paths from $(0, 0)$ to (k, n) , which also parametrize SL_r -invariants. The corresponding enumeration is easily obtained in terms of Schur functions (using Lindström-Gessel-Viennot [12]), giving

$$|\mathbb{P}_{k+1, n}^{(r)}| = s_{k^r}[n+1], \quad (4.6)$$

where $\mathbb{P}_{k, n}^{(r)}$ denotes the set of such families, and k^r stands for the rectangular partition $\underbrace{(k, k, \dots, k)}_r$.

Action of \mathbb{S}_n on \mathcal{R}_n . We may refine the graded enumeration of the ring \mathcal{R}_n , by considering its irreducible decomposition with respect to the action of \mathbb{S}_n , which corresponds to permutations of the n columns of X . Thus we consider the graded Frobenius characteristic

$$\text{Frob}(\mathcal{R}_n)(\mathbf{z}; x) = \sum_{d=0}^{\infty} \text{Frob}(\mathcal{R}_n^{(d)})(\mathbf{z}) x^d, \quad (4.7)$$

⁶The shift in indices is intentional here.

where $\mathcal{R}_n^{(d)}$ stands for the degree d homogeneous component of \mathcal{R}_n .

To calculate this, we follow an approach due to Littlewood for the calculation of the character of the restriction to \mathbb{S}_n of an action of GL_n . It follows from general principles that the character of the action of GL_n on $\mathcal{R}_n^{(d+1)}$, the degree $d+1$ component of \mathcal{R}_n , is equal to the schur function $s_{d^2}(q_1, \dots, q_n)$. The corresponding character value, for the restriction of this action to \mathbb{S}_n , is obtained by evaluating this Schur function at the eigenvalues of permutations. This may be easily calculated, without explicit knowledge of these eigenvalues, in the following manner.

We start by expanding $s_{dd}(q_1, \dots, q_n)$ in terms of the power-sum symmetric function, thus turning the required evaluation into the calculation of the sums of k^{th} -powers of the eigenvalues of any given permutation matrix σ . But this turns out to be equal to the number of fixed points of the k^{th} -power of σ . If μ is the partition giving cycle sizes of σ , this number of fixed points may simply be expressed as

$$\text{fix}(\sigma^k) = \sum_{\mu_i | k} \mu_i. \quad (4.8)$$

Hence, the Frobenius characteristic that we are looking for is given by the formula

$$\mathcal{R}_n^{(d+1)}(\mathbf{z}) = \sum_{\mu \vdash n} s_{dd} \Big|_{p_k \leftarrow \varphi_k(\mu)} \frac{p_\mu(\mathbf{z})}{z_\mu}, \quad (4.9)$$

where we set $\varphi_k(\mu) := \sum_{\mu_i | k} \mu_i$, i.e. the sum of the parts of μ that divide k .

5. LABELLED PARALLELOGRAM POLYOMINOES

A *labelled parallelogram polyomino* is obtained by labeling (as in Section 2) the top path of a parallelogram polyomino. Equivalently, this may be presented in terms of *labelled Motzkin paths*, with labels associated with either north-east steps, or red-east steps. In this case, the relevant increasing-labelling condition simply states that labels should be increasing whenever we they occur on consecutive north-east or red-east steps. Figure 3 gives an example of such a labelling, in both encodings.




FIGURE 7. A labelled parallelogram polyomino.

We say that the underlying parallelogram polyomino is the *shape* of the labelled parallelogram polyomino, and denote by \mathbb{L}_π the set of labelled parallelogram polyominoes of shape π . We also denote by $\mathbb{L}_{k,n}$ ⁷ the set of labelled parallelogram polyominoes of height n and width k , so that

$$\mathbb{L}_{k,n} = \bigcup_{\pi \in \mathbb{P}_{k,n}} \mathbb{L}_\pi.$$

A maximal length sequence of north steps, in a path α , is said to be a *big-step* of α . Clearly, for a height n path α , the sequence of big-step lengths in α forms a composition of n . We denote it by $\gamma(\alpha)$, also setting $\gamma(\pi) := \gamma(\alpha)$ in the case of a parallelogram polyomino $\pi = (\alpha, \beta)$. Recall that it is customary to write $\nu \models n$ when ν is a *composition of n* .

Remark 2. It is surely worth mentioning that, when $k = n$, “classical” Parking Functions may be obtained as special instances of parallelogram polyomino (sequences⁸), in fact in at least three different ways. The first is obtained by the simple device of constraining the lower path to be equal to the *zigzag path*, the one who has height sequence equal to $012 \cdots (n-1)$. In a similar manner, we may fix the lower path to be the one that follows the lower boundary of the surrounding rectangle. Still, an even more natural way is to consider labelled parallelogram polyominoes sequences whose shape is symmetric with respect to the diagonal $x = y$. All of these correspondences are compatible with the action on labels, and area is easy to track. Hence, most of the following formulas may be considered as extensions of analogous formulas for parking functions, potentially in three different ways.

For a height n parallelogram polyomino $\pi = (\alpha, \beta)$, the symmetric group \mathbb{S}_n acts by permutation on the set \mathbb{L}_π , up to a reordering of labels along big-steps. This implies that

$$|\mathbb{L}_\pi| = \binom{n}{\gamma(\pi)}, \tag{5.1}$$

where we use the classical multinomial coefficient notation

$$\binom{n}{\nu} = \frac{n!}{\nu_1! \nu_2! \cdots \nu_k!}, \quad \text{when } \nu = (\nu_1, \nu_2, \dots, \nu_k) \quad \text{and} \quad \nu \models n.$$

It follows (see Proposition 3) that

Proposition 2. *The number of labelled parallelogram polyominoes is given by the formula*

$$|\mathbb{L}_{k,n}| = k^{n-1} \binom{n+k-2}{k-1}, \tag{5.2}$$

and we have the identity

$$k^{n-1} \binom{n+k-2}{k-1} = \sum_{\pi \in \mathbb{P}_{k,n}} \binom{n}{\gamma(\pi)}. \tag{5.3}$$

⁷The top (labelled) path of any element of $\mathbb{L}_{k,n}$ is an element of $\mathbb{L}_{k,n}$ (see Section 2), whence the notation.

⁸This is needed to take into account the possibility that the two paths may touch one another.

It may be worth mentioning that we have the nice generating function

$$\sum_{n=0}^{\infty} k^{n-1} \binom{n+k-2}{k-1} x^n = \left(\frac{1}{1-kx} \right)^k. \quad (5.4)$$

The following result provides a common refinement of formulas (3.3) and (5.2), stated in terms of the Frobenius characteristic of the \mathbb{S}_n -modules \mathbb{L}_π and $\mathbb{L}_{k,n}$ (here considered as free vector spaces over the specified sets).

Writing $\nu = (\nu_1, \nu_2, \dots, \nu_j)$ for $\gamma(\pi)$, it is a classical result that the Frobenius characteristic of \mathbb{L}_π is simply given by the symmetric function $h_\nu(\mathbf{z})$ (in the variables $\mathbf{z} = z_1, z_2, z_3, \dots$). Indeed, the Young subgroup $\mathbb{S}_{\nu_1} \times \dots \times \mathbb{S}_{\nu_j}$ is precisely the stabilizer of the labelled parallelograms polyominoes of shape π . We thus have that

Proposition 3. *The Frobenius characteristic of $\mathbb{L}_{k,n}$ is given by the formula*

$$\text{Frob}(\mathbb{L}_{k,n}) := \sum_{\pi \in \mathbb{P}_{k,n}} h_{\gamma(\pi)}(\mathbf{z}), \quad (5.5)$$

and

$$\text{Frob}(\mathbb{L}_{k,n}) = \frac{1}{k} h_n[k\mathbf{z}] \binom{n+k-2}{k-1} \quad (5.6)$$

$$= \sum_{\lambda \vdash n} k^{\ell(\lambda)-1} \binom{n+k-2}{k-1} \frac{p_\lambda(\mathbf{z})}{z_\lambda} \quad (5.7)$$

$$= \sum_{\mu \vdash n} \frac{1}{k} s_\mu[k] \binom{n+k-2}{k-1} s_\mu(\mathbf{z}), \quad (5.8)$$

where $\ell(\nu)$ (resp. $\ell(\mu)$) stands for the length of a composition ν (resp. partition μ), and k is considered as a constant for the plethystic evaluation in the formulas.

Observe that Formulas (5.2) and (3.3) follow easily from the above expressions, and that (5.6)–(5.8) are polynomial in k . Special values are as follows:

$$\begin{aligned} \text{Frob}(\mathbb{L}_{k,1}) &= h_1(\mathbf{z}) \\ \text{Frob}(\mathbb{L}_{k,2}) &= \binom{k}{2} h_{11}(\mathbf{z}) + k h_2(\mathbf{z}) \\ \text{Frob}(\mathbb{L}_{k,3}) &= 2 \binom{k+1}{4} h_{111}(\mathbf{z}) + 3 \binom{k+1}{3} h_{21}(\mathbf{z}) + \binom{k+1}{2} h_3(\mathbf{z}) \\ \text{Frob}(\mathbb{L}_{k,4}) &= 5 \binom{k+2}{6} h_{1111}(\mathbf{z}) + 10 \binom{k+2}{5} h_{211}(\mathbf{z}) + 4 \binom{k+2}{4} h_{31}(\mathbf{z}) \\ &\quad + 2 \binom{k+2}{4} h_{22}(\mathbf{z}) + \binom{k+2}{3} h_4(\mathbf{z}) \end{aligned}$$

Proof of (5.6). An extension of the ‘‘cyclic lemma’’ of [1] gives a combinatorial proof of Identity (5.6). Consider the action of \mathbb{S}_n , on pairs (ℓ', β') in $\mathbb{L}_{k-1,n} \times \mathbb{P}_{k-1,n-1}$, that

permutes labels in ℓ' (as in Section 2). Assume that α' is the underlying path of ℓ' . Since ℓ' and β are independent, in view of (2.5), the corresponding Frobenius characteristic may be expressed as

$$\text{Frob}(\mathbb{L}_{k-1,n} \times \mathbb{P}_{k-1,n-1})(\mathbf{z}) = h_n[k \mathbf{z}] \binom{n+k-2}{k-1}. \quad (5.9)$$

We are going to show that there is an equivalence relation on the set $\mathbb{L}_{k-1,n} \times \mathbb{P}_{k-1,n-1}$, whose equivalence classes are all of size k , each of which containing one and only one labelled polyomino. Furthermore, for all pairs of a given class, the set partition associated to the labelled path are all the same. Hence, we may conclude that $\text{Frob}(\mathbb{L}_{k,n})$ is equal to the right hand-side of (5.9) divided by k , as formulated in (5.6).




FIGURE 8. There is only one polyomino in the equivalence class of a pair (ℓ, β) .

To this end, we work with slightly modified versions, denoted by α and β , of the paths α' and β' . On one hand, α is obtained by simply adding an east step at the end of α' . On the other hand, we get β by first shifting β' to the right (sending each vertex (a, b) to $(a + 1, b)$), and then adding a final north step. Hence α now goes from $(0, 0)$ to (k, n) , while β goes from $(1, 0)$ to (k, n) . We further denote by ℓ the labelled path version of ℓ' , which is directly obtained by carrying over the labels of α' to α . This makes sense since the only difference between the two is the perforce unlabeled final step. We underline that by construction, ℓ and β end up at the same point, with a final east step for ℓ and a final north step for β .

We now consider the two bi-infinite (labelled in one case) paths, denoted by ℓ^∞ and β^∞ , obtained by repeatedly gluing together copies of the paths ℓ and β at both of their ends. This is illustrated in Figure 9. Two pairs (ℓ'_1, β'_1) and (ℓ'_2, β'_2) , in $\mathbb{L}_{k-1,n} \times \mathbb{P}_{k-1,n-1}$, are considered to be equivalent if they give rise to the same pair of bi-infinite paths, i.e. $(\ell_1^\infty, \beta_1^\infty)$ is equal to $(\ell_2^\infty, \beta_2^\infty)$ up to a shift of the origin. Equivalent pairs correspond to specific portions of the pair of infinite paths. These portions are precisely characterized by points (x, y) where the two paths intersect in such a way that the ℓ^∞ reaches (x, y) by an east step, and β^∞ by a north step. We then say that ℓ^∞ and β^∞ have a *transversal intersection* at (x, y) . Equivalent pairs correspond to portions of ℓ^∞ and β^∞ both lying in the rectangle having south west corner $(x - k, y - n)$ and north east corner (x, y) , for transversal intersection points $p = (x, y)$. We denote by ℓ_p and β_p these portions.

As shown in [1] the two paths ℓ^∞ and β^∞ have exactly k transversal intersections. To finish our proof, we need only observe that, for every transversal intersection point p , the set partition of ℓ_p is equal to that of ℓ . Indeed, the big steps of ℓ_p are simply a cyclic shift of those of ℓ , together with their labels. \square




FIGURE 9. The k transversal intersections of ∞ -iterates of binomial paths.

Remark 3. It is interesting to compare Formula (5.7), evaluated at $k = rn$, with the conjectured expression [4, Formula (17)] for the Frobenius characteristic of the space of trivariate generalized diagonal harmonics. Recall that, up to a sign twist, this expression takes the form

$$\sum_{\lambda \vdash n} (rn + 1)^{\ell(\lambda) - 2} \prod_{j \in \lambda} \binom{(r+1)j}{j} \frac{p_\lambda(\mathbf{z})}{z_\lambda}. \quad (5.10)$$

It has been shown in [6] that this formula gives the Frobenius characteristic for the action of the symmetric group on labelled intervals in the r -Tamari posets. Our point here is that the difference

$$\sum_{\lambda \vdash n} (rn)^{\ell(\lambda) - 1} \binom{(r+1)n - 2}{n - 1} \frac{p_\lambda(\mathbf{z})}{z_\lambda} - \sum_{\lambda \vdash n} (rn + 1)^{\ell(\lambda) - 2} \prod_{j \in \lambda} \binom{(r+1)j}{j} \frac{p_\lambda(\mathbf{z})}{z_\lambda} \quad (5.11)$$

turns out to be h -positive. In terms of the corresponding \mathbb{S}_n -modules, this implies that the latter is contained in the former as a “permutation” \mathbb{S}_n -module. In fact, more appears to be true as discussed in Remark 7.

6. DOUBLY LABELLED POLYOMINOES

We may further add labels to horizontal steps of the lower path, and get an action of the product group $\mathbb{S}_k \times \mathbb{S}_n$ on the resulting set $\mathbb{L}_{k,n}^{(2)}$ of doubly labelled parallelogram polyominoes. We observe experimentally that the associated Frobenius characteristic is

given by the formula

$$\begin{aligned} \text{Frob}(\mathbb{L}_{k,n}^{(2)})(\mathbf{y}, \mathbf{z}) &= \frac{1}{n-1} h_k[(n-1)\mathbf{y}] h_n[k\mathbf{z}] + \frac{1}{k-1} h_k[n\mathbf{y}] h_n[(k-1)\mathbf{z}] \\ &\quad - \frac{n+k-1}{(n-1)(k-1)} h_k[(n-1)\mathbf{y}] h_n[(k-1)\mathbf{z}]. \end{aligned} \quad (6.1)$$

Observe the evident symmetry

$$\text{Frob}(\mathbb{L}_{k,n}^{(2)})(\mathbf{y}, \mathbf{z}) = \text{Frob}(\mathbb{L}_{n,k}^{(2)})(\mathbf{z}, \mathbf{y}), \quad (6.2)$$

which corresponds to exchanging the coordinates in our polyominoes. For example, we have

$$\text{Frob}(\mathbb{L}_{3,2}^{(2)})(\mathbf{y}, \mathbf{z}) = (6 s_3(\mathbf{y}) + 3 s_{21}(\mathbf{y})) s_2(\mathbf{z}) + (3 s_3(\mathbf{y}) + s_{21}(\mathbf{y})) s_{11}(\mathbf{z}).$$

Here, the $s_\mu(\mathbf{y})$ encode irreducibles of \mathbb{S}_k , whereas the $s_\lambda(\mathbf{z})$ encode those of \mathbb{S}_n . Naturally, our previous formula (5.6) appears as the multiplicity of the trivial representation of \mathbb{S}_k in (6.1). More precisely, recalling that

$$\langle h_k[n\mathbf{y}], h_k(\mathbf{y}) \rangle = \binom{n+k-1}{k},$$

we have

$$\begin{aligned} \text{Frob}(\mathbb{L}_{k,n})(\mathbf{z}) &= \langle \text{Frob}(\mathbb{L}_{k,n}^{(2)})(\mathbf{y}, \mathbf{z}), h_k(\mathbf{y}) \rangle \\ &= \frac{1}{n-1} \binom{n+k-2}{k} h_n[k\mathbf{z}] \\ &\quad + \left(\frac{1}{k-1} \binom{n+k-1}{k} - \frac{n+k-1}{(k-1)(n-1)} \binom{n+k-2}{k} \right) h_n[(k-1)\mathbf{z}] \\ &= \frac{1}{k} \binom{n+k-2}{k-1} h_n[k\mathbf{z}], \end{aligned}$$

since the coefficient of $h_n[(k-1)\mathbf{z}]$ vanishes.

It also follows, directly from (6.1), that the total number of doubly labelled parallelogram polyominoes in $\mathbb{L}_{k,n}^{(2)}$ is

$$|\mathbb{L}_{k,n}^{(2)}| = k^n (n-1)^{k-1} + (k-1)^{n-1} n^k - (k-1)^{n-1} (n-1)^{k-1} (n+k-1). \quad (6.3)$$

Considering \mathbb{S}_k -isotypic components of $\mathbb{L}_{k,n}^{(2)}$, we sometimes obtain nice formulas for the Frobenius characteristic of the resulting \mathbb{S}_n -module. For instance, denoting by ρ the rectangular partition r^n , we find that the coefficient of $s_\rho(\mathbf{y})$ in $\mathbb{L}_{rn,n}^{(2)}(\mathbf{y}, \mathbf{z})$ is

$$\mathbb{L}_{rn,n}^{(2)}(\mathbf{y}, \mathbf{z}) \Big|_{s_\rho(\mathbf{y})} = \frac{1}{rn-1} h_n[(rn-1)\mathbf{z}]. \quad (6.4)$$

See remark 6 for a q -analog of this formula.

We also have an alternative way to doubly label parallelogram polyominoes, which has a meaning when such labelled objects encode stable configurations for the sandpile model associated to the complete bipartite graph $K_{k,n}$ [9]. In this setting, we fix the label of the

first step of the blue path to be 1 (in words of the sandpile model, this step corresponds to the sink). With this convention, we may let $\mathbb{S}_n \times \mathbb{S}_{k-1}$ act on the labels, and denote by $\text{Frob}(\mathbb{L}_{k,n}^{(2,*)})$ the corresponding Frobenius characteristic.

Proposition 4. *We have:*

$$\text{Frob}(\mathbb{L}_{k,n}^{(2,*)}) = \frac{1}{k} h_n[k \mathbf{z}] h_{k-1}[n \mathbf{y}] \quad (6.5)$$

which implies that

$$|\mathbb{L}_{k,n}^{(2,*)}| = k^{n-1} n^{k-1}, \quad (6.6)$$

the number of parking functions (or of recurrent configurations, or of spanning trees) of $K_{k,n}$.

Proof. Take the proof of (5.6) and just add labels on the horizontal steps (but one) of the blue path. \square

7. OPERATORS AND MACDONALD POLYNOMIALS

To go on with our presentation, we need to recall a few notions related to the theory of Macdonald polynomials, and operators for which they are joint eigenfunctions. The *integral form (modified) Macdonald polynomials*⁹ $H_\mu(\mathbf{z}; q, t)$, with μ a partition of n , expand as

$$H_\mu(\mathbf{z}; q, t) = \sum_{\lambda \vdash n} K_{\lambda, \mu}(q, t) s_\lambda(\mathbf{z}),$$

where the $K(q, t)$'s are certain rational fractions in q, t , normalized so that $K_{n, \mu} = 1$. They are entirely characterized by this normalization together with the two triangularity properties

- (1) $H_\mu[\mathbf{z}(1 - q)]$ is in the span of the $s_\lambda(\mathbf{z})$, for $\lambda \succeq \mu$ in dominance order,
- (2) $H_\mu[\mathbf{z}(1 - t)]$ is in the span of the $s_\lambda(\mathbf{z})$, for $\lambda \succeq \mu'$ in dominance order.

It may be worth recalling that we have the special case

$$H_n(\mathbf{z}; q, t) = \prod_{i=1}^n (1 - q^i) h_n \left[\frac{\mathbf{z}}{1 - q} \right]. \quad (7.1)$$

Observe that the right hand side is independent of t . This is helpful when we specialize the H_μ 's at $t = 1$, since in that case we have the following multiplicative property

$$H_\mu(\mathbf{z}; q, 1) = \prod_{i=1}^r H_{\mu_i}(\mathbf{z}; q, 1), \quad \text{for} \quad \mu = \mu_1 \mu_2 \cdots \mu_r. \quad (7.2)$$

Thus, we can evaluate this last right hand side using (7.1). Observe also that this states that $H_\mu(\mathbf{z}; q, 1)$ is proportional to $h_\mu[\mathbf{z}/(1 - q)]$.

⁹In previous papers, the notation \tilde{H}_μ is often used for these, to distinguish them from other polynomials that are not used here.

Another often used specialization is at $t = 1/q$. In this case we have (see [10])

$$H_\mu(\mathbf{z}; q, 1/q) = q^{-n(\mu)} \prod_{c \in \mu} (1 - q^{\text{hook}(c)}) s_\mu \left[\frac{\mathbf{z}}{1 - q} \right], \quad (7.3)$$

where $n(\mu) = \sum_i (i-1)\mu_i$, the sum is taken over the cells c in the Ferrers diagram of μ , and $\text{hook}(c)$ is the hook-length of the cell c in μ , *i.e.* the number of cells of μ above c in the same column or to its right in the same row, including c itself.

Operators. For a symmetric function f , we consider the linear operator Δ_f , on symmetric functions, such that

$$\Delta_f(H_\mu) := f[B_\mu(q, t)] H_\mu, \quad (7.4)$$

where $f[B_\mu]$ stands for the evaluation of f in the $q^i t^j$, with (i, j) varying in the set of cells of μ . It is customary to denote by ∇ the operator Δ_{e_n} restricted to symmetric functions of degree n . Observe that the operator Δ_{h_k} (resp. Δ_{e_k}) appears as the coefficient of u^k in the linear operator Φ such that

$$\Phi(H_\mu) = \left(\prod_{(i,j) \in \mu} \frac{1}{1 - q^i t^j u} \right) H_\mu, \quad \text{resp.} \quad \Psi(H_\mu) = \left(\prod_{(i,j) \in \mu} (1 + q^i t^j u) \right) H_\mu. \quad (7.5)$$

One of the interests of the operator ∇ is that it allows a very compact formulation of the Frobenius characteristic of the \mathbb{S}_n -module of bivariate diagonal hamonic polynomials, in the form $\nabla(e_n(\mathbf{w}))$ (see [15] for more on this).

We will also denote $\tilde{\Delta}_f$ (resp. $\bar{\Delta}_f$) the specialization at $t = 1$ (resp $t = 1/q$) of the operator Δ_f . It turns out that the operators $\tilde{\nabla}$, $\tilde{\Phi}$, and $\tilde{\Psi}$ are multiplicative, since the associated eigenvalues are multiplicative. In view of our previous observations (see (7.1)), the set

$$\{h_\mu[\mathbf{z}/(1-q)] \mid \mu \vdash n\} \quad (7.6)$$

constitute a basis of eigenfunctions for the operators $\tilde{\Delta}_f$.

Similarly, from (7.3), it follows that the $s_\mu[\mathbf{z}/(1-q)]$'s are eigenfunctions of $\bar{\Delta}_f$, since they are proportional to the $H_\mu(\mathbf{z}; q, 1/q)$'s. The corresponding eigenvalues are $f[B_\mu(q, 1/q)]$. Observe that in the case of hook shapes, *i.e.* $\mu = s_{(n-r), 1^r}$, these are simply

$$f[B_{(n-r), 1^r}(q, 1/q)] = f[q^{-r}[n]_q]. \quad (7.7)$$

More operators. Recall from [14] that we may introduce symmetric functions $E_{n,r}$ by means of the plethystic identity

$$e_n \left[\mathbf{z} \frac{1-x}{1-q} \right] = \sum_{r=1}^n \frac{(x; q)_r}{(q; q)_r} E_{n,r}(\mathbf{z}). \quad (7.8)$$

In particular, setting $z = q$, we find that

$$e_n(\mathbf{z}) = \sum_{r=1}^n E_{n,r}(\mathbf{z}). \quad (7.9)$$

These play a key role in a generalized version of the ‘‘Shuffle Conjecture’’, together with another family of linear operators on symmetric functions defined as follows. For each integer a , and any symmetric function $f(\mathbf{z})$, we set

$$C_a f(\mathbf{z}) = (-q)^{1-a} f\left[\mathbf{z} - \frac{q-1}{qx}\right] \left(\sum_{k=0}^{\infty} h_k(\mathbf{z}) x^k\right) \Big|_{x^a}. \quad (7.10)$$

For example, we have

$$\begin{aligned} C_a(1) &= (-q)^{1-a} s_a(\mathbf{z}), & C_a(s_1(\mathbf{z})) &= (-q)^{1-a} (s_{a,1}(\mathbf{z}) + q^{-1} s_{a+1}(\mathbf{z})), & \text{and} \\ C_a(s_2(\mathbf{z})) &= (-q)^{1-a} (s_{a,2}(\mathbf{z}) + q^{-1} s_{a+1,1}(\mathbf{z}) + q^{-2} s_{a+2}(\mathbf{z})), & \text{if } a &\geq 2. \end{aligned}$$

As shown in [14], the operators C_a satisfy the commutativity relations

$$q(C_b C_a + C_{a-1} C_{b+1}) = C_a C_b + C_{b+1} C_{a-1}, \quad \text{whenever } a > b. \quad (7.11)$$

Recall also that, for a composition of n into r parts, we may consider the symmetric functions recursively defined as

$$E_\gamma(\mathbf{z}) := \begin{cases} C_a(E_\nu(\mathbf{z})) & \text{if } \gamma = a\nu, \\ 1 & \text{if } \gamma = 0, \end{cases} \quad (7.12)$$

where we denote by 0 the empty composition; and, we have

$$E_{n,r}(\mathbf{z}) = \sum_{\gamma \in \text{Comp}(n,r)} E_\gamma(\mathbf{z}). \quad (7.13)$$

Remark 4. An explicit enumerative description of $\nabla(E_\gamma(\mathbf{z}))$ is conjectured, in [14, Conj. 4.5], in terms of parking functions whose underlying Dyck path has prescribed returns to the diagonal. It appears that a similar statement holds for $\nabla^r(E_\gamma(\mathbf{z}))$, with the heights of returns to the diagonal specified by the composition γ . This is what we call the *r-generalized shuffle conjecture*. In the particular case of $\gamma = n$, since $E_n(\mathbf{z}) = (-q)^{-n+1} h_n(\mathbf{z})$, this conjecture implies that $\omega \tilde{\nabla}^r((-q)^{-n+1} h_n(\mathbf{z}))$ is the q -graded Frobenius characteristic of the \mathbb{S}_n -module generated by r -parking functions whose underlying r -Dyck path never returns to the ‘‘diagonal’’ (of slope $1/r$), except at its endpoints.

Observe that the commutativity relation (7.11) implies that, for any length r composition γ of n , we may express $E_\gamma(\mathbf{z})$ as a linear combination of the $E_\mu(\mathbf{z})$'s, for μ varying in the set of partitions of n having r parts. These last symmetric functions are in fact proportional to a specialization of Macdonald polynomials, that is

$$E_\mu(\mathbf{z}) = q^{-n(\mu)} (-1/q)^{n-r} H_{\mu'}(\mathbf{z}; q, 0), \quad (7.14)$$

where, as usual, $n(\mu)$ stands for $\sum_{i=1}^r (i-1)\mu_i$. Notice that the right hand side of this last identity is expressed in terms of the conjugate partition μ' .

8. FORMULA FOR THE q -FROBENIUS CHARACTERISTIC OF THE LABELLED POLYOMINO MODULE

Michele D’Adderio has proposed [7] the following explicit formula for the q -weighted area counting version of the Frobenius characteristics

$$\text{Frob}(\mathbb{L}_{k+1,n})(\mathbf{z}; q) = \omega \tilde{\Delta}_{h_k}(e_n(\mathbf{z})). \quad (8.1)$$

Here $\tilde{\Delta}_{h_k}$ stands the specialization at $t = 1$ of the operator Δ_{h_k} . This implies that $\tilde{\Delta}_{h_k}(e_n(\mathbf{z}))$ is e -positive, i.e. with coefficients in $\mathbb{N}[q]$ when expanded in the basis of elementary symmetric functions. In generating function format, we may reformulate (8.1) as

$$\begin{aligned} \sum_{k \geq 1} \text{Frob}(\mathbb{L}_{k,n})(\mathbf{z}; q) u^k &= \omega(u \tilde{\Phi}(e_n(\mathbf{z}))) \\ &= \omega \sum_{\mu \vdash n} f_\mu[1 - q] \prod_{i=1}^{\ell(\mu)} \frac{u h_{\mu_i}[\mathbf{z}/(1 - q)]}{(1 - u) \cdots (1 - u q^{\mu_i - 1})} \end{aligned}$$

where we denote by $f_\mu(\mathbf{z})$ the *forgotten* symmetric functions, that are dual to the elementary symmetric functions. Small values are as follows:

$$\begin{aligned} \sum_{k \geq 1} \text{Frob}(\mathbb{L}_{k,1})(\mathbf{z}; q) u^k &= \frac{u}{1 - u} h_1(\mathbf{z}), \\ \sum_{k \geq 1} \text{Frob}(\mathbb{L}_{k,2})(\mathbf{z}; q) u^k &= \frac{u}{(1 - u)(1 - qu)} \left(h_2(\mathbf{z}) + \frac{u}{1 - u} h_{11}(\mathbf{z}) \right), \\ \sum_{k \geq 1} \text{Frob}(\mathbb{L}_{k,3})(\mathbf{z}; q) u^k &= \frac{u}{(1 - u)(1 - uq)(1 - uq^2)} \\ &\quad \left(h_3(\mathbf{z}) + \frac{u(2 + q)}{1 - u} h_{21}(\mathbf{z}) + \frac{u^2(1 + q)}{(1 - u)^2} h_{111}(\mathbf{z}) \right). \end{aligned}$$

Even if Formula (8.1) is still conjectural, the following two formulas should help understand how $\omega \Delta_{h_k}(e_n(\mathbf{z}))$ could be interpreted as a bivariate Frobenius characteristic for labelled polyominoes, with a second statistic accounting for the parameter t , on top of q which already accounts for area. This also confirms indirectly that (8.1) should hold.

Proposition 5. *The q -free component of $\omega \Delta_{h_k}(e_n(\mathbf{z}))$ is given by the following formula*

$$\omega \Delta_{h_k}(e_n(\mathbf{z}))|_{q=0} = \frac{1}{t^k} \left(h_n \left[\mathbf{z} \frac{1 - t^{k+1}}{1 - t} \right] - h_n \left[\mathbf{z} \frac{1 - t^k}{1 - t} \right] \right). \quad (8.2)$$

Moreover, we have

$$\omega \Delta_{h_{k-1}}(e_n(\mathbf{z}))|_{t=1/q} = \frac{q^{n+k-nk-1}}{[k]_q} h_n \left[\mathbf{z} \frac{1 - q^k}{1 - q} \right] \begin{bmatrix} n + k - 2 \\ k - 1 \end{bmatrix}_q \quad (8.3)$$

which is a direct q -analog of (5.6).

For a combinatorial interpretation of the right hand side of (8.2), see Remark 5.

Proof. To prove (8.2), we first observe that $\Delta_{h_k}(e_n(\mathbf{z}))$ is symmetric in q and t , we may thus exchange their role. We may also remove ω , so that we need only show that

$$\widehat{\Delta}_{h_k}(e_n(\mathbf{z})) = \frac{1}{q^k} \left(e_n \left[\mathbf{z} \frac{1-q^{k+1}}{1-q} \right] - e_n \left[\mathbf{z} \frac{1-q^k}{1-q} \right] \right), \quad (8.4)$$

where $\widehat{\Delta}_{h_k} := \Delta_{h_k}|_{t=0}$. Evaluating (7.8) at $z = q^{k+1}$, we get

$$e_n \left[\mathbf{z} \frac{1-q^{k+1}}{1-q} \right] = \sum_{r=1}^n \begin{bmatrix} k+r \\ r \end{bmatrix}_q E_{n,r}(\mathbf{z}). \quad (8.5)$$

Using (8.5), and well known properties of q -binomial coefficients, we can calculate that the right hand side of (8.4) is

$$\widehat{\Delta}_{h_k}(e_n(\mathbf{z})) = \sum_{r=1}^n \frac{1}{q^k} \left(\begin{bmatrix} k+r \\ r \end{bmatrix}_q - \begin{bmatrix} k+r-1 \\ r \end{bmatrix}_q \right) E_{n,r}(\mathbf{z}) \quad (8.6)$$

$$= \sum_{r=1}^n \begin{bmatrix} k+r-1 \\ r-1 \end{bmatrix}_q E_{n,r}(\mathbf{z}). \quad (8.7)$$

Now, recalling (7.9), this last equation readily follows from the surprising fact that the $E_{n,r}(\mathbf{z})$'s are eigenfunctions of the operator $\widehat{\Delta}_{h_k}$. Indeed, as we will see, we have

$$\widehat{\Delta}_{h_k} E_{n,r}(\mathbf{z}) = \begin{bmatrix} k+r-1 \\ r-1 \end{bmatrix}_q E_{n,r}(\mathbf{z}). \quad (8.8)$$

This follows from the even stronger fact that for all compositions γ , having r parts, we have

$$\widehat{\Delta}_{h_k} E_\gamma(\mathbf{z}) = \begin{bmatrix} k+r-1 \\ r-1 \end{bmatrix}_q E_\gamma(\mathbf{z}). \quad (8.9)$$

In view of the observation following (7.13), we need only show that (8.9) holds for $E_\mu(\mathbf{z})$, with μ a length r partition. Since we have already seen recalled (see (7.14)) that these E_μ 's are essentially specializations of the H_μ 's, it only remains to check that

$$\widehat{\Delta}_{h_k} H_{\mu'}(\mathbf{z}; q, 0) = \begin{bmatrix} k+r-1 \\ r-1 \end{bmatrix}_q H_{\mu'}(\mathbf{z}; q, 0).$$

But, by definition of Δ_{h_k} , this is equivalent to the classical fact that

$$\begin{bmatrix} k+r-1 \\ r-1 \end{bmatrix}_q = h_k \left[\frac{1-q^r}{1-q} \right].$$

This ends our proof of (8.2).

To prove (8.3), we start by recalling from above (see (7.3)) that the $s_\mu[\mathbf{z}/(1-q)]$'s are eigenfunctions of $\overline{\Delta}_{h_{k-1}} = \Delta_{h_{k-1}}|_{t=1/q}$, with eigenvalue $h_{k-1}[B_\mu(q, 1/q)]$. Once again, we

remove ω , to turn the identity we have to show into

$$\overline{\Delta}_{h_{k-1}}(e_n(\mathbf{z})) = \frac{q^{n+k-nk-1}}{[k]_q} e_n \left[\mathbf{z} \frac{1-q^k}{1-q} \right] \begin{bmatrix} n+k-2 \\ k-1 \end{bmatrix}_q. \quad (8.10)$$

Cauchy's identity implies that

$$e_n \left[\mathbf{z} \frac{1-q^k}{1-q} \right] = \sum_{\mu \vdash n} s_{\mu'} [1-q^k] s_{\mu} \left[\frac{\mathbf{z}}{1-q} \right].$$

Since $s_{\nu}[1-u]$ is only non-zero when ν is a hook shape, and $s_{(n-r),1^r}[1-u] = (-u)^r(1-u)$, we may rewrite the above identity as as

$$e_n \left[\mathbf{z} \frac{1-q^k}{1-q} \right] = \sum_{r=0}^{n-1} (1-q^k) (-q^k)^{n-r-1} s_{(n-r),1^r} \left[\frac{\mathbf{z}}{1-q} \right]. \quad (8.11)$$

Using this identity at $k=1$, we may expand $e_n(\mathbf{z})$ in terms of eigenfunctions for the operator $\overline{\Delta}_{h_{k-1}}$, and thus we calculate (using (7.7)) that

$$\begin{aligned} \overline{\Delta}_{h_{k-1}}(e_n(\mathbf{z})) &= \overline{\Delta}_{h_{k-1}} \sum_{r=0}^{n-1} (1-q) (-q)^{n-r-1} s_{(n-r),1^r} \left[\frac{\mathbf{z}}{q-1} \right] \\ &= \sum_{r=0}^{n-1} (1-q) (-q)^{n-r-1} q^{-r(k-1)} h_{k-1} \left[\frac{1-q^n}{1-q} \right] s_{(n-r),1^r} \left[\frac{\mathbf{z}}{q-1} \right] \\ &= \begin{bmatrix} n+k-2 \\ k-1 \end{bmatrix}_q \sum_{r=0}^{n-1} (-1)^{n-r-1} (1-q) q^{n-rk-1} s_{(n-r),1^r} \left[\frac{\mathbf{z}}{q-1} \right] \\ &= \frac{q^{n+k-nk-1}}{[k]_q} \begin{bmatrix} n+k-2 \\ k-1 \end{bmatrix}_q \sum_{r=0}^{n-1} (1-q^k) (-q^k)^{n-r-1} s_{(n-r),1^r} \left[\frac{\mathbf{z}}{q-1} \right] \end{aligned}$$

which shows (8.10), using (8.11) to evaluate the sum. \square

Remark 5. The right-hand side of (8.2) may be interpreted as a graded Frobenius characteristic of ribbon shaped labeled parallelogram polyominoes of size $n \times k$. The needed grading corresponds to the dinv statistic for unlabeled parallelogram polyominoes, as defined in [2]. Recall that, in the \mathbb{A} -word encoding $w = w_1 w_2 \cdots w_k$ of a parallelogram polyomino (see Figure 5), the dinv statistic is the number of pairs (w_i, w_j) , with $i < j$ and j is the successor of i in \mathbb{A} . For any ribbon-shaped labeled polyomino, we may simply set its dinv -statistic to be equal to that of the underlying parallelogram polyomino. With this definition, we observe that the dinv -value coincides with the area below the lower path of the polyomino. This readily implies that

$$\left(\text{Frob}(\mathbb{L}_{k,n})(\mathbf{z}; q) \right) \Big|_{q=0} = t^n \left(h_n \left[\mathbf{z} \frac{1-t^k}{1-t} \right] - h_n \left[\mathbf{z} \frac{1-t^{k-1}}{1-t} \right] \right). \quad (8.12)$$

Remark 6. Experiments suggest that we have the following area enumerating analog of Formula (6.4)

$$\mathbb{L}_{r,n}^{(2)}(\mathbf{y}, \mathbf{z}; q) \Big|_{s_{\rho}(\mathbf{y})} = \omega \tilde{\nabla}^r ((-q)^{-n+1} h_n), \quad (8.13)$$

where ρ is the partition consisting of n rows of size r . In fact, this identity follows from the r -generalized shuffle conjecture (see Remark 4), considering paths that never come back to the diagonal of slope $1/r$. It would be interesting to describe all isotypic components of $\mathbb{L}_{rn,n}^{(2)}$ in this manner.

Toward a general divv statistic. A divv-statistic¹⁰ on general labelled polyominoes would make possible an analog of the ‘‘Shuffle Conjecture’’ of [13], in the form:

$$\Delta_{h_k}(e_n(\mathbf{z})) = \sum_{\pi} q^{\text{area}(\pi)} t^{\text{divv}(\pi)} \mathbf{z}_{\pi}, \quad (8.14)$$

with π varying in the set $\mathbb{L}_{k,n}$. More explicitly, this means that we would consider more general labellings of polyominoes; with labels in \mathbb{N} and allowing for repetition of labels, while keeping the strictly increasing condition on runs of vertical steps. In the above expression, \mathbf{z}_{π} denotes the product of the variables z_i , for i running through all labels (with multiplicities) of π . In a sense, this should be a natural extension of the original Shuffle Conjecture, since it appears (observed experimentally) that

$$\Delta_{h_{r_{n-1}}}(e_n(\mathbf{z})) - q^{\binom{n-1}{2}} \nabla^r(e_n(\mathbf{z})) \quad \text{lies in} \quad \mathbb{N}[q, t] \{s_{\mu} | \mu \vdash n\}. \quad (8.15)$$

As we have already seen (see Remark 2), parking functions can be considered as special cases of labelled parallelogram polymominoes, making (8.15) a natural phenomenon.

Remark 7. Explicit computer calculations suggest that even more seems to hold. Indeed, up to the same multiplicative factor of $q^{\binom{n-1}{2}}$, the difference between $\Delta_{h_{n-1}}(e_n(\mathbf{z}))$ and the ‘‘Shuffle-like Conjecture Formula’’ for the Frobenius characteristic of the space of trivariate diagonal harmonics (see [4, Formula (15)]), appears to always be Schur-positive (with coefficients in $\mathbb{N}[q, t]$). Not only is this stronger than (8.15), but it suggest that a third statistic might play a role here.

Remark 8. Along these lines, it is interesting to observe that, when $k \leq n$,

$$q^{kn - \binom{k+1}{2}} \overline{\Delta}_{e_k}(e_n(\mathbf{z})) = \frac{1}{[k+1]_q} \begin{bmatrix} n \\ k \end{bmatrix}_q e_n \left[\mathbf{z} \frac{q^{k+1} - 1}{q - 1} \right]. \quad (8.16)$$

Hence, the Frobenius characteristic (2.8) may be expressed as

$$\text{Frob}(\mathbb{L}_{k,n})(\mathbf{z}; q) = q^{kn - \binom{k+1}{2}} [k+1]_q \begin{bmatrix} n \\ k \end{bmatrix}_q^{-1} \omega \overline{\Delta}_{e_k}(e_n(\mathbf{z})). \quad (8.17)$$

Observe the similarity between this expression and (8.1).

The bounce statistic. Angela Hicks [2] has recently given a nice bijection from which it follows that we have the following explicit formula due to Garsia and D’Adderio

$$\mathbb{P}_{k,n}(q, t) := \sum_{\pi \in \mathbb{P}_{k,n}} q^{\text{area}(\pi)} t^{\text{b}(\pi)} = \langle \nabla(e_{k+n-2}), h_{k-1} h_{n-1} \rangle \quad (8.18)$$

¹⁰Which we do not know how to define yet.

where $b(\pi)$ denotes the *bounce* statistic of a polyomino π , introduced in [9]. In other words, up to a multiplicative factor, the right hand side is the coefficient of the monomial symmetric function of $m_{k-1,n-1}$ in the expansion of $\nabla(e_{k+n-2})$. It immediately follows that $\mathbb{P}_{k,n}(q,t)$ is symmetric, both in q,t and n,k , as already observed in [9]. Moreover, specializing t at $1/q$, we get the following direct q -analogs of (3.3)

$$q^{(k-1)(n-1)} \mathbb{P}_{k,n}(q, 1/q) = \frac{1}{[n+k]_q} \begin{bmatrix} n+k \\ n \end{bmatrix}_q \begin{bmatrix} n+k-2 \\ n-1 \end{bmatrix}_q \quad (8.19)$$

$$= q^{-(k-1)} s_{(k-1,k-1)}(1, q, q^2, \dots, q^n). \quad (8.20)$$

To prove this, we proceed as follows. Recall from [10] that

$$q^{\binom{n}{2}} \nabla(e_n) \Big|_{t \leftarrow 1/q} = \frac{1}{[n+1]_q} e_n[\mathbf{z}[n+1]_q],$$

so that (8.18) may be reformulated as

$$q^{\binom{n+k-2}{2}} \mathbb{P}_{k,n}(q, 1/q) = \left\langle \frac{1}{[n+k-1]_q} e_{n+k-2}[\mathbf{z}[n+k-1]_q], h_{k-1} h_{n-1} \right\rangle. \quad (8.21)$$

Now, Cauchy formula gives

$$e_{n+k-2}[\mathbf{z}[n+k-1]_q] = \sum_{\lambda \vdash n+k-2} s'_\lambda([n+k-1]_q) s_\lambda(\mathbf{z}),$$

leading us to

$$q^{\binom{n+k-2}{2}} \mathbb{P}_{k,n}(q, 1/q) = \frac{1}{[n+k-1]_q} e_{k-1}([n+k-1]_q) e_{n-1}([n+k-1]_q). \quad (8.22)$$

Finally, since

$$e_j([n]_q) = q^j \begin{bmatrix} n \\ j \end{bmatrix}_q,$$

we conclude that

$$q^{\binom{n+k-2}{2}} \mathbb{P}_{k,n}(q, 1/q) = \frac{1}{[n+k-1]_q} q^{k-1} \begin{bmatrix} n+k-1 \\ k-1 \end{bmatrix}_q q^{n-1} \begin{bmatrix} n+k-1 \\ n-1 \end{bmatrix}_q. \quad (8.23)$$

A direct calculation shows that this is equivalent to (8.19).

We cannot fail to notice that since the right hand side of (8.19) has potentially also a combinatorial interpretation, we should naturally wonder what that may be in this case.

This has been carried out recently by Angela Hicks and Emily Leven [16] who also were able to obtain a direct bijective proof of (8.19).

9. THANKS AND FUTURE CONSIDERATIONS

We would like to thank Vic Reiner for reminding us that we should consider doubly labelled polyominoes in the specific manner we describe here. We would also like to thank Nolan Wallach for putting us on the right track for the interpretation of parrallelogram polyominoes as indexing sets for SL_2 -invariants.

In our view, the main question that begs to be addressed in the future, seems to find a natural context in which $\Delta_{h_k}(e_n)$ would appear as the bigraded Frobenius characteristic of an \mathbb{S}_n -module similar to the \mathbb{S}_n -module of diagonal harmonic polynomials. This would help truly explain all our observations above.

REFERENCES

- [1] J.-C. AVAL, M. D'ADDERIO, M. DUKES AND Y. LE BORGNE, *The Sandpile Model on $K_{m,n}$ and a Cyclic Lemma*, submitted.
- [2] J.-C. AVAL, M. D'ADDERIO, M. DUKES, A. HICKS AND Y. LE BORGNE, *Statistics on parallelogram polyominoes and a q, t -analogue of the Narayana numbers*, J. of Combin. Theory Ser. A , Vol. 123 (2014), 271–286.
- [3] F. BERGERON, *Algebraic Combinatorics and Coinvariant Spaces*, CMS Treatise in Mathematics, CMS and A.K.Peters, 2009.
- [4] F. BERGERON, L.-F. PRÉVILLE-RATELLE, *Higher Trivariate Diagonal Harmonics via Generalized Tamari Posets*, J. of Combin. Vol. 3 (2012), 317–341.
- [5] M. BOUSQUET-MÉLOU, *Polyominoes and Polygons*, Jerusalem Combinatorics '93, Contemporary Mathematics, AMS, 55–70, 1994. (see also Habilitation 1996, [Bousquet-Mélou publications](#)).
- [6] M. BOUSQUET-MÉLOU, G. CHAPUY, L.-F. PRÉVILLE-RATELLE, *The representation of the symmetric group on m -Tamari intervals*, Adv. Math. 247 (2013) 309342.
- [7] M. D'ADDERIO, *Personnal communication*.
- [8] M.-P. DELEST, AND G. VIENNOT, *Algebraic Languages and Polyominoes Enumeration*, Theoret. Comput. Sci. 34, 169–206, 1984.
- [9] M. DUKES AND Y. LE BORGNE, *Parallelogram Polyominoes, the Sandpile Model on a Bipartite Graph, and a q, t -Narayana Polynomial*, J. of Combin. Theory Ser. A Volume 120 (2013) 816–842.
- [10] A.M. GARSIA AND M. HAIMAN, *A Remarkable q, t -Catalan Sequence and q -Lagrange inversion*, J. Algebraic Combin. **5** (1996), no. 3, 191–244.
- [11] I. GESSEL, *A Noncommutative Generalization and q -Analog of the Lagrange Inversion Formula*, Trans. Amer. Math. Soc. 257, 455–482, 1980.
- [12] I. GESSEL, AND X. G. VIENNOT, *Binomial determinants, paths, and hook length formulae*, Adv. in Math. 58, 300–321, 1985.
- [13] J. HAGLUND, M. HAIMAN, N. LOEHR, J. REMMEL, AND A. ULYANOV, *A Combinatorial Formula for the Character of the Diagonal Coinvariants*, Duke Math. J. Volume 126, Number 2 (2005), 195-232.
- [14] J. HAGLUND, J. MORSE, AND M. ZABROCKI, *A Compositional Shuffle Conjecture Specifying Touch Points of the Dyck Path*, Canad. J. Math. Vol. 64 (2012) 822–844.
- [15] M. HAIMAN, *Combinatorics, symmetric functions and Hilbert schemes*, In CDM 2002: Current Developments in Mathematics in Honor of Wilfried Schmid & George Lusztig, International Press Books (2003) 39–112.
- [16] A. HICKS, AND E. LEVEN *A refinement of the Shuffle Conjecture with cars of two sizes and $t = 1/q$* J. of Combinatorics, 5.1 (2014), 31–50.
- [17] I. G. MACDONALD, *Symmetric Functions and Hall polynomials*, second ed., Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1995, With contributions by A. Zelevinsky, Oxford Science Publications.
- [18] E. MILLER AND B. STURMFELS, *Combinatorial Commutative Algebra*, Graduate Texts in Mathematics **227**, Springer, 2005.
- [19] S. MUKAI, *An Introduction to Invariants and Moduli*, Cambridge University Press, 2003.

LABRI, CNRS, UNIVERSITÉ DE BORDEAUX, 351 COURS DE LA LIBÉRATION, BORDEAUX.

DÉPARTEMENT DE MATHÉMATIQUES, UQAM, C.P. 8888, SUCC. CENTRE-VILLE, MONTRÉAL, H3C 3P8, CANADA.

DEPARTMENT OF MATHEMATICS, UCSD.