

# IDEALS OF QUASI-SYMMETRIC FUNCTIONS AND SUPER-COINVARIANT POLYNOMIALS FOR $\mathcal{S}_n$

J.-C. AVAL, F. BERGERON, AND N. BERGERON

ABSTRACT. The aim of this work is to study the quotient ring  $\mathbf{R}_n$  of the ring  $\mathbb{Q}[x_1, \dots, x_n]$  over the ideal  $\mathcal{J}_n$  generated by non-constant homogeneous quasi-symmetric functions. This article is a sequel of [2], in which we investigated the case of infinitely many variables. We prove here that the dimension of  $\mathbf{R}_n$  is given by  $C_n$ , the  $n^{\text{th}}$  Catalan number. This is also the dimension of the space  $\mathbf{SH}_n$  of super-covariant polynomials, defined as the orthogonal complement of  $\mathcal{J}_n$  with respect to a given scalar product. We construct a basis for  $\mathbf{R}_n$  whose elements are naturally indexed by Dyck paths. This allows us to understand the Hilbert series of  $\mathbf{SH}_n$  in terms of number of Dyck paths with a given number of factors.

## 1. INTRODUCTION

We study, in this paper, a natural analog of the space  $\mathbf{H}_n$  of covariant polynomials of  $\mathcal{S}_n$ . Let  $X$  denote the  $n$  variables  $x_1, \dots, x_n$  and  $\mathbb{Q}[X]$  denote the ring of polynomials in the variables  $X$ . Let  $\mathcal{I}_n$  denote the ideal of  $\mathbb{Q}[X]$  generated by all symmetric polynomials with no constant term. That is

$$\mathcal{I}_n = \langle h_k(X), k > 0 \rangle,$$

where  $h_k(X)$  is the  $k^{\text{th}}$  homogeneous symmetric polynomials in the variables  $X$ . (*cf.* [18]). We consider the following scalar product on  $\mathbb{Q}[X]$ :

$$(1.1) \quad \langle P, Q \rangle = P(\partial X)Q(X)|_{X=0}$$

where  $\partial X$  stands for  $\partial x_1, \dots, \partial x_n$  and in the same spirit  $X = 0$  stands for  $x_1 = \dots = x_n = 0$ . The space  $\mathbf{H}_n$  is defined as the orthogonal complement, denoted by  $\mathcal{I}_n^\perp$ , of the ideal  $\mathcal{I}_n$  in  $\mathbb{Q}[X]$ .

Equivalently (*cf.* [11], Proposition I.2.3), covariant polynomials (also known as  $\mathcal{S}_n$ -harmonic polynomials) can be defined as polynomials  $P$  such that  $Q(\partial X)P = 0$ , for any symmetric polynomial  $Q$  with no constant term. Since elements of  $\mathbf{H}_n$  satisfy the Laplace equation

$$(\partial x_1^2 + \dots + \partial x_n^2) P = \Delta P = 0,$$

every covariant polynomial is also harmonic.

Classical results [1, 22] state that the space  $\mathbf{H}_n$  affords a graded  $\mathcal{S}_n$ -module structure and is isomorphic (as a representation of  $\mathcal{S}_n$ ) to the left regular representation.

---

*Date:* December 12, 2002.

F. Bergeron is supported in part by NSERC and FCAR.

N. Bergeron is supported in part by CRC, NSERC and PREA.

Furthermore, as a graded  $\mathcal{S}_n$ -module,  $\mathbf{H}_n$  is isomorphic to the quotient

$$Q_n = \mathbb{Q}[X]/\mathcal{I}_n.$$

The space  $Q_n$  appears naturally in other contexts; for instance, as the cohomology ring of the variety of complete flags [7]. The discussion above implies that

$$(1.2) \quad \dim \mathbf{H}_n = n!.$$

Part of the interesting results surrounding the study of  $\mathbf{H}_n$  involve the fact that it can also be described as the linear span of all partial derivatives of the Vandermonde determinant. This is a special case of a general result for finite groups generated by reflections [22].

By analogy, we consider here the space  $\mathbf{SH}_n = \mathcal{J}_n^\perp$  of *super-covariant* polynomials, where  $\mathcal{J}_n$  is the ideal generated by *quasi-symmetric* polynomials with no constant term. Since the ring of symmetric polynomials is a subring of the ring of quasi-symmetric polynomials, we have  $\mathcal{I}_n \subseteq \mathcal{J}_n$  hence  $\mathcal{J}_n^\perp \subseteq \mathcal{I}_n^\perp$ , thus

$$\mathbf{SH}_n \subseteq \mathbf{H}_n,$$

which justifies the terminology. Quasi-symmetric polynomials were introduced by Gessel in 1984 [13] and have since appeared as a crucial tool in many interesting algebraico-combinatorial contexts (*cf.* [6, 12, 19, 20, 21]).

As in the corresponding symmetric setup, we have a graded isomorphism

$$(1.3) \quad \mathbf{SH}_n \simeq \mathbf{R}_n = \mathbb{Q}[X]/\mathcal{J}_n$$

and the approach used in the following work concentrates on this alternate description. We construct a basis of  $\mathbf{R}_n$  by giving an explicit set of monomial representatives. This set is naturally indexed by *Dyck paths* of length  $n$ , hence we obtain the following main theorem.

**Theorem 1.1.** *The dimension of  $\mathbf{SH}_n$  is given by the well known Catalan numbers:*

$$(1.4) \quad \dim \mathbf{SH}_n = \dim \mathbf{R}_n = C_n = \frac{1}{n+1} \binom{2n}{n}.$$

*In fact, taking into account the grading (with respect to degree), we have the Hilbert series*

$$(1.5) \quad \sum_{k=0}^{n-1} \dim \mathbf{SH}_n^{(k)} t^k = \sum_{k=0}^{n-1} \frac{n-k}{n+k} \binom{n+k}{k} t^k.$$

The article contains five sections. In Section 2 we recall useful definitions and basic properties. In Section 3 we construct a family  $\mathcal{G}$  of generators for the ideal  $\mathcal{J}_n$  and state useful properties of this set. Section 4 is devoted to the proof of the first part of Theorem 1.1. We construct an explicit basis for  $\mathbf{R}_n$  which allows us in Section 5 to obtain the Hilbert series of  $\mathbf{SH}_n$ .

Before we start, let us remark that F. Hivert [15] has developed an action of the Hecke algebra on  $\mathbb{Q}[X]$  for which a polynomial is invariant if and only if it is quasi-symmetric. One way to reformulate his result is to consider the generators  $e_i = \frac{q-T_i}{(1+q)}$

of the Hecke algebra, where  $T_i$  are the standard generators and  $q$  is an arbitrary parameter. Then

$$(1.6) \quad e_i e_{i\pm 1} e_i - \frac{q}{(1+q)^2} e_i$$

acts, via Hivert's action, as zero on the polynomial ring. Hence, the quotient of the Hecke algebra by the relation (1.6), classically known as the Temperley-Lieb algebra  $TL_n(q)$  (cf. [16]), faithfully acts on polynomials. At  $q = 1$  this is a subalgebra of the symmetric group algebra. The ring of quasi-symmetric polynomials is then identified as the invariants  $\mathbb{Q}[X]^{TL_n(q)}$  of the algebra  $TL_n(q)$  for any value of  $q$ . The algebra  $TL_n(q)$  is known to have dimension equal to  $C_n$ . It is striking to discover in this context that the space  $\mathbf{R}_n$  above also has dimension  $C_n$ . In analogy to the classical covariant theory, one would like to see an action of  $TL_n(q)$  on  $R_n$  which would identify it as the left regular representation. Unfortunately, Hivert's action is not compatible with multiplication and does not preserve the ideal  $\mathcal{J}_n$ , hence it does not induce the desired action on the quotient. This observation raises many new avenues of investigation.

## 2. BASIC DEFINITIONS

A *composition*  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$  of a positive integer  $d$  is an ordered list of positive integers ( $> 0$ ) whose sum is  $d$ . We denote this by  $\alpha \models d$  and also say that  $\alpha$  is a composition of *size*  $d$  and denote this by  $|\alpha|$ . The integers  $\alpha_i$  are the *parts* of  $\alpha$ , and the length  $\ell(\alpha)$  is set to be the number of parts of  $\alpha$ . We denote by  $0$  the unique empty composition of size  $d = 0$ .

There is a natural one-to-one correspondence between compositions of  $d$  and subsets of  $\{1, 2, \dots, d-1\}$ . Let  $S = \{a_1, a_2, \dots, a_k\}$  be such a subset, with  $a_1 < \dots < a_k$ , then the composition associated to  $S$  is  $\alpha_d(S) = (a_1 - a_0, a_2 - a_1, \dots, a_{k+1} - a_k)$ , where we set  $a_0 := 0$  and  $a_{k+1} := d$ . We denote by  $D(\alpha)$  the set associated to  $\alpha$  through this correspondence. For compositions  $\alpha$  and  $\beta$ , we say that  $\beta$  is a *refinement* of  $\alpha$ , if  $D(\alpha) \subset D(\beta)$ , and denote this by  $\beta \succeq \alpha$ .

We use vector notation for monomials. More precisely, for  $\nu = (\nu_1, \dots, \nu_n) \in \mathbb{N}^n$ , we denote  $X^\nu$  the monomial

$$(2.1) \quad x_1^{\nu_1} x_2^{\nu_2} \cdots x_n^{\nu_n}.$$

For a polynomial  $P \in \mathbb{Q}[X]$ , we further denote  $[X^\nu]P(X)$  as the coefficient of the monomial  $X^\nu$  in  $P(X)$ .

For a *vector*  $\nu \in \mathbb{N}^n$ , let  $c(\nu)$  represent the composition obtained by erasing zeros (if any) in  $\nu$ . A polynomial  $P \in \mathbb{Q}[X]$  is said to be *quasi-symmetric* if and only if, for any  $\nu$  and  $\mu$  in  $\mathbb{N}^n$ , we have

$$[X^\nu]P(X) = [X^\mu]P(X)$$

whenever  $c(\nu) = c(\mu)$ . The space of quasi-symmetric polynomials in  $n$  variables is denoted by  $Qsym_n$ . The space  $Qsym_n^{(d)}$  of homogeneous quasi-symmetric polynomials of degree  $d$  admits as linear basis the set of *monomial* quasi-symmetric polynomials

indexed by compositions of  $d$ . More precisely, for each composition  $\alpha$  of  $d$  with at most  $n$  parts, we set

$$(2.2) \quad M_\alpha = \sum_{c(\nu)=\alpha} X^\nu.$$

For the 0 composition, we set  $M_0 = 1$ . Another important linear basis is that of the *fundamental* quasi-symmetric polynomials (cf. [13]):

$$(2.3) \quad F_\alpha = \sum_{\beta \succeq \alpha} M_\beta$$

with  $\alpha \models n$  and  $\ell(\alpha) \leq n$ . For example, with  $n = 4$ ,

$$\begin{aligned} F_{21}(x_1, x_2, x_3, x_4) &= M_{21}(x_1, x_2, x_3, x_4) + M_{1111}(x_1, x_2, x_3, x_4) \\ &= x_1^2 x_2 + x_1^2 x_3 + x_1^2 x_4 + x_2^2 x_3 + x_2^2 x_4 + x_3^2, x_4 \\ &\quad + x_1 x_2 x_3 + x_1 x_2 x_4 + x_1 x_3 x_4 + x_2 x_3 x_4. \end{aligned}$$

Part of the interest of fundamental quasi-symmetric functions comes from the following properties. The first is trivial, but very useful and the second comes from the theory of  $P$ -partitions [20, 21].

**Proposition 2.1.** For  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k) \models d$ ,

$$(2.4) \quad F_\alpha(X) = \begin{cases} x_1 F_{(\alpha_1-1, \alpha_2, \dots, \alpha_k)}(X) + F_\alpha(x_2, \dots, x_n) & \text{if } \alpha_1 > 1, \\ x_1 F_{(\alpha_2, \alpha_3, \dots, \alpha_k)}(x_2, \dots, x_n) + F_\alpha(x_2, \dots, x_n) & \text{if } \alpha_1 = 1. \end{cases}$$

Let  $u = u_1 \cdots u_\ell \in \mathcal{S}_\ell$  and  $v = v_1 \cdots v_m \in \mathcal{S}_{[\ell+1, \ell+m]}$ . Let  $u \omega v$  denote the set of *shuffles* of the words  $u$  and  $v$ , i.e.  $u \omega v$  is the set of all permutations  $w$  of  $\ell + m$  such that  $u$  and  $v$  are subwords of  $w$ . In particular  $u \omega v$  contains  $\binom{\ell+m}{m}$  permutations. Let  $\mathcal{D}(u) = \{i, u_i > u_{i+1}\}$  denote the *descent set* of  $u$ . If  $\beta$  and  $\gamma$  are the two compositions such that  $D(\beta) = \mathcal{D}(u)$  and  $D(\gamma) = \mathcal{D}(v)$ , then

**Proposition 2.2** ([21], Exercise 7.93).

$$(2.5) \quad F_\beta F_\gamma = \sum_{w \in u \omega v} F_{\alpha_{\ell+m}(D(w))}.$$

In (2.1), the monomials are in correspondence with vectors  $\nu \in \mathbb{N}^n$ . Just as for compositions, the size  $\nu_1 + \cdots + \nu_n$  of  $\nu$  is denoted by  $|\nu|$ . It is also convenient to denote by  $\ell(\nu)$  the position of its last non-zero component. As usual,  $\nu + \mu$  is the componentwise addition of vectors.

For ease of reading, we reserve the use of  $\alpha$ ,  $\beta$  and  $\gamma$  to represent compositions, and the other Greek letters to represent vectors. We use the same symbol  $\alpha$  for both the composition  $(\alpha_1, \dots, \alpha_\ell)$  and the word  $\alpha_1 \cdots \alpha_\ell$ , likewise for vectors. In general, the length of vectors (or number of variables) is fixed and equal to  $n$ . If  $w$  is a word of integers (that is an element of  $\mathbb{N}^k$  for  $0 \leq k \leq n$ ) we denote by  $w0^* = w0^{n-k}$  the vector whose first  $k$  parts are the *letters* of  $w$ , to which are added  $n - k$  zeros at the end. If  $u = u_1 \cdots u_k$  and  $v = v_1 \cdots v_m$  are words of integers, the word


$$uv := u_1 \cdots u_k v_1 \cdots v_m$$

is the *concatenation* of  $u$  and  $v$ .

We next associate to any vector  $\nu$  a path  $\pi(\nu)$  in the  $\mathbb{N} \times \mathbb{N}$  plane with steps going north or east as follows. If  $\nu = (\nu_1, \dots, \nu_n)$ , the path  $\pi(\nu)$  is


$$(0, 0) \rightarrow (\nu_1, 0) \rightarrow (\nu_1, 1) \rightarrow (\nu_1 + \nu_2, 1) \rightarrow (\nu_1 + \nu_2, 2) \rightarrow \dots \\ \rightarrow (\nu_1 + \dots + \nu_n, n - 1) \rightarrow (\nu_1 + \dots + \nu_n, n).$$

For example the path associated to  $\nu = (2, 1, 0, 3, 0, 1)$  is



Observe that the height of the path is always  $n$ , whereas its width is  $|\nu|$ .

We distinguish two kinds of paths, thus two kinds of vectors, with respect to their “behavior” with respect to the diagonal  $y = x$ . If the path remains above the diagonal, we call it a *Dyck path*, and say that the corresponding vector is *Dyck*. If not, we say that the path (or equivalently the associated vector) is *transdiagonal*. For example  $\eta = (0, 0, 1, 2, 0, 1)$  is Dyck and  $\varepsilon = (0, 2, 1, 0, 2, 2)$  is transdiagonal.



Observe that  $\nu = \nu_1 \cdots \nu_n$  is transdiagonal if and only if there exists  $1 \leq m \leq n$  such that

$$(2.6) \quad m < \nu_1 + \dots + \nu_m.$$

Recall that the classical lexicographic order, on monomials of same degree, is

$$(2.7) \quad X^\nu \geq_{\text{lex}} X^\mu \quad \text{iff} \quad \nu \geq_{\text{lex}} \mu,$$

where we say that  $\nu$  is lexicographically larger than  $\mu$ ,  $\nu >_{\text{lex}} \mu$ , if the first non-zero part of the vector  $\nu - \mu$  is positive. For example

$$x_1^3 >_{\text{lex}} x_1^2 x_2 >_{\text{lex}} x_1 x_2^2 >_{\text{lex}} x_2^3 \quad \text{since} \quad (3, 0) >_{\text{lex}} (2, 1) >_{\text{lex}} (1, 2) >_{\text{lex}} (0, 3).$$

3. THE  $\mathcal{G}$  BASIS

Following [2], we exploit relations (2.4) to construct a family

$$\mathcal{G} = \{G_\varepsilon\} \subset \mathcal{J}_n$$

indexed by vectors that are transdiagonal. For  $\alpha$  any composition of  $k \leq n$ , the polynomial  $G_\varepsilon$ , with  $\varepsilon := \alpha 0^*$ , is defined to be

$$(3.1) \quad G_\varepsilon := F_\alpha.$$

When  $\alpha \neq 0$ , the vector  $\varepsilon = \alpha 0^*$  is clearly transdiagonal. For a general vector  $\varepsilon$  (not of the form  $\alpha 0^*$ ), the polynomial  $G_\varepsilon$  is defined recursively in the following way. Let  $\varepsilon = w 0 a \beta 0^*$  be the unique factorization of  $\varepsilon$  such that  $w$  is a word of  $k - 1$  non-negative integers,  $a > 0$  is a positive integer, and  $\beta$  is a composition (parts  $> 0$ ). Then we set

$$(3.2) \quad G_\varepsilon = G_{w a \beta 0^*} - x_k G_{w(a-1)\beta 0^*}.$$

By induction on the length of the indexing vectors, both terms on the right of (3.2) are well defined, and we have

- $\ell(w a \beta 0^*) = \ell(w(a-1)\beta 0^*) = \ell(\varepsilon) - 1$ ;
- $w a \beta 0^*$  and  $w(a-1)\beta 0^*$  are transdiagonal as soon as  $\varepsilon$  is transdiagonal.

In fact, let  $m$  be the first ordinate where  $\pi(\varepsilon)$  crosses the diagonal, this is to say the smallest integer such that  $m < \varepsilon_1 + \dots + \varepsilon_m$ . Then the second assertion follows from

$$\varphi_1 + \dots + \varphi_m > \psi_1 + \dots + \psi_m = \varepsilon_1 + \dots + \varepsilon_m - 1 > m - 1,$$

where  $\varphi = w a \beta 0^*$  and  $\psi = w(a-1)\beta 0^*$ .

For example,

$$\begin{aligned} G_{1020} &= G_{1200} - x_2 G_{1100} \\ &= F_{12}(x_1, x_2, x_3, x_4) - x_2 F_{11}(x_1, x_2, x_3, x_4) \\ &= x_1 x_2^2 + x_1 x_3^2 + x_1 x_4^2 + x_2 x_3^2 + x_2 x_4^2 + x_3 x_4^2 + x_1 x_2 x_3 + x_1 x_2 x_4 \\ &\quad + x_1 x_3 x_4 + x_2 x_3 x_4 - x_2(x_1 x_2 + x_1 x_3 + x_1 x_4 + x_2 x_3 + x_2 x_4 + x_3 x_4) \\ &= x_1 x_3^2 + x_1 x_3 x_4 + x_1 x_4^2 - x_2^2 x_3 - x_2^2 x_4 + x_2 x_3^2 + x_2 x_4^2 + x_3 x_4^2. \end{aligned}$$

We observe on this example that the leading monomial (in lex order) of  $G_{1020}$  is  $X^{1020} = x_1^1 x_2^0 x_3^2 x_4^0$ . This holds in general for the  $\mathcal{G}$  family as stated in the following proposition, for which all technical details can be found in [2].

**Proposition 3.1** ([2], Corollary 3.4). *The leading monomial  $LM(G_\varepsilon)$  of  $G_\varepsilon$  is  $X^\varepsilon$ .*

## 4. PROOF OF THE MAIN THEOREM

We now give an explicit basis for the space  $\mathbf{R}_n$  naturally indexed by Dyck paths. This proves the first part of Theorem 1.1.

**Theorem 4.1.** *The set of monomials*

$$(4.1) \quad \mathcal{B}_n = \{X^\eta \mid \pi(\eta) \text{ is a Dyck path}\}$$

*is a basis of the space  $\mathbf{R}_n$ .*

The proof is achieved in a few steps. We start with the following lemma.

**Lemma 4.2.** *Any  $\mathcal{P}(X) \in \mathbb{Q}[X]$  is in the linear span of  $\mathcal{B}_n$  modulo  $\mathcal{J}_n$ . That is*

$$(4.2) \quad P(X) \equiv \sum_{X^\eta \in \mathcal{B}_n} c_\eta X^\eta \pmod{\mathcal{J}_n}.$$

*Proof.* It clearly suffices to show that (4.2) holds for any monomial  $X^\nu$ , with  $\nu$  transdiagonal. We assume that there exists  $X^\nu$  not reducible of the form (4.2) and we choose  $X^\varepsilon$  to be the smallest amongst them with respect to the lexicographic order. Let us write

$$\begin{aligned} X^\varepsilon &= LM(G_\varepsilon) \\ &= (X^\varepsilon - G_\varepsilon) + G_\varepsilon \\ &\equiv X^\varepsilon - G_\varepsilon \pmod{\mathcal{J}_n}. \end{aligned}$$

All monomials in  $(X^\varepsilon - G_\varepsilon)$  are lexicographically smaller than  $X^\varepsilon$ , thus they are reducible. This contradicts our assumption on  $X^\varepsilon$  and completes our proof.  $\square$

Thus  $\mathcal{B}_n$  spans the space  $\mathbf{R}_n$ . We now prove its linear independence. This is equivalent to showing that the set  $\mathcal{G}$  is a Gröbner basis of the ideal  $\mathcal{J}_n$ . A crucial lemma is the following one, which is the quasi-symmetric analogue of a classical result is the case of symmetric polynomials ([11], Theorem II.2.2).

**Lemma 4.3.** *If we denote by  $\mathcal{L}[S]$  the linear span of a set  $S$ , then*

$$(4.3) \quad \mathbb{Q}[X] = \mathcal{L}[X^\eta F_\alpha \mid X^\eta \in \mathcal{B}_n, \alpha \models r \geq 0].$$

*Proof.* We have already obtained the following reduction for any monomial  $X^\varepsilon$  in  $\mathbb{Q}[X]$ .

$$X^\varepsilon \equiv \sum_{X^\eta \in \mathcal{B}_n} c_\eta X^\eta \pmod{\mathcal{J}_n},$$

which is equivalent to

$$(4.4) \quad X^\varepsilon = \sum_{X^\eta \in \mathcal{B}_n} c_\eta X^\eta + \sum_{\alpha \models r \geq 1} Q_\alpha F_\alpha.$$

We then apply the reduction (4.4) to each monomial of the  $Q_\alpha$ 's and use Proposition 2.2 to reduce products of fundamental quasi-symmetric functions. We obtain (4.3) in a finite number of operations since degrees strictly decrease at each operation, because  $\alpha \models r \geq 1$  implies  $\deg Q_\alpha < |\varepsilon|$ .  $\square$

The next lemma is the final step in our proof of the Theorem 4.1.

**Lemma 4.4.** *The set  $\mathcal{G}$  is a linear basis of the ideal  $\mathcal{J}_n$ , i.e.*

$$(4.5) \quad \mathcal{J}_n = \mathcal{L}[G_\varepsilon \mid \varepsilon \text{ transdiagonal}].$$

*Proof.* Let us denote by  $\mathcal{A}_n$  the set

$$(4.6) \quad \mathcal{A}_n = \{X^\xi \mid x_1^{\xi_n} x_2^{\xi_{n-1}} \cdots x_n^{\xi_1} \in \mathcal{B}_n\}.$$

Now the algebra endomorphism of  $\mathbb{Q}[X]$  that reverses the variables,

$$x_i \mapsto x_{n-i+1},$$

clearly fixes the subalgebra  $Qsym$ . In fact it maps  $F_\alpha$  to  $F_{\alpha'}$ , where  $\alpha'$  is the reverse composition.

It follows from Lemma 4.3 and the endomorphism above that:

$$(4.7) \quad \mathbb{Q}[X] = \mathcal{L}[X^\xi F_\alpha \mid X^\xi \in \mathcal{A}_n, \alpha \models r \geq 0].$$

Now to prove Lemma 4.4, we reduce the problem as follows. We first use (4.7) and Proposition 2.2 to write

$$\begin{aligned} \mathcal{J}_n &= \langle F_\alpha, \alpha \models s \geq 0 \rangle_{\mathbb{Q}[X]} = \mathcal{L}[X^\xi F_\alpha F_\beta \mid X^\xi \in \mathcal{A}_n, \alpha \models s \geq 0, \beta \models t \geq 1] \\ &= \mathcal{L}[X^\xi F_\gamma \mid X^\xi \in \mathcal{A}_n, \gamma \models r \geq 1]. \end{aligned}$$

It is now sufficient to prove that for all  $X^\xi \in \mathcal{A}_n$  and all  $\gamma \models r \geq 1$

$$(4.8) \quad X^\xi F_\alpha \in \mathcal{L}[G_\varepsilon \mid \varepsilon \text{ transdiagonal}].$$

But Lemma 4.2 implies that any monomial of degree greater than  $n$  is in  $\mathcal{J}_n$ . Hence to prove (4.8), we need only show it for  $\xi$  and  $\gamma$  such that  $|\xi| + |\gamma| \leq n$ . To do that, we reduce the product

$$(4.9) \quad x_n^{\xi_n} (x_{n-1}^{\xi_{n-1}} (\dots (x_2^{\xi_2} (x_1^{\xi_1} F_\alpha))))$$

recursively, using

$$(4.10) \quad x_k G_{wb\beta 0^*} = G_{w(b+1)\beta 0^*} - G_{w0(b+1)\beta 0^*}$$

or

$$(4.11) \quad x_k G_{w0^*00^*} = G_{w0^*10^*} - G_{w0^*010^*}.$$

Relations (4.10) and (4.11) are immediate consequences of the definition of the  $\mathcal{G}$  basis (Relation (3.2)).

We have to show that the vectors  $\varepsilon$  generated in this process are all transdiagonal and that the length  $\ell(\varepsilon)$  always remains at most equal to  $n$ . Let us first check that the transdiagonal part. This is obvious in the case of relation (4.11). In the other case (relation (4.10)), for  $\varphi = wb\beta 0^*$ , it is sufficient to observe that if  $m$  is such that

$$\varphi_1 + \dots + \varphi_m > m$$

with  $m > \ell(w)$  (if not, it is evident), then

$$\varphi'_1 + \dots + \varphi'_m > m + 1 > m \quad \text{and} \quad \varphi''_1 + \dots + \varphi''_{m+1} > m + 1$$

where  $\varphi' = w(b+1)\beta 0^*$ , and  $\varphi'' = w0(b+1)\beta 0^*$ . We shall now prove that the length of the  $\varepsilon$ 's always remains at most equal to  $n$ . For this we need to keep track of the term  $\varepsilon_{\ell(\varepsilon)}$ . Two cases have to be considered.

- First case:  $\varepsilon_{\ell(\varepsilon)}$  comes from  $\alpha_{\ell(\alpha)}$  that has shifted to the right by Relation (4.10). It could move at most  $|\xi|$  steps to the right, whence

$$\ell(\varepsilon) \leq \ell(\alpha) + |\xi| \leq |\alpha| + |\xi| \leq n.$$



- Second case:  $\varepsilon_{\ell(\varepsilon)}$  is a “1” generated by relation (4.11) that has shifted to the right. If it is generated by a multiplication by  $x_k$ , then we consider the vector

$$\eta = \xi_n \xi_{n-1} \cdots \xi_k 0^*.$$

Since  $X^\xi \in \mathcal{A}_n$  implies  $\pi(\eta)$  is a Dyck path, we have

$$|\eta| < \ell(\eta) = n - k + 1$$

hence the generated “1” can shift at most to position

$$k + |\eta| \leq k + n - k = n.$$

□

The recursive process used to reduce a product of form (4.9) is illustrated in the following example, where  $n = 5$ .

$$\begin{aligned} x_1 x_3 F_{21} &= x_3(x_1 F_{21}) \\ &= x_3(G_{31000} - G_{03100}) \\ &= x_3 G_{31000} - x_3 G_{03100} \\ &= G_{31100} - G_{31010} - G_{03200} + G_{03020}. \end{aligned}$$

*End of proof of Theorem 4.1.* By Lemma 4.2, the set  $\mathcal{B}_n$  spans the quotient  $\mathbf{R}_n$ . Assume we have a linear dependence relation modulo  $\mathcal{J}_n$ , i.e. there exists  $P$

$$P = \sum_{X^\xi \in \mathcal{B}_n} a_\xi X^\xi \in \mathcal{I}_n.$$

By Lemma 4.4,  $\mathcal{J}_n$  is linearly spanned by the  $G_\varepsilon$ 's, thus

$$P = \sum_{\varepsilon \text{ transdiagonal}} b_\varepsilon G_\varepsilon.$$

This implies  $LM(P) = X^\varepsilon$ , with  $\varepsilon$  transdiagonal, which is absurd. □

A consequence of Lemma 4.4 and Theorem 4.1 is that the set  $\mathcal{G}$  is a Gröbner basis of  $\mathcal{J}_n$  with respect to the lex order. From this we see below that a minimal Gröbner basis of  $\mathcal{J}_n$  is obtained from  $\mathcal{G}$  if we select the  $G_\varepsilon \in \mathcal{G}$  such that  $\pi(\varepsilon)$  has exactly one step under the line  $y = x$  and no other horizontal steps after that.

**Corollary 4.5.** *A minimal Gröbner basis for  $\mathcal{J}_n$  is given by*

$$(4.12) \quad \{G_\varepsilon \in \mathcal{G} \mid \varepsilon = w0^*, \ell(w) = |w| + 1, w_1 + \cdots + w_s \leq s, \text{ for } s < \ell(w)\}.$$


*Proof.* Theorem 4.1 implies that the monomial ideal  $LT(\mathcal{J}_n)$  of leading terms of  $\mathcal{J}_n$  is generated by all monomials  $X^\eta$  where  $\pi(\eta)$  is transdiagonal. For any such  $\eta$  let  $m$  be the smallest integer such that  $m < \eta_1 + \cdots + \eta_m$  and let  $\varepsilon = \eta_1 \cdots \eta_{m-1} a 0^*$  where  $a = m - 1 - \eta_1 - \cdots - \eta_{m-1}$ . The monomial  $X^\varepsilon$  divides  $X^\eta$  which shows that  $LT(\mathcal{J}_n)$  is generated by the leading monomials of the  $G_\varepsilon$  in (4.12). This gives that (4.12) is a Gröbner basis. To show minimality, consider  $X^\xi$  a monomial that strictly divides the leading monomial of a  $G_\varepsilon$  in (4.12). Since  $\pi(\varepsilon)$  has exactly one step under the line  $y = x$ , we have that  $\pi(\xi)$  is not transdiagonal and  $X^\xi \notin LT(\mathcal{J}_n)$ . Hence the leading monomials of the  $G_\varepsilon$  in (4.12) is a minimal set of generators for  $LT(\mathcal{J}_n)$ . □

## 5. HILBERT SERIES

Since Theorem 4.1 gives us an explicit basis for the quotient  $\mathbf{R}_n$ , which is isomorphic to  $\mathbf{SH}_n$  as a graded vector space, we are able to refine relation (1.4) by giving the Hilbert series of the space of super-covariant polynomials. For  $k \in \mathbb{N}$ , let  $\mathbf{SH}_n^{(k)}$  and  $\mathbf{R}_n^{(k)}$  denote the projections

$$(5.1) \quad \mathbf{SH}_n^{(k)} = \mathbf{SH}_n \cap \mathbb{Q}^{(k)}[X] \simeq \mathbf{R}_n \cap \mathbb{Q}^{(k)}[X] = \mathbf{R}_n^{(k)}$$

where  $\mathbb{Q}^{(k)}[X]$  is the vector space of homogeneous polynomials of degree  $k$  together with zero. Here, we represent Dyck paths horizontally, with  $n$  rising steps  $(1, 1)$  and  $n$  falling steps  $(1, -1)$ . Let us denote by  $D_n^{(k)}$  the number of Dyck paths of length  $2n$  ending by exactly  $k$  falling steps and by  $C_n^{(k)}$  the number of Dyck paths of length  $2n$  which have exactly  $k$  factors, *i.e.*  $k + 1$  points on the axis. The next figure gives an example of a Dyck path of length 28, ending with 4 falling steps and made of 3 factors.



It is well known that

$$(5.2) \quad D_n^{(k)} = C_n^{(k)} = \frac{k(2n-k-1)!}{n!(n-k)!},$$

where the first equality is classical (*cf.* [23] for example for a bijective proof), and the second corresponds to [17], formula (7).

Let us denote by  $F_n(t)$  the Hilbert series of  $\mathbf{SH}_n$ , *i.e.*

$$(5.3) \quad F_n(t) = \sum_{k \geq 0} \dim \mathbf{SH}_n^{(k)} t^k.$$

**Theorem 5.1.** *For  $0 \leq k \leq n - 1$ , the dimension of  $\mathbf{SH}_n^{(k)}$  is given by*

$$(5.4) \quad \dim \mathbf{SH}_n^{(k)} = \dim \mathbf{R}_n^{(k)} = D_n^{(n-k)} = C_n^{(n-k)} = \frac{n-k}{n+k} \binom{n+k}{k}.$$

*For  $k \geq n$  the dimension of  $\mathbf{SH}_n^{(k)}$  is 0.*

*Proof.* By Theorem 4.1, we know that the set

$$\mathcal{B}_n = \{X^\eta \mid \pi(\eta) \text{ is a Dyck path}\}$$

is a basis for  $\mathbf{R}_n$ . It is then sufficient to observe that the path  $\pi(\eta)$  associated to  $\eta$  ends by exactly  $n - |\eta|$  falling steps.  $\square$

For example, we have:

$n$	$F_n(t)$
1	1
2	$1 + t$
3	$1 + 2t + 2t^2$
4	$1 + 3t + 5t^2 + 5t^3$
5	$1 + 4t + 9t^2 + 14t^3 + 14t^4$
6	$1 + 5t + 14t^2 + 28t^3 + 42t^4 + 42t^5$
7	$1 + 6t + 20t^2 + 48t^3 + 90t^4 + 132t^5 + 132t^6$

This gives

$$(5.5) \quad F_n(t) = \sum_{k=0}^{n-1} \frac{n-k}{n+k} \binom{n+k}{k} t^k.$$

from which one easily deduces that the generating series for the  $F_n(t)$ 's is

$$(5.6) \quad \sum_n F_n(t) x^n = \frac{1 - \sqrt{1 - 4tx} - 2x}{2(t+x-1)}.$$

**Remark 5.2.** The study of various filtrations of the space  $\mathbb{Q}[X]$ , with respect to family of ideals of quasi-symmetric polynomials, will be the object of a forthcoming paper [3].

#### ACKNOWLEDGMENTS

We thank Adriano Garsia for stimulating discussions and suggestions about this work. We also thank Mike Zabrocki for his careful reading.

#### REFERENCES

- [1] E. ARTIN, *Galois Theory*, Notre Dame Mathematical Lecture **2** (1944), Notre Dame, IN.
- [2] J.-C. AVAL AND N. BERGERON, *Catalan Paths and Quasi-Symmetric Functions*, Proc. Amer. Math. Soc., to appear.
- [3] J.-C. AVAL, F. BERGERON, N. BERGERON AND A. GARSIA, *Ideals of Quasi-symmetric polynomials and related varieties*, in preparation.
- [4] F. BERGERON, N. BERGERON, A. GARSIA, M. HAIMAN AND G. TESLER, *Lattice Diagram Polynomials and Extended Pieri Rules*, Adv. Math., **142** (1999), 244-334.
- [5] F. BERGERON, A. GARSIA AND G. TESLER, *Multiple Left Regular Representations Generated by Alternants*, J. of Comb. Th., Series A, **91**, 1-2 (2000), 49-83.
- [6] N. BERGERON, S. MYKYTIUK, F. SOTTILE, AND S. VAN WILLIGENBURG, *Pieri Operations on Posets*, J. of Comb. Theory, Series A, **91** (2000), 84-110.
- [7] A. BOREL, *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes des groupes de Lie compacts*, Ann. of Math., **57** (1953), 115-207.
- [8] C. DE CONCINI AND C. PROCESI, *Symmetric functions, conjugacy classes and the flag variety*, Invent. Math. **64** (1981), 203-230.
- [9] D. COX, J. LITTLE AND D. O'SHEA, *Ideals, Varieties, and Algorithms*, Springer-Verlag, New-York, 1992.
- [10] A. M. GARSIA AND M. HAIMAN, *A graded representation model for Macdonald's polynomials*, Proc. Nat. Acad. Sci. U.S.A. **90** (1993), no. 8, 3607-3610.
- [11] A. Garsia, M. Haiman, *Orbit Harmonics and Graded Representations*, Éditions du Lacim, to appear.

- [12] I. GELFAND, D. KROB, A. LASCoux, B. LECLERC, V. RETAKH, AND J.-Y. THIBON, *Non-commutative symmetric functions*, Adv. in Math., **112** (1995), 218–348.
- [13] I. GESSEL, *Multipartite  $P$ -partitions and products of skew Schur functions*, in Combinatorics and Algebra (Boulder, Colo., 1983), C. Greene, ed., vol. 34 of Contemp. Math., AMS, 1984, pp. 289–317.
- [14] M. HAIMAN, *Hilbert schemes, polygraphs, and the Macdonald positivity conjecture*, J. Amer. Math. Soc., **14** (2001), 941–1006.
- [15] F. HIVERT, *Hecke algebras, difference operators, and quasi-symmetric functions*, Adv. in Math., **155** (2000), 181–238.
- [16] V. JONES, *Index for subfactors*, Inv. Math. **72** (1983) 1–25.
- [17] G. KREWERAS, *Sur les éventails de segments*, Cahiers du BURO, **15** (1970), 3–41.
- [18] I. MACDONALD, *Symmetric Functions and Hall Polynomials*, Oxford Univ. Press, 1995, second edition.
- [19] C. MALVENUTO AND C. REUTENAUER, *Duality between quasi-symmetric functions and the Solomon descent algebra*, Journal of Algebra, **177** (1995), 967–982.
- [20] R. STANLEY, *Enumerative Combinatorics, Vol. 1*, Wadsworth and Brooks/Cole, 1986.
- [21] R. STANLEY, *Enumerative Combinatorics Vol. 2*, no. 62 in Cambridge Studies in Advanced Mathematics, Cambridge University Press, 1999. Appendix 1 by Sergey Fomin.
- [22] R. STEINBERG, *Differential equations invariant under finite reflection groups*, Trans. Amer. Math. Soc., **112** (1964), 392–400.
- [23] J. VALLÉ, *Une bijection explicative de plusieurs propriétés remarquables des ponts*, Europ. J. Combinatorics, **18** (1997), 117–124.

(J.-C. Aval) LABORATOIRE A2X, UNIVERSITÉ BORDEAUX 1, 351 COURS DE LA LIBÉRATION,  
33405 TALENCE CEDEX, FRANCE

*E-mail address:* aval@math.u-bordeaux.fr

(F. Bergeron) DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DU QUÉBEC À MONTRÉAL,  
MONTRÉAL, QUÉBEC, H3C 3P8, CANADA

*E-mail address:* bergeron.francois@uqam.ca

(N. Bergeron) DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO,  
ONTARIO M3J 1P3, CANADA

*E-mail address:* bergeron@mathstat.yorku.ca

*URL:* <http://www.math.yorku.ca/bergeron>