

Uniform bounds for the exponential moments of the maximum of a Dyck path

Oleksiy Khorunzhiy

Université de Versailles

45 avenue des Etats-Unis

78035 Versailles cedex

khorunzhiy@math.uvsq.fr

Jean-François Marckert

CNRS, LaBRI

Université Bordeaux

351 cours de la Libération

33405 Talence cedex

marckert@labri.fr

Abstract

Let D_{2n} be a Dyck path chosen uniformly in the set of Dyck paths with $2n$ steps. The aim of this note is to show that for any $\lambda > 0$ the sequence $\mathbb{E}(\exp(\lambda(2n)^{-1/2} \max D_{2n}))$ converges, and therefore is bounded uniformly in n . The uniform bound justifies an assumption used in literature to prove certain estimates of high moments of large random matrices.

2000 Mathematics Subject Classification: 60C05; 60G70; 60F99

Key Words: Dyck paths, Bernoulli bridge, simple random walks, random matrices

1 Introduction

Let $\mathbb{N} = \{0, 1, 2, 3, \dots\}$ be the set of non-negative integers. For any $n \in \mathbb{N}$, we denote by \mathcal{W}_n the set of Bernoulli chains with n steps :

$$\mathcal{W}_n = \{\mathbf{S} = (S_i)_{0 \leq i \leq n} : S_0 = 0, S_{i+1} = S_i \pm 1 \text{ for any } i \in \llbracket 0, n-1 \rrbracket\},$$

where $\llbracket a, b \rrbracket = [a, b] \cap \mathbb{N}$. The set of Dyck paths \mathcal{D}_n (sometimes called simple or Bernoulli excursions) is defined by

$$\mathcal{D}_n = \{\mathbf{S} : \mathbf{S} \in \mathcal{W}_n, S_n = 0, S_i \geq 0 \text{ for any } i \in \llbracket 0, n \rrbracket\}.$$

Of course \mathcal{D}_n is empty for odd n , and one has

$$\#\mathcal{D}_{2n} = \frac{1}{n+1} \binom{2n}{n} \tag{1}$$

the n th Catalan number. Let $\mathbb{P}_n^{(w)}$ and $\mathbb{P}_{2n}^{(d)}$ be the uniform distributions on \mathcal{W}_n and \mathcal{D}_{2n} . The expectations with respect to these measures will be denoted by $\mathbb{E}_n^{(w)}$ and $\mathbb{E}_{2n}^{(d)}$. The aim of this note is to prove the following statement.

Theorem 1 For any $\lambda > 0$, we have

$$\mathbb{E}_{2n}^{(d)} \left(\exp \left(\lambda \frac{\max \mathbf{S}}{\sqrt{2n}} \right) \right) \rightarrow_n \mathbb{E} \left(\exp \left(\lambda \max_{t \in [0,1]} \mathbf{e}(t) \right) \right) \quad (2)$$

where $(\mathbf{e}(t), t \in [0, 1])$ is the normalized Brownian excursion. In particular, for any $\lambda > 0$

$$\sup_n \mathbb{E}_{2n}^{(d)} (\exp(\lambda n^{-1/2} \max \mathbf{S})) < +\infty. \quad (3)$$

We may notice that the right hand side is finite for every λ using the computation of Chung [2]:

$$\mathbb{P} \left(\max_{t \in [0,1]} \mathbf{e}(t) \leq x \right) = 1 + 2 \sum_{j \geq 1} (1 - 4j^2 x^2) e^{-2j^2 x^2}, \quad \text{for } x > 0,$$

and then $\max_{t \in [0,1]} \mathbf{e}(t)$ possesses all exponential moments.

1.1 Relations with the spectral theory of large random matrices

Dyck paths play a central role in combinatorics and arise in many situations (see for instance 66 examples of the appearance of the Catalan numbers in combinatorics in Stanley [10, ex. 6.19 p.219]). In the present case, the motivation for establishing Theorem 1 comes from the study of the moments of random real symmetric (or hermitian) matrices known as the Wigner ensemble [13, 16]. In these settings, the Catalan numbers (1) represent the moments of the eigenvalue distribution of random matrices of Wigner ensemble A_N in the limit of their infinite dimension $N \rightarrow \infty$ [16]. Recent studies [13, 15] of high moments of large Wigner random matrices have used the exponential moments of the maxima of the Dyck paths. More precisely, it was shown that [7, 15]

$$\mathbb{E} (\text{Tr } A_N^{2n}) \leq C_1 \frac{e^{C_2 t^{1/3}}}{t^{1/6}} Q_n(C_3 t^{1/6}) (1 + o(1)), \quad n = \lfloor tN^{2/3} \rfloor, \quad N \rightarrow \infty, \quad (4)$$

where Tr denotes the trace of the square matrix, C_1 , C_2 and C_3 are certain constants, and

$$Q_n(\lambda) = \mathbb{E}_{2n}^{(d)} \left(\exp \left(\lambda \frac{\max \mathbf{S}}{\sqrt{2n}} \right) \right).$$

It was assumed in [7, 13, 15] that $\limsup_n Q_n(\lambda)$ is bounded. Theorem 1 above shows that this assertion is true.

2 Proof of the Theorem

Before proving the Theorem, we first discuss the appearance of $\max_{t \in [0,1]} \mathbf{e}(t)$ and the non-triviality of the result. Let $C[0,1]$ be the set of continuous functions defined on $[0,1]$ with real values. For any $\mathbf{S} \in \mathcal{W}_n$, denote by $u_n = u_n^{\mathbf{S}}$ the function in $C[0,1]$ obtained from \mathbf{S} by interpolation and rescaling:

$$u_n(t) = \frac{1}{\sqrt{n}}(S(\lfloor nt \rfloor) + \{nt\}(S(\lceil nt \rceil) - S(\lfloor nt \rfloor))) \text{ for any } t \in [0,1]. \quad (5)$$

It is known that under $\mathbb{P}_{2n}^{(d)}$, $u_{2n} \xrightarrow[n]{(d)} \mathbf{e}$ in $C[0,1]$ endowed with the topology of uniform convergence where $\xrightarrow[n]{(d)}$ means the convergence in distribution (see e.g. Kaigh [6] where this result is shown for general increment distributions). By continuity of the map $f \mapsto e^{\lambda \max f}$ from $C[0,1]$ into \mathbb{R} , under $\mathbb{P}_{2n}^{(d)}$,

$$\exp\left(\frac{\lambda \max \mathbf{S}}{\sqrt{2n}}\right) = \exp(\lambda \max u_{2n}) \xrightarrow[n]{(d)} \exp\left(\lambda \max_{t \in [0,1]} \mathbf{e}(t)\right). \quad (6)$$

Then the uniform integrability argument is sufficient: given $\lambda > 0$, in order to prove that (6) implies (2), it suffices to show that

$$\sup_n \mathbb{E}_{2n}^{(d)} \left(\exp\left(\left(\lambda + \varepsilon\right) \frac{\max \mathbf{S}}{\sqrt{2n}}\right) \right) < +\infty, \quad (7)$$

for some $\varepsilon > 0$ (see Billingsley [1, Section 16]). Hence to prove the Theorem, using (6), only the second assertion (3) (weaker in appearance) needs to be proved. This is what we will do.

Remark 1 *Smith & Diaconis [14] proved that $\mathbb{P}_{2n}^{(d)}\left(\frac{\max S}{\sqrt{2n}} \leq y\right) = \mathbb{P}_{2n}^{(d)}(\max \mathbf{e} \leq y) + O(n^{1/2})$, and the convergence of moments of $\frac{\max S}{\sqrt{2n}}$ (under $\mathbb{P}_{2n}^{(d)}$) to those of $\max \mathbf{e}$ is also known (see Flajolet & Odlyzko [5] and references therein, where this is stated in link with the convergence of the height of random trees). These convergence results are not strong enough to obtain Theorem 1.*

The strategy will be at first to transform the question in terms of Bernoulli bridges, and then to transform the question in terms of simple random walks where the answer is easy. The steps follow some ideas developed in Janson & Marckert [8] in their proof of their Lemma 1.

2.1 From Dyck paths to Bernoulli bridges

Let us introduce the set \mathcal{B}_n of “Bernoulli bridges” with n steps

$$\mathcal{B}_n = \{\mathbf{S} : \mathbf{S} \in \mathcal{W}_n, S_n = -1\}.$$

The quotes around “Bernoulli bridges” are there to signal that often the terms “Bernoulli bridges” concerns walks ending at 0 instead at -1 . Clearly, \mathcal{B}_n is empty for even n and it is easy to see that $\#\mathcal{B}_{2n+1} = \binom{2n+1}{n}$; we denote by $\mathbb{P}_{2n+1}^{(b)}$ the uniform distribution on \mathcal{B}_{2n+1} , and by $\mathbb{E}_{2n+1}^{(b)}$ the expectation with respect to $\mathbb{P}_{2n+1}^{(b)}$.

The cycle Lemma (introduced by Dvoretzky & Motzkin [3], see also Raney [9] and also Pitman [12], Section 6.1) allows to relate quantities on Dyck paths and on Bernoulli bridges, and among other explains why

$$(2n + 1)\#\mathcal{D}_{2n} = \#\mathcal{B}_{2n+1}. \quad (8)$$

Consider the set of Dyck paths with size $2n$ with an additional last step -1 :

$$\mathcal{D}_{2n+1}^* := \{\mathbf{S} : \mathbf{S} \in \mathcal{W}_{2n+1}, S_i \geq 0 \text{ for any } i \in \llbracket 0, 2n \rrbracket, S_{2n} = 0, S_{2n+1} = -1\}.$$

Obviously there is a canonical correspondence between \mathcal{D}_{2n+1}^* and \mathcal{D}_{2n} , and this correspondence conserves the value of the maximum of the paths. Now, the left hand side of (8) is viewed to be the cardinality of $\mathcal{D}_{2n+1}^* \times \llbracket 1, 2n+1 \rrbracket$.

We state the cycle Lemma as follows:

Lemma 2 (*Cycle Lemma*) *There exists a one-to-one correspondence Ψ_{2n+1} between $\mathcal{D}_{2n+1}^* \times \llbracket 1, 2n+1 \rrbracket$ and \mathcal{B}_{2n+1} and such that if $\mathbf{S}' = \Psi_{2n+1}(\mathbf{S}, k)$ for some $k \in \llbracket 1, 2n+1 \rrbracket$ then*

$$|(\max \mathbf{S}) - (\max \mathbf{S}' - \min \mathbf{S}')| \leq 1. \quad (9)$$

We provide a proof of this classical result for reader’s convenience.

Proof. For any walk \mathbf{S} in \mathcal{W}_{n+1} , let

$$\Delta_j(\mathbf{S}) = \mathbf{S}_{j+1} - \mathbf{S}_j, \quad j \in \llbracket 0, n \rrbracket$$

denote the list of increments of \mathbf{S} . For a fixed (\mathbf{S}, k) element in $\mathcal{D}_{2n+1}^* \times \llbracket 1, 2n+1 \rrbracket$, we let $\Psi_{2n+1}(\mathbf{S}, k)$ be the walk which list of increments is $(\Delta_{(i+k) \bmod (2n+1)}(\mathbf{S}), i = 0 \dots, 2n)$.

For any k in $\llbracket 1, 2n+1 \rrbracket$, $\mathbf{S}' := \Psi_{2n+1}(\mathbf{S}, k)$ is indeed a bridge since the sum of the increments is -1 . We now explain why Ψ_{2n+1} is a bijection from $\mathcal{D}_{2n+1}^* \times \llbracket 1, 2n+1 \rrbracket$ onto \mathcal{B}_{2n+1} . For a fixed element $\mathbf{S} \in \mathcal{D}_{2n+1}^*$, let

$$\Psi_{2n+1}(\mathbf{S}) := \left\{ \Psi_{2n+1}^{(k)}(\mathbf{S}), k \in \llbracket 0, 2n \rrbracket \right\},$$

be “a rotation class”. It is easy to see that $\Psi_{2n+1}(\mathbf{S}, k)$ reaches its minimum for the first time at time $2n+1-k$. Hence, $\Psi_{2n+1}(\mathbf{S}, k) \neq \Psi_{2n+1}(\mathbf{S}, k')$ if $k \neq k'$, and each rotation class $\Psi_{2n+1}(\mathbf{S})$ contains a unique Dyck path. It remains to explain why each bridge belongs to a unique rotation class: take a bridge \mathbf{S} that reaches its minimum for the first time at time k . The walk \mathbf{S}' which list of increments is $(\Delta_{(i+2n+1-k) \bmod (2n+1)}(\mathbf{S}), i = 0, \dots, 2n)$ is a Dyck path. Thus, $\Psi_{2n+1}(\mathbf{S}', k) = \mathbf{S}$, and then \mathbf{S} belongs to the rotation class of \mathbf{S}' (and only to this one).

As a conclusion, each rotation class contains a unique element of \mathcal{D}_{2n+1}^* , has cardinality $2n+1$, and of course each element of \mathcal{D}_{2n+1}^* belongs to a rotation class. Now, it is easy to see that for \mathbf{S} in \mathcal{D}_{2n+1}^* , for any $k \in \llbracket 0, 2n \rrbracket$, the bridges $\mathbf{S}' = \Psi_{2n+1}(\mathbf{S}, k)$ satisfies $|(\max \mathbf{S}) - (\max \mathbf{S}' - \min \mathbf{S}')| \leq 1$. \square

Hence, the uniform distribution on \mathcal{B}_{2n+1} is the push-forward measure of the uniform distribution on $\mathcal{D}_{2n+1}^* \times \llbracket 1, 2n+1 \rrbracket$ by Ψ_{2n} (which amounts to first choosing a Dyck path uniformly, and then a rotation). It follows from all these considerations that

$$\sup_n \mathbb{E}_{2n}^{(d)} \left(e^{\lambda \frac{\max \mathbf{S}}{\sqrt{2n}}} \right) < +\infty \text{ if and only if } \sup_n \mathbb{E}_{2n+1}^{(b)} \left(e^{\lambda \frac{\max \mathbf{S} - \min \mathbf{S}}{\sqrt{2n}}} \right) < +\infty.$$

We now show that this second assertion holds.

2.2 From bridges to simple walks

For any walk \mathbf{S} let

$$Y_{[a,b]}^{\mathbf{S}} := \max_{a \leq k \leq b} S_k - \min_{a \leq k \leq b} S_k.$$

We have, using $Y_{[0,2n+1]}^{\mathbf{S}} \leq Y_{[0,n]}^{\mathbf{S}} + Y_{[n,2n+1]}^{\mathbf{S}}$ and the Cauchy-Schwarz inequality

$$\mathbb{E}_{2n+1}^{(b)} \left(e^{\lambda \frac{\max \mathbf{S} - \min \mathbf{S}}{\sqrt{2n}}} \right) = \mathbb{E}_{2n+1}^{(b)} \left(e^{\lambda \frac{Y_{[0,2n+1]}^{\mathbf{S}}}{\sqrt{2n}}} \right) \quad (10)$$

$$\leq \left[\mathbb{E}_{2n+1}^{(b)} \left(e^{2\lambda \frac{Y_{[0,n]}^{\mathbf{S}}}{\sqrt{2n}}} \right) \mathbb{E}_{2n+1}^{(b)} \left(e^{2\lambda \frac{Y_{[n,2n+1]}^{\mathbf{S}}}{\sqrt{2n}}} \right) \right]^{1/2} \quad (11)$$

The idea here is to work on the half of the trajectory where the conditioning $S_{2n+1} = -1$ will appear to be "not so important". Since a time reversal of Bernoulli bridges with size $2n + 1$, followed by a symmetry with respect to the x -axis send \mathcal{B}_{2n+1} onto \mathcal{B}_{2n+1} and exchange the "two halves" of the trajectory, we just have to prove that for $a = 0$ and $a = 1$

$$\sup_n \mathbb{E}_{2n+1}^{(b)} \left(e^{2\lambda Y_{[0,n+a]}^{\mathbf{S}}} \right) < +\infty. \quad (12)$$

We will treat the case $a = 0$ the other one being very similar.

We equip the space \mathcal{W}_{2n+1} with the filtration $\mathcal{F} := (\mathcal{F}_k)$ where \mathcal{F}_k is generated by the random variables (S_1, \dots, S_k) .

Lemma 3 *Let A_n be an \mathcal{F}_n -measurable event. We have*

$$\mathbb{P}_{2n+1}^{(b)}(A_n) \leq C_0 \mathbb{P}_{2n+1}^{(w)}(A_n) = C_0 \mathbb{P}_n^{(w)}(A_n), \quad (13)$$

for a constant C_0 valid for all n (and all A_n).

Proof . The equality in this formula is clear since under $\mathbb{P}_{2n+1}^{(w)}$, \mathbf{S} is a Markov chain. Only the existence of C_0 is needed to be proved. In the following computations, we will use that since both $\mathbb{P}_{2n+1}^{(w)}$ and $\mathbb{P}_{2n+1}^{(b)}$ are the uniform distributions on their respective set, we have

$$\mathbb{P}_{2n+1}^{(b)} = \mathbb{P}_{2n+1}^{(w)}(\cdot | S_{2n+1} = -1).$$

We will also use that under both $\mathbb{P}_{2n+1}^{(w)}$ and $\mathbb{P}_{2n+1}^{(b)}$, \mathbf{S} is a Markov chain.

Let A be $\mathcal{F}_n^{(S)}$ -measurable, we have

$$\mathbb{P}_{2n+1}^{(w)}(A | S_n = k, S_{2n+1} = -1) = \mathbb{P}_{2n+1}^{(w)}(A | S_n = k) = \mathbb{P}_n^{(w)}(A | S_n = k). \quad (14)$$

We have the following chain of equalities

$$\mathbb{P}_{2n+1}^{(b)}(A) = \mathbb{P}_{2n+1}^{(w)}(A | S_{2n+1} = -1) = \frac{\mathbb{P}_{2n+1}^{(w)}(A, S_{2n+1} = -1)}{\mathbb{P}_{2n+1}^{(w)}(S_{2n+1} = -1)} \quad (15)$$

$$= \sum_k \mathbb{P}_n^{(w)}(A | S_n = k) \mathbb{P}_{2n+1}^{(w)}(S_n = k | S_{2n+1} = -1). \quad (16)$$

(this last equality coming from (14)). Let us denote by $N(m, j)$ the number of trajectories of simple walks of the length m that end at j . Clearly $N(m, j) = \binom{m}{(m+j)/2}$, when m and j have the same parity, zero if not. Also it is easy to see that the number

of trajectories \mathbf{S} in \mathcal{B}_{2n+1} such that $S_n = j$ is $N(n, j)N(n+1, j+1)$. Then one obtains by simple counting arguments that

$$\begin{aligned} \mathbb{P}_{2n+1}^{(w)}(S_n = k | S_{2n+1} = -1) &= \frac{N(n, k)N(n+1, k+1)}{N(2n+1, -1)} \\ &= \frac{N(n, k)}{2^n} \frac{2^n N(n+1, k+1)}{N(2n+1, -1)} \\ &= \mathbb{P}_n^{(w)}(S_n = k) \frac{2^n N(n+1, k+1)}{N(2n+1, -1)} \leq C_0 \mathbb{P}_n^{(w)}(S_n = k), \end{aligned}$$

where

$$C_0 = \sup_{n \geq 1} \sup_k \frac{2^n N(n, k+1)}{N(2n+1, -1)} = \sup_{n \geq 1} 2^n \binom{n}{\lfloor n/2 \rfloor} \binom{2n}{n}^{-1},$$

is indeed finite (as one may check using the Stirling formula). Hence, the right hand side in (16) is bounded by

$$\sum_k \mathbb{P}_n^{(w)}(A | S_n = k) C_0 \mathbb{P}_n^{(w)}(S_n = k) = C_0 \mathbb{P}_n^{(w)}(A).$$

This ends the proof of the Lemma. \square

To conclude the proof of Theorem 1, we explain why (12) holds true. Using Lemma 3, we have

$$\sup_n \mathbb{E}_{2n+1}^{(b)} \left(e^{2\lambda Y_{[0, n]}^{\mathbf{S}}} \right) \leq C_0 \sup_n \mathbb{E}_n^{(w)} \left(e^{2\lambda Y_{[0, n]}^{\mathbf{S}}} \right), \quad (17)$$

since $Y_{[0, n]}^{\mathbf{S}}$ is \mathcal{F}_n -measurable. The right hand side of (17) is much simpler than the left, since it deals with simple random walks under the uniform distribution. Then using again Cauchy-Schwarz, it suffices to show that

$$\sup_n \mathbb{E}_n^{(w)} \left(e^{4\lambda \frac{\max_{0 \leq k \leq n} S_k}{\sqrt{2n}}} \right) < +\infty \quad (18)$$

and the same thing for max replaced by min (which gives the same quantity). Now, by the André reflexion principle (see Feller [4], page 72), we have

$$\mathbb{P}_n^{(w)} \left(\max_{0 \leq k \leq n} S_k \geq x \right) = 2\mathbb{P}_n^{(w)}(S_n > x) + \mathbb{P}_n^{(w)}(S_n = x) \leq 2\mathbb{P}_n^{(w)}(S_n \geq x).$$

Now, the use of Hoeffding's inequality yields directly (18).

References

- [1] P. Billingsley, *Probability and measure*, second Edition, (1986).
- [2] K.L. Chung, *Maxima in Brownian excursions*. Bull. Amer. Math. Soc. 81 (1975), 742–745.
- [3] A. Dvoretzky and Th. Motzkin, *A problem of arrangements*, Duke Math. J. **14**(1947), 305–313.
- [4] W. Feller, *An introduction to probability theory and its applications.*, Vol. I. Third edition John Wiley & Sons, Inc., New York-London-Sydney (1968).
- [5] Ph. Flajolet, and A. Odlyzko, *The average height of binary trees and other simple trees*, J. Comput. System Sci., 25, 2, (1982) 171–213.
- [6] W.D. Kaigh, *An invariance principle for random walk conditioned by a late return to zero*, Ann. Probability 4, No. 1, (1976) 115–121.
- [7] O. Khorunzhiy and V. Vengerovsky, *Ewen walks and estimates of high moments of large Wigner random matrices*, preprint arXiv:0806.0157.
- [8] J.F. Marckert and S. Janson, *Convergence of Discrete Snakes*, Journal of Theoretical Probability **18** No. 3 , (2005) 615– 645.
- [9] G. N. Raney, *Functional composition patterns and power series reversion*, Trans. Amer. Math. Soc. **94** (1960) 441–451.
- [10] R.P. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge University Press. (1999)
- [11] M.L. Mehta, *Random Matrices*, Academic Press, Second edition, (1991).
- [12] J. Pitman, *Combinatorial stochastic processes*, Lecture Notes in Mathematics Vol. 1875, Springer-Verlag, Berlin (2006). Lectures from the 32nd Summer School on Probability Theory held in Saint-Flour, July 7–24, 2002.
- [13] Ya. G. Sinai and A. Soshnikov, *A refinement of Wigner’s semicircle law in a neighborhood of the spectrum edge for random symmetric matrices*, Func. Anal. Appl. **32** (1998) 114–131.
- [14] L. Smith, and P. Diaconis, *Honest Bernoulli excursions*, J. Appl. Probab., **25**, 3, (1988), 464–477.
- [15] A. Soshnikov, *Universality of the edge of the spectrum in Wigner random matrices*, Commun. Math. Phys. **207** (1999) 697–733.
- [16] E. Wigner, *Characteristic vectors of bordered matrices with infinite dimensions*, Ann. Math. **62** (1955) 548–564.