


# Combinatoire des graphes de Strebel

Bertrand Eynard, IPHT CEA Saclay,



Journées Combinatoires de Bordeaux, février 2014

# 1. Introduction

# Quantum gravity

Idea: "2D quantum gravity"  $\Leftrightarrow$  "summing over surfaces"

$$\mathbb{E}(\text{"observable"}) = \int_{\mathcal{M}} d[S] e^{-\text{Energy}(S)} \text{ observable}(S)$$

"Energy"=measure. "observable" can be "Area", "curvature", "geodesic distance between marked points", or anything ...

Claimed by physicists:  $\Leftrightarrow$  Liouville CFT (implies KdV, KPZ, GFF, SLE)

Ensemble:  $\mathcal{M}$ = "set of all Riemann surfaces", possibly of given topology, modulo automorphisms.

Given topology:  $\mathcal{M} = \mathcal{M}_{g,n}$ = moduli space of Riemann surfaces of genus  $g$  with  $n$  marked points.

$\mathcal{M}_{g,n}$  is a complex manifold (orbifold) of dimension

$$\dim_{\mathbb{R}} \mathcal{M}_{g,n} = 2 \times (3g - 3 + n)$$

Un-marking points (forgetful maps):

$$\mathcal{M}_{g,n} \subset \mathcal{M}_{g,n+1} \subset \cdots \subset \mathcal{M}_{g,n+k} \subset \dots$$

# Quantum gravity

Idea: "2D quantum gravity"  $\Leftrightarrow$  "summing over surfaces"

$$\mathbb{E}(\text{"observable"}) = \int_{\mathcal{M}} d[S] e^{-\text{Energy}(S)} \text{ observable}(S)$$

"Energy"=measure. "observable" can be "Area", "curvature", "geodesic distance between marked points", or anything ...

Claimed by physicists:  $\Leftrightarrow$  Liouville CFT (implies KdV, KPZ, GFF, SLE)

Ensemble:  $\mathcal{M}$ = "set of all Riemann surfaces", possibly of given topology, modulo automorphisms.

Given topology:  $\mathcal{M} = \mathcal{M}_{g,n}$ = moduli space of Riemann surfaces of genus  $g$  with  $n$  marked points.

$\mathcal{M}_{g,n}$  is a complex manifold (orbifold) of dimension

$$\dim_{\mathbb{R}\mathbb{C}} \mathcal{M}_{g,n} = 2 \times (3g - 3 + n)$$

Un-marking points (forgetful maps):

$$\mathcal{M}_{g,n} \subset \mathcal{M}_{g,n+1} \subset \cdots \subset \mathcal{M}_{g,n+k} \subset \dots$$

# Quantum gravity

Idea (in the 70's-90's): Discretize.

Family of discrete subsets  $\mathbb{M}_{g,n} \subset \mathcal{M}_{g,n}$ .  $\mathbb{M}_{g,n}$  = set of maps of genus  $g$  with  $n$  marked faces.

Ex:  $\mathbb{M}_{0,1}$  = set of rooted planar maps.

Forgetful maps       $\mathbb{M}_{g,n} \subset \mathbb{M}_{g,n+1} \subset \cdots \subset \mathbb{M}_{g,n+k} \subset \cdots$

$$\mathbb{E}(\text{"observable"}) \stackrel{?}{=} \lim_{\text{continue } k \rightarrow \infty} \sum_{\mathbb{M}_{g,n+k}} e^{-\text{Energy}_k(S)} \text{observable}(S)$$

Remark on observables:

- some observables are "global" (ex: area) easily obtained by a derivative of the energy.
- some observables are "topological" independent of a metric (ex: average curvature, or Chern classes)
- some observables more subtle: geodesic distance between marked points. Need an "embedding" with a metric.

## In the 80's: Witten's KdV conjecture

- maps  $\Leftrightarrow$  *formal* matrix models (BIPZ78)
- matrix models  $\Leftrightarrow$  orthogonal polynomials (Dyson, Gaudin, Mehta,...60's). *True only for not formal.*
- "double scaling limit" of orthogonal polynomials  $\Leftrightarrow$  KdV Tau-function (Kazakov,...80's) + KPZ scaling exponents. *Ansatz not proved before the 2000's.*

→ led Witten's to conjecture:

Continuum limit of partition function of maps = KdV Tau–function

In 1992: M. Kontsevich proved that topological gravity = KdV:

$$\int_{\mathcal{M}_{g,n}} \text{Chern classes} = \mathcal{T}_{\text{KdV}}$$

Kontsevich's proof: *without using discretization !*

## In the 80's: Witten's KdV conjecture

- maps  $\Leftrightarrow$  *formal* matrix models (BIPZ78)
- matrix models  $\Leftrightarrow$  orthogonal polynomials (Dyson, Gaudin, Mehta,...60's). *True only for not formal.*
- "double scaling limit" of orthogonal polynomials  $\Leftrightarrow$  KdV Tau-function (Kazakov,...80's) + KPZ scaling exponents. *Ansatz not proved before the 2000's.*

→ led Witten's to conjecture:

Continuum limit of partition function of maps = KdV Tau–function

In 1992: M. Kontsevich proved that topological gravity = KdV:

$$\int_{\mathcal{M}_{g,n}} \text{Chern classes} = \mathcal{T}_{\text{KdV}}$$

Kontsevich's proof: *without using discretization !*

## In the 80's: Witten's KdV conjecture

- maps  $\Leftrightarrow$  *formal* matrix models (BIPZ78)
- matrix models  $\Leftrightarrow$  orthogonal polynomials (Dyson, Gaudin, Mehta,...60's). *True only for not formal.*
- "double scaling limit" of orthogonal polynomials  $\Leftrightarrow$  KdV Tau-function (Kazakov,...80's) + KPZ scaling exponents. *Ansatz not proved before the 2000's.*

→ led Witten's to conjecture:

Continuum limit of partition function of maps = KdV Tau–function

In 1992: M. Kontsevich proved that topological gravity = KdV:

$$\int_{\mathcal{M}_{g,n}} \text{Chern classes} = \mathcal{T}_{\text{KdV}}$$

Kontsevich's proof: *without using discretization !*

## In the 80's: Witten's KdV conjecture

- maps  $\Leftrightarrow$  *formal* matrix models (BIPZ78)
- matrix models  $\Leftrightarrow$  orthogonal polynomials (Dyson, Gaudin, Mehta,...60's). *True only for not formal.*
- "double scaling limit" of orthogonal polynomials  $\Leftrightarrow$  KdV Tau-function (Kazakov,...80's) + KPZ scaling exponents. *Ansatz not proved before the 2000's.*

→ led Witten's to conjecture:

Continuum limit of partition function of maps = KdV Tau–function


In 1992: M. Kontsevich proved that topological gravity = KdV:

$$\int_{\mathcal{M}_{g,n}} \text{Chern classes} = \mathcal{T}_{\text{KdV}}$$

Kontsevich's proof: *without using discretization !*

# Strebel Graphs

Kontsevich's method: Canonical bijection with **Strebel graphs**



**Bijection :**  $\mathcal{M}_{g,n} \times \mathbb{R}_+^n \longleftrightarrow \bigoplus_{G \in \mathcal{G}_{g,n}} G \times \mathbb{R}_+^{\#\text{edges}}$

$\mathcal{G}_{g,n}$  = set of metric graphs of genus  $g$ , with  $n$  marked faces, and edge lengths  $\ell_e$ .



For each marked face  $L_f$  =perimeter:  $L_f = \sum_{e \rightarrow f} \ell_e$ .

Remark: for 3-valent graphs:  $\#\text{edges} = 6g - 6 + 3n$ .

Higher valent vertices obtained when some  $\ell_e \rightarrow 0$ .

# Strebel Graphs


Metric graph  $\mapsto$  Riemann surface



Ex :  $G \in \mathcal{G}_{1,1}$


$$\text{Perimeter } L = 2l_1 + 2l_2 + 2l_3.$$

For fixed perimeter, an element of  $\mathcal{M}_{1,1}$  is thus parametrized by  $(l_1, l_2) \in \mathbb{R}_+^2$ . Indeed, genus 1 Riemann surfaces are parametrized by a complex modulus  $\tau \in \mathbb{C}$



# Strebel Graphs


Metric graph  $\mapsto$  Riemann surface



Ex :  $G \in \mathcal{G}_{1,1}$


Perimeter  $L = 2l_1 + 2l_2 + 2l_3$ .

For fixed perimeter, an element of  $\mathcal{M}_{1,1}$  is thus parametrized by  $(l_1, l_2) \in \mathbb{R}_+^2$ . Indeed, genus 1 Riemann surfaces are parametrized by a complex modulus  $\tau \in \mathbb{C}$



# Strebel Graphs


Metric graph  $\mapsto$  Riemann surface



Ex :  $G \in \mathcal{G}_{1,1}$

$$\text{Perimeter } L = 2\ell_1 + 2\ell_2 + 2\ell_3.$$

For fixed perimeter, an element of  $\mathcal{M}_{1,1}$  is thus parametrized by  $(\ell_1, \ell_2) \in \mathbb{R}_+^2$ . Indeed, genus 1 Riemann surfaces are parametrized by a complex modulus  $\tau \in \mathbb{C}$



# Strebel's bijection

Riemann surface  $(S, p_1, \dots, p_n) \mapsto$  Metric graph


**Strebel's theorem:** there exists a unique "metric"  $\sqrt{|\Omega|^2}$  on  $S$ , singular (double poles) at the  $p_i$ 's, with residues  $-L_i^2$ :

$$\Omega = \sum_{i=1}^n \frac{-L_i^2}{(x - p_i)^2} dx^2 + \text{at most simple poles}$$


such that all closed geodesics are singular.

The lines  $\text{Im } \Phi(x) = \text{constant}$  are geodesics:

$$\Phi(x) = \int^x \sqrt{\Omega}$$



near a pole  $p_i$  :



, near a zero :

# Strebel's bijection


Example in  $\mathcal{M}_{0,3}$ .

Take the 3 marked points to be  $(0, 1, \infty)$ , chose 3 perimeter lengths  $L_0, L_1, L_\infty$ . Then the Strebel differential is:


$$\Omega(x) = -\frac{L_\infty^2 x^2 - (L_\infty^2 + L_0^2 - L_1^2)x + L_0^2}{x^2(x-1)^2} dx^2 = -L_\infty^2 \frac{(x-a)(x-b)}{x^2(x-1)^2} dx^2$$

The vertices are located at the zeroes  $a, b$  of  $\Omega$ :

$$a, b = \frac{1}{2L_\infty^2} \left( L_\infty^2 + L_0^2 - L_1^2 \pm \sqrt{L_0^4 + L_1^4 + L_\infty^4 - 2L_0^2L_1^2 - 2L_0^2L_\infty^2 - 2L_1^2L_\infty^2} \right)$$



or



Remark:  $\Omega$ =stress energy tensor of classical Liouville theory.

# Back to Quantum Gravity

We thus have

$$\int \prod_i dL_i \int_{\mathcal{M}_{g,n}} \text{observable} = \sum_G \frac{1}{\#\text{Aut}(G)} \int_{\mathbb{R}_+^{\#\text{edges}}} \text{observable} \prod_e d\ell_e$$

Theorem Kontsevich 92: The measure  $\prod_e d\ell_e$  is topological:

$$(3g-3+n)! 2^{5g-5+2n} \prod_e d\ell_e = \left( \sum_i L_i^2 c_1(T^* p_i) \right)^{3g-3+n} \prod_{i=1}^n dL_i$$

$\psi_i = c_1(T^* p_i)$  = Chern class of cotangent bundle at marked point  $p_i$ .

$$\psi_i = \sum_{e < e' \text{ around face } i} d\frac{\ell_e}{L_i} \wedge d\frac{\ell_{e'}}{L_i}$$

$\psi_i$  is a 2-form on the cotangent space of  $\mathcal{M}_{g,n} \times \mathbb{R}_+^n$ .

# Laplace transform

Multiply by  $e^{-\sum_{i=1}^n \lambda_i L_i}$  and integrate over  $\ell_e$ .

$$e^{-\sum_i \lambda_i L_i} = e^{-\sum_{e=(e_+, e_-) \in \text{edges}} \ell_e (\lambda_{e_+} + \lambda_{e_-})}$$

For an observable polynomial in the  $\ell_e$ 's:

$$\sum_G \int d\ell_e \prod_e \ell_e^{k_e} e^{-\sum_i \lambda_i L_i} = \sum_G \prod_{e=(e_+, e_-)} \frac{k_e!}{(\lambda_{e_+} + \lambda_{e_-})^{k_e+1}}$$

In principle, any measure  $d\mu$  on  $\mathcal{M}_{g,n}$  can be realized as a function of the edge lengths  $f(\{\ell_e\}) \prod_e d\ell_e$ , or in Laplace transform as a function of the  $\lambda_f$ 's.

In particular: discrete measures can be realized. Embedding  $\mathbb{M}_{g,n} \subset \mathcal{M}_{g,n}$  with the discrete measure, can be realized by a suitable system of weights  $w(G; \{\lambda_f\})$  on graphs  $G \in \mathcal{G}_{g,n}$ .

We shall consider the following set of graphs:

$\mathcal{G}_{g,n}$ =set of graphs of genus  $g$ , with  $n$  labeled marked faces.

- Marked face  $i$  carries a variable  $z_i$ .
- Unmarked faces carry a variable in a set  $\{\lambda_1, \dots, \lambda_N\}$ .
- an edge  $e$  between two faces  $e_+, e_-$ , has weight


$$\frac{1}{(\text{variable}(e_+) + \text{variable}(e_-))}.$$

- 3-valent vertices: weight =1.
- 1-valent vertices only for marked faces: weight =-1.

Weight of Feynman type (Wick's theorem type):

$$w(G) = N^{-\#\text{unmarked faces}} \prod_{e=(e_+, e_-) \in \text{edges}} \frac{1}{\lambda_{e_+} + \lambda_{e_-}} \prod_{v \in \text{vertices}} \text{weight}(v)$$

Example:  $G \in \mathcal{G}_{0,1}$  with one marked face (exterior), and one unmarked face (interior):




$$\rightarrow \text{weight} = \frac{-1}{N} \frac{1}{z+z} \frac{1}{z+\lambda_a}$$

# Generating functions

Define : 
$$W_{g,n}(z_1, \dots, z_n) = \sum_{G \in \mathcal{G}_{g,n}} w(G) - \delta_{g,0} \delta_{n,1} z_1$$


where we sum over graphs with  $n$  labeled marked faces, each having one 1-valent vertex, and with  $z_i$  in marked face  $i$ .

Example: graphs contributing to  $W_{0,1}(z)$  with up to 5 edges:



$$\begin{aligned} W_{0,1}(z) &= -z + \frac{1}{N} \sum_{a=1}^N \frac{-1}{2z(z + \lambda_a)} \\ &\quad - \frac{1}{N^2} \sum_{a,b} \frac{1}{(2z)^3 (z + \lambda_a)(z + \lambda_b)} + \frac{1}{(2z)^2, (z + \lambda_a)^2 (z + \lambda_b)} \end{aligned}$$

## Other example: $W_{0,2}$ : graphs with up to 7 edges






$$W_{0,2}(z, z') = \frac{1}{4zz'(z + z')^2} + 0 + \dots \quad (0.1)$$

Remark: huge cancellations.


# Tutte's recursion

Tutte's recursion = deleting edges.



- consider the 1-valent vertex of face 1,
- it is attached to an edge  $\frac{1}{2z_1}$ ,
- which is attached to a 3-valent vertex.
- delete the right edge of the 3-valent vertex:
  - the face on the other side is the same face ( $z_1$ ) and the graph gets disconnected, or  $g \rightarrow g - 1$
  - the face on the other side is another face, marked or unmarked,

# Tutte's recursion



Case: – the face on the other side is another face, marked or unmarked,

Lemma: if  $F(z) = \prod_{i=1}^k \frac{1}{z+c_i}$ , then

$$\frac{F(z) - F(a)}{z - a} = \sum_{j=1}^k \prod_{i=1}^j \frac{1}{z + c_i} \quad \prod_{i=j}^k \frac{1}{a + c_i}$$

# Tutte's recursion

It follows:

Theorem (Tutte's recursion formula):

$$\begin{aligned} & 2z(W_{g,n}(z, z_2, \dots, z_n) + z\delta_{g,0}\delta_{n,1}) \\ = & W_{g-1,n+1}(z, z, z_2, \dots, z_n) \\ & + \sum_{h+h'=g} \sum_{I \cup I' = \{z_2, \dots, z_n\}} W_{h,1+|I|}(z, I) W_{h',1+|I'|}(z, I') \\ & + \frac{1}{N} \sum_{a=1}^N \frac{W_{g,n}(z, z_2, \dots, z_n) - W_{g,n}(\lambda_a, z_2, \dots, z_n)}{z^2 - \lambda_a^2} \\ & + \sum_{j=2}^n \frac{1}{2z_j} \frac{d}{dz_j} \frac{W_{g,n-1}(z, z_2, \dots, \hat{z}_j, \dots, z_n) - W_{g,n-1}(z_2, \dots, z_n)}{z^2 - z_j^2} \end{aligned}$$

# Disc amplitude


Planar maps with 1 marked face (rooted planar map):

$$2z(W_{0,1}(z) + z) = (W_{0,1}(z) + z)^2 + \frac{1}{N} \sum_{a=1}^N \frac{W_{0,1}(z) - W_{0,1}(\lambda_a)}{z^2 - \lambda_a^2}$$

**Theorem:** Solution (we assume  $t_1 = \frac{1}{N} \sum_{a=1}^N \frac{1}{\lambda_a} = 0$ ):

$$W_{0,1}(z) = -z - \frac{1}{N} \sum_{a=1}^N \frac{1}{2z(z + \lambda_a)}$$

i.e. only these graphs contribute:



# Annulus amplitude


Planar maps with 2 marked faces:

$$\begin{aligned} 2z W_{0,2}(z, z') &= 2 W_{0,1}(z) W_{0,2}(z, z') \\ &\quad + \frac{1}{N} \sum_{a=1}^N \frac{W_{0,2}(z, z') - W_{0,1}(\lambda_a, z')}{z^2 - \lambda_a^2} \\ &\quad + \frac{1}{2z'} \frac{d}{dz'} \frac{W_{0,1}(z) - W_{0,1}(z')}{z^2 - z'^2} \end{aligned}$$

**Theorem:** Solution (we assume  $t_1 = \sum_{a=1}^N \frac{1}{\lambda_a} = 0$ ):

$$W_{0,2}(z, z') = \frac{1}{4zz' (z + z')^2}$$

i.e. only this graph contributes:



# (0,3) amplitude

Planar maps with 3 marked faces:

$$\begin{aligned} 2z W_{0,3}(z_1, z_2, z_3) = & 2 W_{0,1}(z_1) W_{0,3}(z_1, z_2, z_3) + 2 W_{0,2}(z_1, z_2) W_{0,2}(z_1, z_3) \\ & + \frac{1}{N} \sum_{a=1}^N \frac{W_{0,3}(z_1, z_2, z_3) - W_{0,3}(\lambda_a, z_2, z_3)}{z_1^2 - \lambda_a^2} \\ & + \frac{1}{2z_2} \frac{d}{dz_2} \frac{W_{0,2}(z_1, z_3) - W_{0,2}(z_2, z_3)}{z_1^2 - z_2^2} \\ & + \frac{1}{2z_3} \frac{d}{dz_3} \frac{W_{0,2}(z_1, z_2) - W_{0,2}(z_2, z_3)}{z_1^2 - z_3^2} \end{aligned}$$

**Theorem:** Solution (we assume  $t_1 = \frac{1}{N} \sum_{a=1}^N \frac{1}{\lambda_a} = 0$ ):

$$W_{0,3}(z_1, z_2, z_3) = \frac{1}{8(2-t_3) z_1^3 z_2^3 z_3^3}$$

where  $t_k = \frac{1}{N} \sum_{a=1}^N \lambda_a^{-k}$ .

# (1,1) amplitude

Genus 1 maps with 1 marked face:

$$2z W_{1,1}(z) = 2 W_{0,1}(z) W_{1,1}(z) + W_{0,2}(z, z) + \frac{1}{N} \sum_{a=1}^N \frac{W_{1,1}(z) - W_{1,1}(\lambda_a)}{z^2 - \lambda_a^2}$$

**Theorem:** Solution (we assume  $t_1 = \frac{1}{N} \sum_{a=1}^N \frac{1}{\lambda_a} = 0$ ):

$$W_{1,1}(z) = \frac{1}{16(2-t_3)} \frac{1}{z^5} + \frac{t_5}{16(2-t_3)^2} \frac{1}{z^3}$$

where  $t_k = \frac{1}{N} \sum_{a=1}^N \lambda_a^{-k}$ .

# Topological recursion

From Tutte's equation, one proves by recursion ( $W_{g,n}$  = rational with poles at  $z_i = 0$ , expand at 0 instead of  $\infty$ ):

Theorem (the  $W_{g,n}$ 's satisfy the topological recursion (EO07))

$$\begin{aligned} & W_{g,n}(z_1, z_2, \dots, z_n) \\ = & \operatorname{Res}_{z \rightarrow 0} K(z_1, z) \left[ W_{g-1, n+1}(z, z, z_2, \dots, z_n) \right. \\ & + \sum'_{h+h'=g, I \cup I'=\{z_2, \dots, z_n\}} \left. W_{h, 1+|I|}(z, I) W_{h', 1+|I'|}(z, I') \right] dz \end{aligned}$$

where  $K(z_1, z) = \frac{z}{z_1} \frac{1}{z_1^2 - z^2} (2 - \sum_{k=0}^{\infty} t_{2k+3} z^{2k})^{-1}$  ,  $t_k = \frac{1}{N} \sum_{a=1}^N \lambda_a^{-k}$

This theorem computes recursively  $W_{g,n}$  in terms of  $W_{g',n'}$  with  $2g' + n' - 2 < 2g + n - 2$ .

# Topological recursion

Example of computation:

$$\begin{aligned} W_{1,1}(z_1) &= \operatorname{Res}_{z \rightarrow 0} K(z_1, z) W_{0,2}(z, z) dz \\ &= \operatorname{Res}_{z \rightarrow 0} \frac{z dz}{z_1} \frac{1}{z_1^2 - z^2} (2 - t_3 - t_5 z^2 + \dots)^{-1} \frac{1}{4z^2(z+z)^2} \\ &= \frac{1}{16(2-t_3)z_1^3} \operatorname{Res}_{z \rightarrow 0} \frac{dz}{z^3} \left(1 + \frac{z^2}{z_1^2} + O(z^4)\right) \left(1 + \frac{t_5}{2-t_3} z^2\right) \\ &= \frac{1}{16(2-t_3)z_1^3} \left( \frac{1}{z_1^2} + \frac{t_5}{(2-t_3)} \right) \end{aligned}$$

→ agrees with known result



By recursion on  $g$  and  $n$ , we find:

## Lemma

$$W_{g,n} = \text{Polynomial} \in \mathbb{Q}[1/z_i, 1/(2 - t_3), t_k]$$

$$\text{where } t_k = \frac{1}{N} \sum_a \lambda_a^{-k}.$$

Instead of  $t_k$ 's, we prefer to use their Schur transforms  $\hat{t}_k$ 's:

$$\text{Def : } \sum_{k \geq 0} \hat{t}_k u^{-k} = -\ln \left( 2 - \sum_{k \geq 0} (2k+1)!! t_{2k+3} u^{-k} \right)$$

For example we have

$$\hat{t}_0 = -\ln(2 - t_3), \hat{t}_1 = \frac{3t_5}{2-t_3}, \hat{t}_2 = \frac{15t_7}{2-t_3} + \frac{9t_5^2}{2(2-t_3)^2}, \text{ etc...}$$

By recursion on  $g$  and  $n$ , we find:

## Lemma

$$W_{g,n} = \text{Polynomial} \in \mathbb{Q}[1/z_i, 1/(2 - t_3), t_k]$$

$$\text{where } t_k = \frac{1}{N} \sum_a \lambda_a^{-k}.$$

Instead of  $t_k$ 's, we prefer to use their Schur transforms  $\hat{t}_k$ 's:

$$\text{Def : } \sum_{k \geq 0} \hat{t}_k u^{-k} = -\ln \left( 2 - \sum_{k \geq 0} (2k+1)!! t_{2k+3} u^{-k} \right)$$

For example we have

$$\hat{t}_0 = -\ln(2 - t_3), \hat{t}_1 = \frac{3t_5}{2-t_3}, \hat{t}_2 = \frac{15t_7}{2-t_3} + \frac{9t_5^2}{2(2-t_3)^2}, \text{ etc...}$$

## Lemma

$$W_{g,n} = \frac{(2-t_3)^{2-2g-n}}{\prod_i 2z_i^3} \text{ Polynomial } \in \mathbb{Q}[1/z_i^2, \{\hat{t}_k\}_{k \geq 1}]^{\deg=3g-3+n}$$

(homogeneity  $\deg(1/z_i^2) = 1$ ,  $\deg(\hat{t}_k) = k$ )

Instead of  $t_k$ 's, we prefer to use their Schur transforms  $\hat{t}_k$ 's:

$$\text{Def : } \sum_{k \geq 0} \hat{t}_k u^{-k} = -\ln \left( 2 - \sum_{k \geq 0} (2k+1)!! t_{2k+3} u^{-k} \right)$$

For example we have

$$\hat{t}_0 = -\ln(2-t_3), \hat{t}_1 = \frac{3t_5}{2-t_3}, \hat{t}_2 = \frac{15t_7}{2-t_3} + \frac{9t_5^2}{2(2-t_3)^2}, \text{ etc...}$$

# Topological recursion and Virasoro constraints

## Lemma

$$W_{g,n} = \frac{(2-t_3)^{2-2g-n}}{\prod_i 2z_i^3} \text{ Polynomial } \in \mathbb{Q}[1/z_i^2, \{\hat{t}_k\}_{k \geq 1}]^{\deg=3g-3+n}$$

(homogeneity  $\deg(1/z_i^2) = 1$ ,  $\deg(\hat{t}_k) = k$ )

$$\begin{aligned} W_{g,n} &= \frac{(2-t_3)^{2-2g-n}}{\prod_i 2z_i} \sum_{d_1+\dots+d_n+\sum_{k \geq 1} kn_k=3g-3+n} \\ &\quad \prod_i \frac{(2d_i+1)!!}{z_i^{2d_i+2}} \prod_{k \geq 1} \frac{\hat{t}_k^{n_k}}{n_k!} C_{g,n}(d_1, \dots, d_n; n_1, n_2, n_3, \dots) \end{aligned}$$

Witten's Notation (intersection numbers):

$$C_{g,n}(d_1, \dots, d_n; n_1, \dots, n_{3g-3+n}) \stackrel{\text{def}}{=} \langle \tau_{d_1} \dots \tau_{d_n} \kappa_1^{n_1} \kappa_2^{n_2} \kappa_3^{n_3} \dots \rangle_g$$



# Topological recursion and Virasoro constraints

We thus write:

$$W_{g,n} = \frac{(2-t_3)^{2-2g-n}}{\prod_i 2z_i} \sum_{d_1+\dots+d_n+\sum_k kn_k=3g-3+n} \prod_i \frac{(2d_i+1)!!}{z_i^{2d_i+2}} \prod_{k\geq 1} \frac{\hat{t}_k^{n_k}}{n_k!} C_{g,n}(d_1, \dots, d_n; n_1, n_2, n_3, \dots)$$

Witten's Notation (intersection numbers):

$$C_{g,n}(d_1, \dots, d_n; n_1, n_2, \dots) \stackrel{\text{def}}{=} <\tau_{d_1} \dots \tau_{d_n} \kappa_1^{n_1} \kappa_2^{n_2} \kappa_3^{n_3} \dots>_g$$

and we define  $C_{g,n}(\{d_i\}; \{n_i\}) = <\prod_i \tau_{d_i} \prod_i \kappa_i^{n_i}>_g = 0$  if  $\sum_i d_i + \sum_k kn_k \neq 3g - 3 + n$ .

# Topological recursion and Virasoro constraints

We thus write:

$$W_{g,n} = \frac{(2-t_3)^{2-2g-n}}{\prod_i 2z_i} \sum_{d_1, \dots, d_n, n_1, n_2, \dots,} \\ \prod_i \frac{(2d_i + 1)!!}{z_i^{2d_i+2}} \prod_{k \geq 1} \frac{\hat{t}_k^{n_k}}{n_k!} C_{g,n}(d_1, \dots, d_n; n_1, n_2, n_3, \dots)$$

Witten's Notation (intersection numbers):

$$C_{g,n}(d_1, \dots, d_n; n_1, n_2, \dots) \stackrel{\text{def}}{=} <\tau_{d_1} \dots \tau_{d_n} \kappa_1^{n_1} \kappa_2^{n_2} \kappa_3^{n_3} \dots>_g$$

and we define  $C_{g,n}(\{d_i\}; \{n_i\}) = <\prod_i \tau_{d_i} \prod_i \kappa_i^{n_i}>_g = 0$  if  $\sum_i d_i + \sum_k k n_k \neq 3g - 3 + n$ .

# Topological recursion and Virasoro constraints

We thus write:

$$W_{g,n} = \frac{(2-t_3)^{2-2g-n}}{\prod_i 2z_i} \sum_{d_1, \dots, d_n, n_1, n_2, \dots,} \\ \prod_i \frac{(2d_i+1)!!}{z_i^{2d_i+2}} \prod_{k \geq 1} \frac{\hat{t}_k^{n_k}}{n_k!} < \tau_{d_1} \dots \tau_{d_n} \prod_k \kappa_k^{n_k} >_g$$

Witten's Notation (intersection numbers):

$$C_{g,n}(d_1, \dots, d_n; n_1, n_2, \dots) \stackrel{\text{def}}{=} < \tau_{d_1} \dots \tau_{d_n} \kappa_1^{n_1} \kappa_2^{n_2} \kappa_3^{n_3} \dots >_g$$

and we define  $C_{g,n}(\{d_i\}; \{n_i\}) = < \prod_i \tau_{d_i} \prod_i \kappa_i^{n_i} >_g = 0$  if  $\sum_i d_i + \sum_k k n_k \neq 3g - 3 + n$ .

# Topological recursion and Virasoro constraints

We thus write:

$$W_{g,n} = \frac{(2-t_3)^{2-2g-n}}{\prod_i 2z_i} \sum_{d_1, \dots, d_n} \prod_i \frac{(2d_i+1)!!}{z_i^{2d_i+2}} \langle \tau_{d_1} \dots \tau_{d_n} e^{\sum_k \hat{t}_k \kappa_k} \rangle_g$$

Witten's Notation (intersection numbers):

$$C_{g,n}(d_1, \dots, d_n; n_1, n_2, \dots) \stackrel{\text{def}}{=} \langle \tau_{d_1} \dots \tau_{d_n} \kappa_1^{n_1} \kappa_2^{n_2} \kappa_3^{n_3} \dots \rangle_g$$

and we define  $C_{g,n}(\{d_i\}; \{n_i\}) = \langle \prod_i \tau_{d_i} \prod_i \kappa_i^{n_i} \rangle_g = 0$  if  $\sum_i d_i + \sum_k k n_k \neq 3g - 3 + n$ .

# Topological recursion and Virasoro constraints

- Example  $(g, n) = (1, 1)$ :

$$W_{1,1} = \frac{(2 - t_3)^{-1}}{2z} \left( \frac{1}{24} \frac{3}{z^4} + \frac{1}{24} \frac{\hat{t}_1}{z^2} \right)$$

i.e.

$$C_{1,1}(1; 0, 0, \dots) = \langle \tau_1 \rangle_1 = \frac{1}{24}$$

$$C_{1,1}(0; 1, 0, \dots) = \langle \tau_0 \kappa_1 \rangle_1 = \frac{1}{24}$$

- Example  $(g, n) = (0, 3)$ :

$$W_{0,3} = \frac{(2 - t_3)^{-1}}{8 z_1^3 z_2^3 z_3^3}$$

i.e.

$$C_{0,3}(0, 0, 0; 0, 0, \dots) = \langle \tau_0 \tau_0 \tau_0 \rangle_0 = 1$$

# Topological recursion and Virasoro constraints

Notational remark: Algebraic topology

$\langle \tau_{d_1} \dots \tau_{d_n} \kappa_1^{n_1} \kappa_2^{n_2} \kappa_3^{n_3} \dots \rangle_g$  is not only a notation, it is actually an integral of cohomology classes.

$$\langle \tau_{d_1} \dots \tau_{d_n} \kappa_1^{n_1} \kappa_2^{n_2} \kappa_3^{n_3} \dots \rangle_g = \int_{\overline{\mathcal{M}}_{g,n}} \prod_{i=1}^n \tau_{d_i} \prod_k \kappa_k^{n_k}$$

This can be proved [Kontsevich 1992, E 2008] from  $\prod_e d\ell_e \propto (\sum_i L_i^2 \psi_i)^{3g-3+n} \prod_i dL_i$ , and Laplace transform.

The classes  $\tau$  and  $\kappa$  are:

- $\tau_d = \psi^d$ , where  $\psi = c_1(T^*p) =$  1st Chern class of the cotangent bundle at marked point  $p$ .  $\tau_d$  is called "**Witten's class**".
- $\kappa_d$  = Schur transform of  $\tau_{d+1}$ , is called "**Mumford's class**".

# Topological recursion and Virasoro constraints

The topological recursion (implied by Tutte's recursion) implies some relations among the coefficients

$$C_{g,n} = \langle \tau_{d_1} \dots \tau_{d_n} \prod_k \kappa_K^{n_k} \rangle_g:$$

## Theorem (Virasoro)

*The intersection numbers satisfy "Witten's Virasoro conjecture" [EHX1997].*

This theorem was proved recently [Mulase, Safnuk 2006], [Liu, Xu...], using Mirzakhani's results from hyperbolic geometry, proof long and difficult.

Here we find a combinatorial proof from Tutte's recursion.

# Topological recursion and Virasoro constraints

Case where all  $t_k = 0$ .

In that case we have

$$W_{g,n}(z_1, \dots, z_n) = \frac{2^{2-2g-n}}{\prod_i 2z_i} \sum_{d_1, \dots, d_n} \prod_{i=1}^n \frac{(2d_i + 1)!!}{z_i^{2d_i+2}} < \tau_{d_1} \dots \tau_{d_n} >_g$$

Define:

$$\psi_\hbar(z) = e^{\sum_{g,n} \frac{\hbar^{2g-2+n}}{n!} \int_\infty^z \dots \int_\infty^z W_{g,n} dz_1 \dots dz_n}$$

Using Tutte's equations, one can prove that

Theorem

$\psi(z)$  satisfies Airy equation with respect to  $x = z^2$ :

$$\hbar^2 \frac{d^2}{dx^2} \psi = x \psi$$

This is another formulation of KdV Witten's conjecture.



# 3. Higher spin

# Witten's r-spin conjecture

$\mathcal{M}_{g,n}^{1/r}$  = moduli space spin  $r$  bundles over Riemann surfaces of genus  $g$  with  $n$  marked points.

→ analogous of colored maps in the discrete version, Potts model, Tutte's chromatic polynomials.

Witten's conjecture: integrals of Chern classes over  $\mathcal{M}_{g,n}^{1/r}$ , make a Tau-function for the r-KdV hierarchy.

Proved [Faber, Shadrin, Zvonkine 2006].

But the Virasoro conjecture yet unproved, also many Faber-Pandarpande conjectures yet unproved.

# Witten's r-spin conjecture

Kontsevich's r-spin combinatorial model:

Consider a "potential"  $V(z) = \sum_{k=1}^{r+1} \frac{v_k}{k} z^k$ .

Define for  $2 \leq m \leq r + 1$ :

$$\mathcal{V}_m(z_1, \dots, z_m) = - \sum_{i=1}^m \frac{V'(z_i)}{\prod_{j \neq i} (z_i - z_j)} = - \sum_{j \geq m} v_j \sum_{k_1 + \dots + k_m = j-m} \prod_i z_i^{k_i}$$

and  $\mathcal{P}(z_1, z_2) = - \frac{1}{\mathcal{V}_2(z_1, z_2)} = \frac{z_1 - z_2}{V'(z_1) - V'(z_2)}$

Case  $r = 2$  and  $V(z) = z^3/3$  gives back Strebel's graphs

$$\mathcal{V}_3(z_1, z_2, z_3) = 1 \quad , \quad \mathcal{P}(z_1, z_2) = \frac{1}{z_1 + z_2}$$

# Witten's r-spin conjecture


$$W_{g,n}(z_1, \dots, z_n) = \sum_G \frac{1}{\#\text{Aut } G} \prod_{v=\text{vertex}} \mathcal{V}_{\text{valence}(v)}(\{\lambda_f\}_{f \mapsto v}) \prod_{e=\text{edges}} \mathcal{P}(\lambda_{e_+}, \lambda_{e_-})$$

(Most often we shall assume that

$$\sum_{a=1}^N \frac{\lambda_a^k}{V''(\lambda_a)} = 0 \quad \forall k \in [0, , r-2] \quad )$$

- The case  $r = 2$ ,  $V(z) = z^3/3$  is the Strebel-graphs model.
- The case  $r \geq 3$  and  $V(z) = z^{r+1}/(r+1)$  is the r-spin Strebel graphs model.

# Tutte's equations




Define  $W_0(z) = W_{0,1}(z) - z$  and for  $k \geq 1$ :

$$W_k(z; a_1, \dots, a_k) = \sum_{i=1}^k \frac{W_{0,1}(a_i) - W_{0,1}(z)}{(V'(a_i) - V'(z)) \prod_{j \neq i} (V'(a_i) - V'(a_j))}$$

We have Tutte's recursion (planar rooted map case):

$$V''(z) W_{0,1}(z) = \sum_{k \geq 2} \sum_{L_1, \dots, L_k} \mathcal{V}_{k+1+\sum_i \# L_i}(\overbrace{z, \dots, z}^{k+1}, L_1, \dots, L_k)$$
$$\prod_{i=1}^k W_k(z; L_i)$$

# Tutte's equations


$$W_3(z; a, b, c) =$$


Example  $r = 3$

$$\begin{aligned} V''(z) W_{0,1}(z) &= \mathcal{V}_4(z, z, z, z) W_0(z)^3 \\ &\quad + 2 \sum_a \mathcal{V}_4(z, z, z, \lambda_a) W_0(z) W_1(z; \lambda_a) \\ &\quad + \sum_{a,b} \mathcal{V}_4(z, z, \lambda_a, \lambda_b) W_2(z; \lambda_a, \lambda_b) \\ &\quad + \mathcal{V}_3(z, z, z) W_0(z)^2 + \sum_a \mathcal{V}_3(z, z, \lambda_a) W_1(z; \lambda_a) \end{aligned}$$

# Tutte's equations

Solution:

Disc :  $W_{0,1}(z) = -z - \sum_a \mathcal{V}_3(z, z, \lambda_a) \mathcal{P}(z, z) \mathcal{P}(z, \lambda_a)$



i.e. only these graphs contribute :

Annulus :  $W_{0,2}(z, z') = \frac{1}{V''(z) V''(z') (z - z')^2} - \frac{1}{(V'(z) - V'(z'))^2}$

# Tutte's equations

Tutte's equations imply topological recursion :

Conjecture (the  $W_{g,n}$ 's satisfy the topological recursion (EO07))

$$\begin{aligned} & W_{g,n}(z_1, z_2, \dots, z_n) \\ = & \operatorname{Res}_{z \rightarrow V''^{-1}(0)} K(z_1, z) \left[ W_{g-1, n+1}(z, \sigma(z), z_2, \dots, z_n) \right. \\ & + \sum'_{h+h'=g, I \sqcup I'=\{z_2, \dots, z_n\}} \left. W_{h, 1+|I|}(z, I) W_{h', 1+|I'|}(\sigma(z)z, I') \right] dz \end{aligned}$$

$$\text{where } K(z_1, z) = \frac{V''(z_1)}{(z_1 - z)} \frac{1}{W_{0,1}(z) - W_{0,1}(\sigma(z))}$$

$$\text{and where } V'(\sigma(z)) = V'(z).$$

This implies the r-spin Virasoro conjecture.

# Tutte's equations

Polynomiality

For  $2g + n - 2 > 0$ :

$$W_{g,n} \in \mathbb{Q}[z_i, 1/V''(z_i), 1/\hat{t}_{j,0}, \hat{t}_{j,k}]$$

where  $\sum_k \hat{t}_{j,k} u^{-k} = \ln \left( \frac{1}{N} \sum_a \int_{\gamma_j} \frac{dz}{z-a} e^{-u V'(z)} \right) \quad j \in [1, r-1]$

$$W_{g,n} = \sum_{\{j_i, d_i\}, \{n_{j,k}\}} \prod_{i=1}^n \frac{z_i^{j_i}}{V''(z_i)^{d_i}} < W_r \prod_{i=1}^n \tau_{j_i, d_i} e^{\sum_{j,k} \hat{t}_{j,k} \kappa_{j,k}} >_g$$

The coefficients are the r-spin intersection numbers

$$< W_r \prod_{i=1}^n \tau_{j_i, d_i} \prod_{j,k} \kappa_{j,k}^{n_{j,k}} >_g$$

# Tutte's equations

Chosing  $V'(z) = z^r - r\epsilon^r z$  and taking the limit  $\epsilon \rightarrow 0$ , implies Faber-Pandaripande's conjectures.

Chosing  $\hat{t}_{j,k} = 0$ , and

Define:

$$\psi_{\hbar}(z) = e^{\sum_{g,n} \frac{\hbar^{2g-2+n}}{n!} \int_{\infty}^z \dots \int_{\infty}^z W_{g,n} dz_1 \dots dz_n}$$

Using Tutte's equations, one can prove that

## Conjecture

$\psi(z)$  satisfies Airy equation with respect to  $x = z^r$ :

$$\hbar^r \frac{d^r}{dx^r} \psi = x \psi$$

This is another formulation of  $r$ -KdV Witten's conjecture.

# 5. Conclusion

## Conclusion:

- Strebel's graphs are a very promising tool for studying "2d quantum gravity". Not only the topological part.
- The combinatorics of Strebel's graphs satisfies Tutte's equations  $\Rightarrow$  topological recursion
- From the Tutte's equations and topological recursion, we can prove many non-trivial topological gravity properties, like Virasoro or r-KdV.

## Further prospects:

- study non-topological properties ? Example: the embedding into  $\mathbb{C}$  is function of  $\ell_e$ 's, in particular, the location of vertices or centers of faces are functions of  $\ell_e$ 's.

The end

Thank you for your attention

**Acknowledgments:**

This work was supported:

by the CRM Montréal, by CERN, by the Quebec government with the FQRNT.