

# Shortest Cycle Cover

André Raspaud

LaBRI  
Université Bordeaux I  
France


Graph Theory 2005, Nyborg, December 1-4, 2005




- ▶ A *circuit* is a connected 2-regular subgraph and a *cycle* is a union of edge-disjoint circuits. A *cycle cover* of a graph  $G$  is a family of cycles of  $G$  such that each edge of  $G$  is contained in at least one cycle of the family.
- ▶ The *length* of a cycle cover  $\mathcal{C}$ , denoted by  $l(\mathcal{C})$ , is the sum of the lengths (number of edges) of all the cycles in  $\mathcal{C}$ .




# Example



# Example



# Example



- ▶ Given a graph  $G$  and an integer  $k$ , does  $G$  have a cycle cover of total length at most  $k$ ?
- ▶ Thomassen ('92) proved that the SCCP is a NP-complete problem








- ▶ Given a graph  $G$  and an integer  $k$ , does  $G$  have a cycle cover of total length at most  $k$ ?
- ▶ Thomassen ('92) proved that the SCCP is a NP-complete problem



## Conjecture

Every bridgeless graph  $G = (V, E)$  has a cycle cover  $\mathcal{C}$  such that:

$$l(\mathcal{C}) \leq \frac{7}{5}|E|.$$



$$l(\mathcal{C}) = 21$$



A *cycle double cover* **CDC** of a graph  $G = (V, E)$  is a family of cycles of  $G$  such that every edge appears in exactly two cycles of this family.

The following double cover conjecture was proposed by several authors.

**Conjecture (Szekeres '73, Seymour '79)**

*Every bridgeless graph has a cycle double cover.*



Jamshi and Tarsi proved ('92):

## Theorem

*If the Conjecture concerning the existence of a cycle cover  $\mathcal{C}$  of length at most  $\frac{7}{5}|E|$  is true then the Cycle double cover conjecture is true.*

First bound : Itai, Rodeh '78:

## Theorem

*Every bridgeless graph  $G = (V, E)$  has a cycle cover  $\mathcal{C}$  such that:*

$$l(\mathcal{C}) \leq |E| + 2|V|\log|V|.$$




## Definition

Let  $M$  be an abelian group,  $\vec{G}$  be an oriented graph,

$\phi : E(\vec{G}) \rightarrow M$  is a *flow* or an  $M$ -*flow* if :

$\sum_{e \in \Delta^+(v)} \phi(e) = \sum_{e \in \Delta^-(v)} \phi(e)$  for every vertex  $v \in V(\vec{G})$ .



If  $M = \mathbb{Z}_2^k$  the condition is  $\sum_{e \in \Delta(v)} \phi(e) = 0$



## Definition

- ▶ The *support*  $\sigma(\phi)$  of a flow  $\phi$  in a digraph  $G = (V, E)$  is the set of arcs  $e$  such that  $\phi(e) \neq 0$ .
- ▶ If  $\sigma(\phi) = E$ ,  $\phi$  is a  $M$ -nowhere zero flow ( $M$ -NZF).
- ▶ A  $\mathbb{Z}$ -flow is a  $k$ -flow if  $\phi(E) \subset [-(k-1), 0 \cup] 0, k-1]$
- ▶  $B \subseteq M$ ,  $-B = B$   
 $\phi$  is a  $B$ -flow,  
 if  $\phi : E \rightarrow M$  is a flow s.t.  $\phi(E) \subseteq B$

If  $M = \mathbb{Z}_2$ ,  $\sigma(\phi)$  is a cycle.



## Definition

- ▶ The *support*  $\sigma(\phi)$  of a flow  $\phi$  in a digraph  $G = (V, E)$  is the set of arcs  $e$  such that  $\phi(e) \neq 0$ .
- ▶ If  $\sigma(\phi) = E$ ,  $\phi$  is a  $M$ -nowhere zero flow ( $M$ -NZF).
- ▶ A  $\mathbb{Z}$ -flow is a  $k$ -flow if  $\phi(E) \subset [-(k-1), 0] \cup [0, k-1]$
- ▶  $B \subseteq M$ ,  $-B = B$   
 $\phi$  is a  $B$ -flow,  
 if  $\phi : E \rightarrow M$  is a flow s.t.  $\phi(E) \subseteq B$

If  $M = \mathbb{Z}_2$ ,  $\sigma(\phi)$  is a cycle.



## Definition

- ▶ The *support*  $\sigma(\phi)$  of a flow  $\phi$  in a digraph  $G = (V, E)$  is the set of arcs  $e$  such that  $\phi(e) \neq 0$ .
- ▶ If  $\sigma(\phi) = E$ ,  $\phi$  is a  $M$ -nowhere zero flow ( $M$ -NZF).
- ▶ A  $\mathbb{Z}$ -flow is a  $k$ -flow if  $\phi(E) \subset [-(k-1), 0 \cup] 0, k-1]$
- ▶  $B \subseteq M$ ,  $-B = B$   
 $\phi$  is a  $B$ -flow,  
 if  $\phi : E \rightarrow M$  is a flow s.t.  $\phi(E) \subseteq B$

If  $M = \mathbb{Z}_2$ ,  $\sigma(\phi)$  is a cycle.



## Definition

- ▶ The *support*  $\sigma(\phi)$  of a flow  $\phi$  in a digraph  $G = (V, E)$  is the set of arcs  $e$  such that  $\phi(e) \neq 0$ .
- ▶ If  $\sigma(\phi) = E$ ,  $\phi$  is a  $M$ -nowhere zero flow ( $M$ -NZF).
- ▶ A  $\mathbb{Z}$ -flow is a  $k$ -flow if  $\phi(E) \subset [-(k-1), 0] \cup [0, k-1]$
- ▶  $B \subseteq M$ ,  $-B = B$   
 $\phi$  is a  $B$ -flow,  
 if  $\phi : E \rightarrow M$  is a flow s.t.  $\phi(E) \subseteq B$

If  $M = \mathbb{Z}_2$ ,  $\sigma(\phi)$  is a cycle.



## Theorem (Tutte '50)

*Let  $G$  be a digraph. For every  $k \geq 2$ , the following conditions are equivalent:*

- 1. There exists a  $\mathbb{Z}_k$ -NZF in  $G$ .*
- 2. For any Abelian group  $M$  of order  $k$ , there exists a  $M$ -NZF in  $G$ .*
- 3. There exists a  $k$ -NZF in  $G$ .*





## Conjecture (Tutte '54)

*Every bridgeless graph without 3-cuts has a 3-NZF.*

## Theorem (Jaeger'79)

*Every bridgeless graph without 3-cuts has a 4-NZF.*

## Conjecture (Tutte '54)

*Every bridgeless graph has a 5-NZF.*

## Theorem

*Every bridgeless graph has:*

- 1. a 8-NZF (Jaeger '79)*
- 2. a 6-NZF (Seymour '81)*



## Conjecture (Tutte '54)

*Every bridgeless graph without 3-cuts has a 3-NZF.*

## Theorem (Jaeger'79)

*Every bridgeless graph without 3-cuts has a 4-NZF.*

## Conjecture (Tutte '54)

*Every bridgeless graph has a 5-NZF.*

## Theorem

*Every bridgeless graph has:*

1. *a 8-NZF (Jaeger '79)*
2. *a 6-NZF (Seymour '81)*



## Conjecture (Tutte '54)

*Every bridgeless graph without 3-cuts has a 3-NZF.*

## Theorem (Jaeger'79)

*Every bridgeless graph without 3-cuts has a 4-NZF.*

## Conjecture (Tutte '54)

*Every bridgeless graph has a 5-NZF.*

## Theorem

*Every bridgeless graph has:*

1. *a 8-NZF (Jaeger '79)*
2. *a 6-NZF (Seymour '81)*



## Conjecture (Peissmann '81, Celmins '84 )

- ▶ *For every graph with no cut-edge, there is a list of five cycles so that every edge is contained in exactly two.*
- ▶ *Every graph with no cut-edge has a B-flow for the set  $B \subseteq \mathbb{Z}_2^5$  consisting of those vectors with exactly two 1's.*



## Conjecture (Jaeger '88)

- ▶ *For every oriented graph with no cut-edge, there is a list of five 2-flows  $\phi_1, \phi_2, \dots, \phi_5$  with  $\sum_{i=1}^5 \phi_i = 0$  such that every edge is in the support of exactly two of these flows.*
- ▶ *Every graph with no cut-edge has a B-flow for the set  $B \subseteq \mathbb{Z}^5$  consisting of those vectors with exactly three 0's, one 1, and one  $-1$ .*



## Conjecture (Fulkerson,'71)


- ▶ *For every cubic graph with no cut-edge, there is a list of 6 perfect matchings so that every edge is contained in exactly two.*
- ▶ *Every graph with no cut-edge has a B-flow for the set  $B \subseteq \mathbb{Z}_2^6$  consisting of those vectors with exactly four 1's.*

## Conjecture (Berge,'79)

*The edges of any cubic graph with no cut-edge can be covered by 5 perfect matchings.*



	$C_1$	$C_2$	$C_3$	$C_4$	$C_5$	$C_6$
1	1	0	0	0	0	0
2	0	1	0	0	0	0
3	0	0	1	0	0	0
4	0	0	0	1	0	0
5	0	0	0	0	1	0
6	0	0	0	0	0	1
7	0	0	1	0	0	1
8	0	1	0	0	1	0
9	0	0	0	1	0	1
10	0	1	1	0	0	0
11	0	0	0	1	1	0
12	1	1	1	1	1	0
13	1	1	0	1	1	1
14	1	0	0	1	0	1
15	1	1	1	0	0	0



## Conjecture (Jaeger)

*Every bridgeless graph has a Petersen flow.*



## Theorem (Bermond, Jackson, Jaeger)

If  $G = (V, E)$  has a 4-NZF then it has a cycle cover  $\mathcal{C}$  such that:

$$l(\mathcal{C}) \leq \frac{4}{3}|E|$$

*Proof:* 4-NZF  $\iff \mathbb{Z}_2 \times \mathbb{Z}_2$ -NZF.

$\Phi, \phi_1, \phi_2$  s.t.  $\Phi(e) = (\phi_1(e), \phi_2(e))$ ,  $\phi_i$  is a  $\mathbb{Z}_2$ -flow,

$E = \sigma(\phi_1) \cup \sigma(\phi_2)$ .

$C_1 = \sigma(\phi_1)$ ,  $C_2 = \sigma(\phi_2)$ ,  $C_3 = C_1 \triangle C_2$





## Theorem (Bermond, Jackson, Jaeger)

If  $G = (V, E)$  has a 4-NZF then it has a cycle cover  $\mathcal{C}$  such that:

$$l(\mathcal{C}) \leq \frac{4}{3}|E|$$

*Proof:* 4-NZF  $\iff \mathbb{Z}_2 \times \mathbb{Z}_2$ -NZF.

$\Phi, \phi_1, \phi_2$  s.t.  $\Phi(e) = (\phi_1(e), \phi_2(e))$ ,  $\phi_i$  is a  $\mathbb{Z}_2$ -flow,

$E = \sigma(\phi_1) \cup \sigma(\phi_2)$ .

$C_1 = \sigma(\phi_1)$ ,  $C_2 = \sigma(\phi_2)$ ,  $C_3 = C_1 \triangle C_2$



## Theorem (Bermond, Jackson, Jaeger)

If  $G = (V, E)$  has a 4-NZF then it has a cycle cover  $\mathcal{C}$  such that:

$$l(\mathcal{C}) \leq \frac{4}{3}|E|$$

*Proof:* 4-NZF  $\iff \mathbb{Z}_2 \times \mathbb{Z}_2$ -NZF.

$\Phi, \phi_1, \phi_2$  s.t.  $\Phi(e) = (\phi_1(e), \phi_2(e))$ ,  $\phi_i$  is a  $\mathbb{Z}_2$ -flow,

$E = \sigma(\phi_1) \cup \sigma(\phi_2)$ .

$C_1 = \sigma(\phi_1)$ ,  $C_2 = \sigma(\phi_2)$ ,  $C_3 = C_1 \triangle C_2$



## Theorem (Bermond, Jackson, Jaeger)

If  $G = (V, E)$  has a 4-NZF then it has a cycle cover  $\mathcal{C}$  such that:

$$l(\mathcal{C}) \leq \frac{4}{3}|E|$$

*Proof:* 4-NZF  $\iff \mathbb{Z}_2 \times \mathbb{Z}_2$ -NZF.

$\Phi, \phi_1, \phi_2$  s.t.  $\Phi(e) = (\phi_1(e), \phi_2(e))$ ,  $\phi_i$  is a  $\mathbb{Z}_2$ -flow,

$E = \sigma(\phi_1) \cup \sigma(\phi_2)$ .

$C_1 = \sigma(\phi_1), C_2 = \sigma(\phi_2), C_3 = C_1 \triangle C_2$



## Theorem (Bermond, Jackson, Jaeger)

If  $G = (V, E)$  has a 4-NZF then it has a cycle cover  $\mathcal{C}$  such that:

$$l(\mathcal{C}) \leq \frac{4}{3}|E|$$

*Proof:* 4-NZF  $\iff \mathbb{Z}_2 \times \mathbb{Z}_2$ -NZF.

$\Phi, \phi_1, \phi_2$  s.t.  $\Phi(e) = (\phi_1(e), \phi_2(e))$ ,  $\phi_i$  is a  $\mathbb{Z}_2$ -flow,

$E = \sigma(\phi_1) \cup \sigma(\phi_2)$ .

$C_1 = \sigma(\phi_1)$ ,  $C_2 = \sigma(\phi_2)$ ,  $C_3 = C_1 \triangle C_2$



$C_1$	$C_2$	$C_3$
0	1	1
1	0	1
1	1	0

$$|C_1| + |C_2| + |C_3| = 2|E|$$

$\exists k \in \{1, 2, 3\}$  s.t.  $|C_k| \geq \frac{2}{3}|E|$

$\{C_i, C_j\}$  ( $i \neq j$ ) is a cycle cover

Let  $\mathcal{C} = \{C_i, C_j\}$ ,  $i \neq j \neq k$ :

$$I(\mathcal{C}) \leq 2|E| - \frac{2}{3}|E|$$

$$I(\mathcal{C}) \leq \frac{4}{3}|E|$$



$C_1$	$C_2$	$C_3$
0	1	1
1	0	1
1	1	0

$|C_1| + |C_2| + |C_3| = 2|E|$   
 $\exists k \in \{1, 2, 3\}$  s.t.  $|C_k| \geq \frac{2}{3}|E|$   
 $\{C_i, C_j\}$  ( $i \neq j$ ) is a cycle cover  
 Let  $\mathcal{C} = \{C_i, C_j\}, i \neq j \neq k$ :

$$I(\mathcal{C}) \leq 2|E| - \frac{2}{3}|E|$$

$$I(\mathcal{C}) \leq \frac{4}{3}|E|$$



$C_1$	$C_2$	$C_3$
0	1	1
1	0	1
1	1	0

$|C_1| + |C_2| + |C_3| = 2|E|$   
 $\exists k \in \{1, 2, 3\}$  s.t.  $|C_k| \geq \frac{2}{3}|E|$   
 $\{C_i, C_j\}$  ( $i \neq j$ ) is a cycle cover  
 Let  $C = \{C_i, C_j\}$ ,  $i \neq j \neq k$ :

$$I(C) \leq 2|E| - \frac{2}{3}|E|$$

$$I(C) \leq \frac{4}{3}|E|$$



$C_1$	$C_2$	$C_3$
0	1	1
1	0	1
1	1	0

$|C_1| + |C_2| + |C_3| = 2|E|$   
 $\exists k \in \{1, 2, 3\}$  s.t.  $|C_k| \geq \frac{2}{3}|E|$   
 $\{C_i, C_j\}$  ( $i \neq j$ ) is a cycle cover  
 Let  $\mathcal{C} = \{C_i, C_j\}, i \neq j \neq k$ :

$$I(\mathcal{C}) \leq 2|E| - \frac{2}{3}|E|$$

$$I(\mathcal{C}) \leq \frac{4}{3}|E|$$





$C_1$	$C_2$	$C_3$
0	1	1
1	0	1
1	1	0

$|C_1| + |C_2| + |C_3| = 2|E|$   
 $\exists k \in \{1, 2, 3\}$  s.t.  $|C_k| \geq \frac{2}{3}|E|$   
 $\{C_i, C_j\}$  ( $i \neq j$ ) is a cycle cover  
 Let  $\mathcal{C} = \{C_i, C_j\}, i \neq j \neq k$ :

$$I(\mathcal{C}) \leq 2|E| - \frac{2}{3}|E|$$

$$I(\mathcal{C}) \leq \frac{4}{3}|E|$$



$c(G)$  the size of a smallest cycle cover of a bridgeless graph  $G$ .

Authors	Results
Itai, Rodeh ('78)	$c(G) \leq  E  + 2 V  \log  V .$
Bermond, Jackson, Jaeger ('83)	$c(G) \leq \frac{5}{3} E .$
Alon, Tarsi ('85)	$c(G) \leq \frac{3}{2} E .$
Jamshy, R. , Tarsi ('87)	If $G$ has a 5-NZF $c_3(G) \leq \frac{8}{5} E .$
Jackson ('90)	If $G$ is $2k$ -edge-connected ( $k \geq 2$ ), then $c(G) \leq \frac{2k+2}{2k+1} E .$



Authors	Results
R. ('91)	If $G$ has a 4-NZF $G \neq K_4$ , then $c_2(G) \leq  E  +  V  - 3$ .
Fan, R. ('92)	If Fulkerson conjecture is true $c_3(G) \leq \frac{22}{15} E $ .
R. ('92)	If Petersen conjecture is true $c_4(G) \leq \frac{7}{5} E $ .
Fan ('93)	If 3-flow conjecture of Tutte is true $c_3(G) \leq \frac{7}{5} E $ .
Fan ('98)	$c(G) \leq  E  +  V  - 1$ .
Shu, Zhang ('04)	$G$ with odd edge-connectivity $r > 3$ $c_3(G) \leq \frac{r+1}{r} E $ .



Let  $G = (V, E)$  be a directed graph.  $p \geq 2q$  integers. A flow  $f : E \rightarrow \{-(p-1), \dots, p-1\}$  is a  $(p, q)$ -flow if for every edge  $e \in E$   $q \leq |f(e)| \leq p - q$ .


A  $(k, 1)$ -flow is a  $k$ -NZF



## Conjecture (R., Zhu '05)

If  $G = (V, E)$  has a  $(p, q)$ -flow then it has a cycle cover  $\mathcal{C}$  with


$$l(\mathcal{C}) \leq \frac{2(p-q)}{p} |E|$$



## Conjecture (R., Zhu '05)

If  $G = (V, E)$  has a  $(p, q)$ -flow then it has a cycle cover  $\mathcal{C}$  with


$$l(\mathcal{C}) \leq \frac{2(p-q)}{p} |E|$$



Let  $G = (V, E)$  be a graph.  $p \geq 2q$  integers.  $G = (V, E)$  is  $(p, q)$ -colorable if there exists a mapping  $f : V \rightarrow \{0, \dots, p-1\}$  s.t. for every edge  $xy \in E$  (Vince '88):

$$p \leq |f(x) - f(y)| \leq p - q$$

$$\chi_c(G) = \inf \left\{ \frac{p}{q} : \text{there is a } (p, q) \text{-coloring of } G \right\}$$




$(5, 2)$ -coloring of  $C_5$

Let  $G = (V, E)$  be a graph.  $p \geq 2q$  integers.  $G = (V, E)$  is  $(p, q)$ -colorable if there exists a mapping  $f : V \rightarrow \{0, \dots, p-1\}$  s.t. for every edge  $xy \in E$  (Vince '88):

$$p \leq |f(x) - f(y)| \leq p - q$$


$$\chi_c(G) = \inf\left\{\frac{p}{q} : \text{there is a } (p, q)\text{-coloring of } G\right\}$$



$(5, 2)$ -coloring of  $C_5$



$K_{\frac{p}{q}}$  be the graph with vertex set  $\{0, \dots, p-1\}$  ( $p \geq 2q$ ) in which  $ij$  is an edge if and only if  $q \leq |j-i| \leq p-q$ .



Circular clique  $K_{\frac{8}{3}}$

- ▶ A CDC  $\mathcal{C} = \{C_0, \dots, C_{p-1}\}$  is said to be  $(p, q)$ -colorable if the intersection graph  $G_{\mathcal{C}}$  is  $(p, q)$ -colorable.
- ▶  $V(G_{\mathcal{C}}) = \{0, \dots, p-1\}$  and  $ij \in E(\mathcal{C})$  iff  $C_i \cap C_j \neq \emptyset$ .
- ▶  $D_{p,q}$  is the subset of  $\mathbb{Z}_2^p$  consisting of elements  $(z_0, \dots, z_{p-1})$  with exactly two elements  $z_i, z_j$  equal to one and all the other equal to 0, with  $p \leq |j - i| \leq p - q$ .

### Proposition

*$G$  has a  $D_{p,q}$ -flow iff  $G$  has a  $(p, q)$ -colorable CDC*




## Theorem (R., Zhu '05)

*If  $G$  has a  $D_{p,q}$ -flow then it has a cycle cover  $\mathcal{C}$  with*


$$l(\mathcal{C}) \leq \frac{2(p-q)}{p} |E|$$




- ▶  $G$  has a  $D_{p,q}$ -flow  $\iff G_C$  the intersection graph of the corresponding CDC is a sub-graph of  $K_{\frac{p}{q}}$
- ▶  $e_{i,j} = \{e \in E : e \in C_i \cap C_j\}$ .
- ▶  $e \in e_{i,j} \iff e$  is of type  $ij$
- ▶ there is a bijection between the  $e_{i,j}$ 's and the edges of  $K_{\frac{p}{q}}$




- ▶  $C_{012} = \{C_3, C_4, C_5, C_6, C_7\}$  is a cycle cover of the graph
- ▶ The edges of types  $e_{i,j}$   $i \in \{0, 1, 2\}$ ,  $j \in \{3, 4, 5, 6, 7\}$  are covered once by this cycle cover  $C_{012}$ .
- ▶ The edges of the remaining types  $e_{3,7}, e_{3,6}, e_{4,7}$  are covered twice.



- ▶  $I(C_{012}) = |E| + |e_{3,7}| + |e_{3,6}| + |e_{4,7}|$
- ▶  $I(C_{012}) = |E| + |R_0|$  ( $R_0$  the edges of the remaining types)
- ▶  $K_{\frac{p}{q}}$  is vertex transitive.  $C_{012}$  is denoted by  $C_0$ .
- ▶  $C_i, i \in \{0, \dots, 7\}$  are 8 cycle covers ( $p$ )
- ▶  $I(C_i) = |E| + |R_i|$



- ▶ Each  $R_i$  contains 3 types of edges ( $\frac{(p-2q+1)(p-2q)}{2}$ )
- ▶  $\sum_0^7 I(C_i) = 8|E| + \sum_0^7 |R_i|$
- ▶  $\sum_0^{p-1} I(C_i) = p|E| + \sum_0^{p-1} |R_i|$
- ▶ each edges of types belonging to the remaining edges are counted twice ( $p - 2q$  times)
- ▶  $\sum_0^7 I(C_i) = 8|E| + 2|E|$
- ▶  $\sum_0^{p-1} I(C_i) = p|E| + (p - 2q)|E|$




- There is  $i \in \{0, \dots, 7\}$  ( $i \in \{0, \dots, p-1\}$ ) such that

$$I(C_i) \leq \frac{5}{4}|E|$$



$$I(C_i) \leq \frac{2(p-q)}{p}|E|$$







# Bicircular flows

## Theorem (Jaeger '81)

If  $G$  has a  $(2k + 1, k)$ -flow then it has a  $D_{2k+1, k}$ -flow

## Corollary

If  $G$  has a  $(2k + 1, k)$ -flow then it has a cycle cover  $\mathcal{C}$  with

$$l(\mathcal{C}) \leq \frac{2k + 2}{2k + 1} |E|$$

