

Partitions of graphs with bounded maximum average degree

André Raspaud

LaBRI
Université Bordeaux I
France

Graph Theory 2011
Kolding,
Denmark

Thursday April 28 - Sunday May 1, 2011

Proper k -coloring

Proper k -coloring

Let G be a graph and k ($k \geq 1$) an integer.

A proper k -coloring of G is a mapping $\phi : V(G) \rightarrow \{1, \dots, k\}$ such that:

- ▶ for every edge xy , $\phi(x) \neq \phi(y)$

In other words, a k -coloring of G is a partition V_1, V_2, \dots, V_k of $V(G)$ such that V_i is an independent set for every i , i.e., the subgraph induced by V_i has maximum degree zero.

d -improper k -coloring

Burr and Jacobson (1985), Cowen, Cowen, and Woodall (1986), Harary and Jones (1985).

d -improper k -coloring

Let G be a graph and k, d ($k, d \geq 1$) integers.

A d -improper k -coloring of G is a mapping $\psi : V(G) \rightarrow \{1, \dots, k\}$ such that :


- ▶ $\forall i, 1 \leq i \leq k$, $G[i]$ has a maximum degree at most d
- ▶ $G[i]$ is the subgraph induced by color i .

Every vertex v has at most d neighbors receiving the same color as v . We

denote: (d, d, \dots, d) -coloring = d -improper k -coloring

- ▶ a $(0, 0, 0, 0)$ -coloring is a proper 4-coloring.
- ▶ a $(2, 2, 2)$ -coloring is a 2-improper 3-coloring.

1-improper 2-coloring



(1, 1)-coloring

Known results

Appel and Haken, 1977

- ▶ Every planar graph is $(0, 0, 0, 0)$ -colorable.

Cowen, Cowen, and Woodall, 1986

- ▶ Every planar graph is 2-improperly 3-colorable, i.e. $(2, 2, 2)$ -colorable.

Xu, 2009

Every plane graph with neither adjacent triangles nor 5-cycles is $(1, 1, 1)$ -colorable.

Known results-Choosability

Definition

A graph G is d -improper m -choosable, or simply $(m, d)^*$ -choosable, if for every list assignment L , where $|L(v)| \geq m$ for every $v \in V(G)$, there exists an L -colouring of G such that each vertex of G has at most d neighbours coloured with the same colour as itself.

Eaton and Hull (1999), Škrekovski (1999)

- ▶ Every planar graph is 2-improper 3-choosable: $(3, 2)^*$ -choosable.

If a graph G is 2-improper 3-choosable then it is $(2, 2, 2)$ -colorable.

Škrekovski proved that for every k , there are planar graphs which are not k -improper 2-colorable.

Known results

Cushing and Kierstead (2009)

Every planar graph is 1-improper 4-choosable ($(4, 1)^*$ -choosable).

Dong and Xu (2009)

Let G be a plane graph without any cycles of length in $\{4, 8\}$, then G is $(3, 1)^*$ -choosable.

Question (Xu and Zhang, 2007)

Is it true that every planar graph without adjacent triangle is $(3, 1)^$ -choosable.*

Every planar graph without adjacent triangle is $(1, 1, 1)$ -colorable?

Definition-Maximum average degree

$$\text{Mad}(G) = \max \left\{ \frac{2 \cdot |E(H)|}{|V(H)|}, H \subseteq G \right\}.$$

Definition-Maximum average degree

$$\text{Mad}(G) = \max \left\{ \frac{2 \cdot |E(H)|}{|V(H)|}, H \subseteq G \right\}.$$

In 1995, Jensen and Toft showed that there is a polynomial algorithm to compute $\text{Mad}(G)$ for a given graph G .

T. R. Jensen and B. Toft, Choosability versus chromaticity, Geombinatorics 5(1995), 45-64.

if G is a planar graph with girth g , then $\text{Mad}(G) < \frac{2g}{g-2}$.

Havet and Sereni, 2006

- ▶ For every $k \geq 0$, every graph G with $Mad(G) < \frac{4k+4}{k+2}$ is k -improperly 2-colorable (in fact k -improperly 2-choosable), i.e. (k, k) -colorable
- ▶ $k = 1$ $Mad(G) < \frac{8}{3}$: $(1, 1)$ -colorable (planar, $g = 8$).
- ▶ $k = 2$ $Mad(G) < 3$: $(2, 2)$ -colorable (planar, $g = 6$).

A more general result:

Theorem (Havet and Sereni)

For every $l \geq 2$ and every $k \geq 0$, all graphs of maximum average degree less than $\frac{l(l+2k)}{l+k}$ are k -improper l -choosable.


it implies (k, \dots, k) -colorable.

(d_1, d_2, \dots, d_k) -coloring

(d_1, d_2, \dots, d_k) -coloring

A graph G is (d_1, d_2, \dots, d_k) -colorable if and only if:

- ▶ it exists a partition of V : $V = V_1 \cup V_2 \cup \dots \cup V_k$ such that $\forall i \in [1, k]$, $\Delta(G[V_i]) \leq d_i$



$(2, 1)$ -coloring

In this talk: $(1, 0)$, $(k, 0)$, $(k, 1)$, (k, j) , $(k, 0, \dots, 0)$

$(1, 0)$ -colorable

Theorem (Glebov and Zambalaeva, 2007)

Every planar graph is $(1, 0)$ -colorable if $g(G) \geq 16$.

Theorem (Borodin and Ivanova, 2009)

Every graph is $(1, 0)$ -colorable if $Mad(G) < \frac{7}{3}$


New technique first introduced by Borodin, Ivanova and Kostochka in 2006.

This implies: A planar graph is $(1, 0)$ -colorable if $g(G) \geq 14$

Improvement

Theorem (Borodin and Kostochka, 2010)


Every graph G with $\text{Mad}(G) \leq \frac{12}{5}$ is $(1,0)$ -colorable and the restriction on $\text{Mad}(G)$ is sharp.



Improvement

Theorem (Borodin and Kostochka, 2010)

Every graph G with $\text{Mad}(G) \leq \frac{12}{5}$ is $(1, 0)$ -colorable and the restriction on $\text{Mad}(G)$ is sharp.



$$\text{Mad}(G_p) = \frac{2|E(G_p)|}{|V(G_p)|} = \frac{12p + 6}{5p + 2} = \frac{12}{5} + \frac{6}{5(5p + 2)}$$

$(k, 0)$ -coloring

$(k, 0)$ -coloring

- ▶ Bipartition V_1, V_2 of $V(G)$
- ▶ $\Delta(G[V_1]) \leq k$ and $G[V_2]$ is a stable set.

Color k : vertices of V_1

Color 0 : vertices of V_2 .

$k = 1$


Theorem (Borodin and Kostochka, 2010)

Every graph is $(1, 0)$ -colorable if $\text{Mad}(G) \leq \frac{12}{5}$

Corollary


A planar graph is $(1, 0)$ -colorable if $g(G) \geq 12$

Outerplanar graphs



Smallest value of k such that G admits a $(k, 0)$ -coloring?


Outerplanar graphs



Smallest value of k such that G admits a $(k, 0)$ -coloring?

$k = 1?$


Outerplanar graphs



Smallest value of k such that G admits a $(k, 0)$ -coloring?

$k = 1?$

Outerplanar graphs



(2, 0)-coloring

Outerplanar graphs


Claim

- ▶ Outerplanar graphs with girth 4 are (2,0)-colorable.
- ▶ Outerplanar graphs with girth 5 are (1,0)-colorable.

(the girth of a graph G is the length of a shortest cycle of G .)

Outerplanar graphs


For outerplanar graphs with girth 3, k is unbounded.



Non $(k, 0)$ -colorable outerplanar graph with girth 3

Outerplanar graphs


For outerplanar graphs with girth 3, k is unbounded.



Non $(k, 0)$ -colorable outerplanar graph with girth 3

Outerplanar graphs

For outerplanar graphs with girth 3, k is unbounded.



Non $(k, 0)$ -colorable outerplanar graph with girth 3

$$\text{Mad}(G_{p,k}) = \frac{2|E(G_{p,k})|}{|V(G_{p,k})|} = \frac{2((3k+2)(p+1)-1)}{(2k+2)(p+1)-1} \xrightarrow{p \rightarrow \infty} \frac{3k+2}{k+1}$$

Sparse graphs

Key concepts:

soft components, feeding area

Theorem (Borodin, Ivanova, Montassier, Ochem, R., 2009)

Let $k \geq 0$ be a integer. Every graph with maximum average degree smaller than $\frac{3k+4}{k+2}$ is $(k, 0)$ -colorable.

- ▶ $Mad(G) < \frac{5}{2} \rightarrow (2, 0)$
- ▶ $Mad(G) < \frac{13}{5} \rightarrow (3, 0)$

Sparse graphs

Key concepts:

soft components, feeding area

Theorem (Borodin, Ivanova, Montassier, Ochem, R., 2009)

Let $k \geq 0$ be an integer. Every graph with maximum average degree smaller than $\frac{3k+4}{k+2}$ is $(k, 0)$ -colorable.

- ▶ $Mad(G) < \frac{5}{2} \rightarrow (2, 0)$
- ▶ $Mad(G) < \frac{13}{5} \rightarrow (3, 0)$

Optimality:

$$\lim_{p \rightarrow \infty} Mad(G_{p,k}) = \frac{3k+2}{k+1} < \frac{3k+4}{k+2} + \frac{1}{k+3}.$$

Corollary

Every planar graph G is:


- ▶ $(1, 0)$ -colorable if $g(G) \geq 14$,
- ▶ $(2, 0)$ -colorable if $g(G) \geq 10$,
- ▶ $(3, 0)$ -colorable if $g(G) \geq 9$,
- ▶ $(4, 0)$ -colorable if $g(G) \geq 8$,
- ▶ $(8, 0)$ -colorable if $g(G) \geq 7$.

Corollary

Every planar graph G is:

- ▶ $(1, 0)$ -colorable if $g(G) \geq 14$,
- ▶ $(2, 0)$ -colorable if $g(G) \geq 10$,
- ▶ $(3, 0)$ -colorable if $g(G) \geq 9$,
- ▶ $(4, 0)$ -colorable if $g(G) \geq 8$,
- ▶ $(8, 0)$ -colorable if $g(G) \geq 7$.

For planar graphs with girth 6, k is unbounded.




Non $(k, 0)$ -colorable planar graph with girth 6

$(k, 1)$ -coloring

Theorem (Borodin, Ivanova, Montassier, R., 2010)

Let $k \geq 2$ be a integer. Every graph with maximum average degree smaller than $\frac{10k+22}{3k+9}$ is $(k, 1)$ -colorable.

$(k, 1)$ -coloring



An example of $G_{n,k}$ with $n = 3$ and $k = 3$.

Non $(k, 1)$ -colorable graph.

$$\text{Mad}(G_{n,k}) = \frac{2|E(G_{n,k})|}{|V(G_{n,k})|} = \frac{2(2n - 1 + 5(k - 1)n + n(2k + 3))}{2n - 1 + 3(k - 1)n + n(k + 2)} = \frac{2(7nk - 1)}{n(4k + 1) - 1}$$
$$\lim_{n \rightarrow \infty} \text{Mad}(G_{n,k}) = \frac{14k}{4k + 1}$$

(k, j) -coloring

Let k, j be two integers with $j \geq 2$, $k \geq j + 1$ and $r_1, r_2, r_3, r_4, r_{51}, r_{52}, m_{k,j}$ be seven non-negative reals. Given k, j , let $\mathcal{P}_{k,j}$ be the linear program described below.

Theorem (Borodin, Ivanova, Montassier, R., 2010)

Let $(r_1, r_2, r_3, r_4, r_{51}, r_{52}, m_{k,j})$ be a solution of $\mathcal{P}_{k,j}$, then every graph with $Mad(G) < m_{k,j}$ is (k, j) -colorable.

$\mathcal{P}_{k,j}$:

$$2 + r_1 + r_2 \geq m_{k,j} \quad (1)$$

$$2 + 2 \cdot r_3 \geq m_{k,j} \quad (2)$$

$$1 \geq r_2 \quad (3)$$

$$j + 2 - (j + 1) \cdot r_2 + r_1 \geq m_{k,j} \quad (4)$$

$$j + 2 - j \cdot r_2 + 2 \cdot r_4 \geq m_{k,j} \quad (5)$$

$$r_2 \geq r_{51} \quad (6)$$

$$j + 2 - (j - 1) \cdot r_2 - 2 \cdot r_{51} \geq m_{k,j} \quad (7)$$

$$r_1 \geq r_3 \quad (8)$$

$$r_1 \geq r_4 + r_{52} \quad (9)$$

$$1 \geq r_1 \quad (10)$$

$$k + 2 - (k + 1) \cdot r_1 + r_{51} \geq m_{k,j} \quad (11)$$

$$k + 2 - (k + 1) \cdot r_1 - r_4 + r_{52} \geq m_{k,j} \quad (12)$$

$$r_1 \geq r_{51} \quad (13)$$

$$k + 2 - k \cdot r_1 - r_4 - r_{51} \geq m_{k,j} \quad (14)$$

$$k + 2 - k \cdot r_1 - 2 \cdot r_4 - r_{52} \geq m_{k,j} \quad (15)$$

$$k + j + 2 - (k + j + 1) \cdot r_1 - r_4 - r_{52} \geq m_{k,j} \quad (16)$$

$$k + j + 2 - (k + j + 1) \cdot r_1 - r_{51} \geq m_{k,j} \quad (17)$$

$$\text{Maximize } m_{k,j} \quad (18)$$

(k, j) -coloring

In particular:

Corollary

- ▶ Every graph G with $Mad(G) < \frac{10}{3}$ is $(13, 2)$ -colorable.
- ▶ Every graph G with $Mad(G) < \frac{317}{95}$ is $(7, 3)$ -colorable.

(k, j) -coloring

In particular:

Corollary

- ▶ Every graph G with $Mad(G) < \frac{10}{3}$ is $(13, 2)$ -colorable.
- ▶ Every graph G with $Mad(G) < \frac{317}{95}$ is $(7, 3)$ -colorable.

For planar graphs:

$g(G)$	$(k, 0)$	$(k, 1)$	$(k, 2)$	$(k, 3)$	$(k, 4)$
3,4	×	×	×	×	×
5	×	?	$(13, 2)$	$(7, 3)$	$(4, 4)$ [HS06]
6	×	$(5, 1)$	$(2, 2)$ [HS06]		
7	$(8, 0)$	$(2, 1)$			
8	$(4, 0)$	$(1, 1)$ [HS06]			
9	$(3, 0)$				
10	$(2, 0)$				
14	$(1, 0)$				

(k, j) -coloring: very last result

Let $F(j, k)$ denote the supremum of x such that every graph G with $\text{Mad}(G) < x$ is (k, j) -colorable. It is easy to see that $F(0, 0) = 2$.

Theorem (Borodin and Kostochka, 2011)

Let $j \geq 0$ and $k \geq 2j + 2$. Then $F(j, k) = 2 \left(2 - \frac{k+2}{(j+2)(k+1)} \right)$.

- ▶ If $\text{Mad}(G) < \frac{3k+2}{k+1}$ ($k \geq 2$) then G is $(k, 0)$ -colorable.
- ▶ If $\text{Mad}(G) < \frac{10k+8}{3k+3}$ ($k \geq 4$) then G is $(k, 1)$ -colorable.

$(0, \dots, 0, k)_l$ -coloring

Definition

A $(0, \dots, 0, k)_l$ -coloring of G is a partition (V_1, V_2, \dots, V_l) ($l \geq 3$) of the vertices of G such that for $1 \leq i \leq l - 1$, $G[V_i]$ is a stable set, and $G[V_l]$ has maximum degree at most k .

Theorem (Kaiser, Montassier, R., Rucky, 2011)

Let $l \geq 3$ and $k \geq 0$ be integers. Every graph G with $\text{Mad}(G) < l + \frac{2k}{2k+l+3}$ is $(0, \dots, 0, k)_l$ -colorable.

If $\text{Mad}(G) < \frac{13}{4}$ then G is $(0, 0, 1)$ -colorable.

Steinberg Conjecture

Conjecture (Steinberg, 1976)

Every planar graph without 4 and 5-cycles is 3-colorable ((0, 0, 0)-colorable)

Can we prove :

- ▶ (0, 0, 1)-colorable?

Steinberg Conjecture

Conjecture (Steinberg, 1976)

Every planar graph without 4 and 5-cycles is 3-colorable ((0, 0, 0)-colorable)

Can we prove :

- ▶ (0, 0, 1)-colorable?
- ▶ (0, 1, 1)-colorable?

Steinberg Conjecture

Conjecture (Steinberg, 1976)

Every planar graph without 4 and 5-cycles is 3-colorable ((0, 0, 0)-colorable)

Can we prove :

- ▶ (0, 0, 1)-colorable?
- ▶ (0, 1, 1)-colorable?
- ▶ (1, 1, 1)-colorable (Xu, 2009)

Steinberg Conjecture

Conjecture (Steinberg, 1976)

Every planar graph without 4 and 5-cycles is 3-colorable ((0, 0, 0)-colorable)

Can we prove :

- ▶ (0, 0, 1)-colorable?
- ▶ (0, 1, 1)-colorable?
- ▶ (1, 1, 1)-colorable (Xu, 2009)
- ▶ (0, 0, 2)-colorable?

Steinberg Conjecture

Conjecture (Steinberg, 1976)

Every planar graph without 4 and 5-cycles is 3-colorable ((0, 0, 0)-colorable)

Can we prove :

- ▶ (0, 0, 1)-colorable?
- ▶ (0, 1, 1)-colorable?
- ▶ (1, 1, 1)-colorable (Xu, 2009)
- ▶ (0, 0, 2)-colorable?

(0, 0, 7) seems ok.....

Thank you for your attention!