

ACYCLIC k -STRONG HYPERGRAPH COLORING OF MAPS ON SURFACES

O.V. Borodin¹, A.V. Kostochka²

Institute of Mathematics, 630090, Novosibirsk, Russia

A. Raspaud and E. Sopena³

LaBRI, Université Bordeaux I, 33405 Talence Cedex, France

Abstract. A vertex coloring of a graph is acyclic if the ends of each edge have different colors and each bicolored subgraph is acyclic. Let each face of size at most k , where $k \geq 3$, in a map on a surface S^N be replaced by a clique on the same vertices. Then the resulting map can be acyclically colored with a number of colors that depends linearly both on N and on k . Such results were previously known only for $1 \leq N \leq 2$ and $3 \leq k \leq 4$.

Keywords. Acyclic coloring, Acyclic strong coloring.

1 Introduction

The problems of coloring graphs embedded on surfaces is an important part of graph theory. In particular, the well-known Four Color Problem belongs to this area.

We denote by $V(G)$ the set of vertices of a graph G and by $E(G)$ its set of edges. A (proper) k -coloring of G is a mapping $f : V(G) \rightarrow \{1, 2, \dots, k\}$ such that $f(x) \neq f(y)$ whenever x and y are adjacent in G .

A vertex coloring of a graph is called k -cyclic or k -strong hypergraph coloring if every two distinct vertices lying in the boundary of the same face of size at most k have different colors. (We use below both terms.) This type of coloring may be looked at as an admissible coloring of a pseudograph obtained by replacing each face of size at most k by a clique on the same vertices. Let $\chi_k(G)$ be the minimum number of colors sufficient to color G k -cyclically, and let $\chi_k(S^N)$ be the minimum number of colors sufficient to color k -cyclically each map on surface S^N with Euler characteristic N . (Sometimes we drop the arguments.)

The case $k = 3$ corresponds to the ordinary vertex colouring, and the Four Color Theorem by Appel and Haken [1] and Heawood's theorem [2] give precise upper bounds for the plane and for the higher surfaces, respectively.

The case $k = 4$ may also be formulated in terms of the simultaneous vertex-face coloring and the vertex coloring of 1-embeddable graphs. For the plane, Borodin [3] proved, confirming Ringel's conjecture in [4], precise upper bound $\chi_4 \leq 6$; for the projective plane, Schumacher [5] proved precise upper bound $\chi_4 \leq 7$; and for the other surfaces, Ringel [6] proved that $\chi_4(S^N) \leq \frac{2H(N)}{\sqrt{3}}$, where $H(N) = \lfloor \frac{7 + \sqrt{49 - 24N}}{2} \rfloor$ is the Heawood number.

¹A part of this research was done during a visit of LaBRI, supported by the NATO Collaborative Research Grant n° 97-1519. It was partially supported by the grant 97-01-01075 of the Russian Foundation for Fundamental Research.

²This work was partially supported by the INTAS grant 97-1001 and by the grant 96-01-01614 of the Russian Foundation for Fundamental Research.

³The work of these two authors was partly supported by the NATO Collaborative Research Grant n° 97-1519.

For arbitrary $k \geq 3$, Ore and Plummer [7] proved that each plane graph G is k -cyclically $2k$ -colorable, and Borodin, Sanders and Zhao [8] improved this bound to $\chi_k(G) \leq \frac{9k}{5}$.

A vertex coloring of a graph is *acyclic* if it is admissible, i.e., the ends of each edge are colored differently, and there is no bicolored cycle. Note that a loop gives a unicolored edge and two multiple edges give a bicolored cycle. We define a vertex coloring of a map to be acyclic provided that no two ends of an edge e are coloured the same unless e is a loop, and no bicolored cycle of length > 2 exists. Borodin [9] proved that each plane graph is acyclically 5-colorable, which bound is best possible. Albertson and Berman [10] proved that each graph embedded on the oriented surface S^N , where $N < 0$, is acyclically $(8 - 2N)$ -colorable. Alon, Mohar and Sanders [11] proved, using the acyclic 5-colorability of plane graphs, that each graph on the projective plane is acyclically 7-colorable, and showed this bound to be best possible. Also, they prove that each graph embedded on an arbitrary surface S^N is acyclically $\mathcal{O}(N^{4/7})$ -colorable, and this bound is tight up to the factor of $(\log N)^{1/7}$.

The acyclic coloring has a number of applications to other coloring problems, for the definitions see [12, 13, 14, 15, 16]. Suppose one has proved that $a(G) \leq a$. Then G has the star chromatic number at most $a2^{a-1}$ (Grünbaum [13]) and the oriented chromatic number at most $a2^{a-1}$ (Raspaud and Sopena [16]); every m -coloring of the edges of G can be homomorphically mapped on that of a graph with at most am^{a-1} vertices (Alon and Marshall [12]); every m -coloring of the edges of mixed graph G combined with its n -coloring can be homomorphically mapped on a graph with at most $a(2n + m)^{a-1}$ vertices (Nešetřil and Raspaud [15]).

Also, Hakimi, Mitchem and Schmeichel [14, p.38-39] proved that $E(G)$ can be partitioned into $a(G)$ star forests (whose every component is a star). Directly using [9], this confirms the conjecture of Algor and Alon [17] that the edges of every planar graph can be partitioned into five star forests.

In this paper we consider a coloring that is both acyclic and k -cyclic. Namely, we color acyclically the result of replacing all faces of size at most k in a map by cliques of the same sizes. So, each face of size at most k is assumed to have all ‘invisible’ diagonals. For $k = 3$ this type of coloring coincides with the acyclic coloring.

Our main result is

Theorem 1 *Every map on a surface S^N is acyclically k -strong hypergraph colorable with $c_N k + d_N$ colors whenever $k \geq 4$ and $N \leq 0$.*

In fact, we prove the result with $c_N = \max\{999, 117 - 471N\}$ and $d_N = 39 - 156N$. The argument is intentionally kept simple; more complicated argument may be used to decrease c_N and d_N .

Corollary 2 *Every map on the plane ($N = 2$) or on the projective plane ($N = 1$) is acyclically k -strong hypergraph colorable with $c_0 k + d_0$ colors whenever $k \geq 4$.*

Proof. Follows from the facts that each plane or projective plane map is also a map on the torus or Klein bottle, respectively. \square

In [18], we prove that each projective plane graph (and hence each plane graph) is acyclically 4-strong hypergraph 20-colorable, i.e., each graph 1-embedded on the projective plane is acyclically 20-colorable. Thus, Theorem 1 and Corollary 2 augment the results of [9] ($N = 2, k = 3$), of [11] ($N \leq 1, k = 3$), and of [18] ($1 \leq N \leq 2, k = 4$).

2 Proof of Theorem 1

The *size*, $s(f)$, of a face f is the number of edges in its boundary, $\partial(f)$, counting multiplicities. For instance, a cut edge appears twice on the boundary of a certain face. To simplify the argument, we restrict ourselves to the case when $\partial(f)$ is connected whenever $f \in F$. The *degree* of a vertex v , i.e., the number of incident edges (loops are counted twice), is denoted by $d(v)$. By a $\geq k$ -vertex we mean that of degree at least k , etc.

Given S^N and k , let P''' be a counterexample with the minimum number of vertices. For brevity, the acyclically k -strong hypergraph coloring with $c_N k + d_N$ colors we are searching for will be called *good*.

Delete from P''' the common edge of two adjacent 3-faces, if any. Repeat this operation until the resulting counterexample, P'' , has no adjacent 3-faces.

Triangulate all $> k$ -faces of P'' by adding diagonals to obtain another counterexample, P' , with the minimum number of vertices. (A good coloring of P' would be good for P'' .)

Remove from P' each loop e that forms a 1-face. This decreases the size of the other face incident with e by 1. Similarly, for a 2-face $f = e_1 e_2$ in P' , remove one of the boundary edges, say e_1 . This results in a face of the same size as the former face that was incident with e_1 and different from f .

If the so obtained graph P were well colorable, then recovering loops and repeated edges would not spoil the good coloring. Thus, P is also a counterexample with the fewest vertices. We have actually proved the following

Lemma 3 *If f is a face of P , then $3 \leq s(f) \leq k$; no two 3-faces of P are adjacent.*

Informally speaking, only adding new nontrivial adjacencies, represented by ‘visible’ edges or ‘invisible diagonals’, can spoil the good coloring. Our concern therefore is not to lose adjacencies while transforming P into a smaller pseudograph, which already admits a good coloring.

The following two observations should be kept in mind during the rest of the proof.

Observation 4 *Contracting an edge $e = vz$ into a vertex $v * z$ decreases the size of each face f in P by 0, 1 or 2, depending on how many times e appears on $\partial(f)$.*

*The obtained map R is well colorable by the minimality of P . Transfer its coloring to P with z colored as $v * z$ and v uncolored. Then in order that to get a good coloring for P , it remains to find a color for v without creating unicolored edges other than loops and bicolored cycles other than 2-cycles.*

Observation 5 *Suppose a vertex v is incident with edges $e_i = vz_i$, where $0 \leq i \leq d(v) \leq k - 1$, in a cyclic order. (Of course, z_i 's are not all necessarily distinct.) Split each nontriangular face $f_i = \dots z_i v z_{i+1}$, where subscripts are taken modulo $d(v)$, into a triangle $z_i v z_{i+1}$ and a face f'_i . Then $s(f'_i) < s(f_i)$. Remove v and all e_i 's, then a new face arises of size $d(v) \leq k$, so that the obtained pseudograph can be well colored. Hence, all distinct z_i 's are colored differently. If we choose a color α for v not appearing on any vertex in $\cup_{0 \leq i \leq k-1} \partial(f_i)$, then the only possibility for the obtained coloring to be bad is a bicolored α, β -cycle (i.e., consisting of vertices alternatively colored with α and β) through v and some vertex u such that u is colored with β and $u \in \partial(f_i) \setminus \{z_i, z_{i+1}\}$ for some f_i .*

Lemma 6 *If $v \in V(P)$ then $d(v) \geq 2$.*

Proof. If $d(v) = 1$, we contract the edge vz , transfer the good coloring of the obtained pseudograph to P , and color v differently from those at most k vertices that belong to the same faces as v . \square

Lemma 7 *Each face in P is incident with at least three vertices (not necessarily distinct) of degree at least three.*

Proof. If a face f in P has only two vertices u, w (not necessarily distinct) of degree greater than 2 in $\partial(f)$, then we remove the longer of the two paths that form $\partial(f)$. This creates a new face f' of size $\leq k$. From a good coloring of the resulting graph, it is easy to obtain a good coloring of P , since all vertices on $\partial(f')$ are colored pairwise differently.

The case that there is only one > 2 -vertex in $\partial(f)$ easily reduces to the previous one. If all vertices in $\partial(f)$ have degree 2, then P is a cycle; a contradiction. \square

If every face of P is an open 2-cell, then Euler's formula says that $|V| - |E| + |F| = N$; otherwise

$$|V| - |E| + |F| \geq N.$$

Using obvious equalities $2|E| = \sum_{v \in V} d(v) = \sum_{f \in F} s(f)$, this may be rewritten as

$$\sum_{v \in V} (d(v) - 4) + \sum_{f \in F} (s(f) - 4) \leq -4N,$$

or

$$-2n_2 + \sum_{v \in V_{3+}} (d(v) - 4) + \sum_{f \in F} (s(f) - 4) < -4N + 1, \quad (1)$$

where n_i is the number of i -vertices in P , and V_{i+} is the set of $\geq i$ -vertices in P .

Denote by $n_2(f)$ the number of 2-vertices on the boundary of face f , counting multiplicities. The *reduced size*, $s^*(f)$, of a face f is defined to be $s(f) - n_2(f)$. In fact, $s^*(f)$ is the number of > 2 -vertices in $\partial(f)$, counting multiplicities. Then (1) may be put as

$$\sum_{v \in V_{3+}} (d(v) - 4) + \sum_{f \in F} (s^*(f) - 4) < -4N + 1. \quad (2)$$

By Lemma 7, $s^*(f) \geq 3$ whenever $f \in F$. Let $f_i^*(v)$ be the number of i^* -faces at v , i.e., those having $s^*(f) = i$. Then

$$\sum_{v \in V_{3+}} (d(v) - 4 - \frac{f_3^*(v)}{3}) + \sum_{f \in F_{4+}^*} (s^*(f) - 4) < -4N + 1, \quad (3)$$

where F_{i+}^* is the set of $\geq i^*$ -faces in P , or

$$\sum_{v \in V_{3+}} (d(v) - 4 - \frac{f_3^*(v)}{3} + \frac{4N - 1}{n^*}) + \sum_{f \in F_{4+}^*} (s^*(f) - 4) < 0, \quad (4)$$

where $n^* = |V_{3+}| = |V| - n_2$.

Put $ch(v) = d(v) - 4 - \frac{f_3^*(v)}{3} + \frac{4N-1}{n^*}$ whenever $v \in V_{3+}$, $ch(f) = 0$ if f is a 3^* -face, and $ch(f) = s^*(f) - 4$ whenever $f \in F_{4+}^*$. Then (4) becomes

$$\sum_{v \in V_{3+}} ch(v) + \sum_{f \in F} ch(f) < 0. \quad (5)$$

The rest of our proof consists in redistributing *the charge* $ch(x)$ on $x \in V_{3+} \cup F$, preserving the sum of charges, so that *the new charge*, $ch^*(x)$, is nonnegative whenever $x \in V_{3+} \cup F$. The contradiction with (5) will complete the proof.

First observe the following fact.

Lemma 8 $n^* > 39(-4N + 1)$.

Proof. Euler's formula (1) for the map P^* obtained from P by contracting each path $uv_1 \dots v_s w$ such that $d(u) \geq 3$, $d(v_1) = \dots = d(v_s) = 2$, $d(w) \geq 3$ into an edge uw , may be written as

$$\sum_{v \in V(P^*)} (d_{P^*}(v) - 6) + \sum_{f \in F(P^*)} (2s_{P^*}(f) - 6) \leq -6N,$$

which implies

$$\sum_{v \in V(P^*)} (d_{P^*}(v) - 6) \leq -6N,$$

or else

$$|E(P^*)| \leq 3n^* - 3N.$$

By Lemma 1,

$$n \leq n^* + k|E(P^*)|.$$

But $n > c_N k + d_N$, for otherwise P has a trivial good coloring (all vertices have different colors), and so

$$n^*(1 + 3k) - 3kN > c_N k + d_N \geq (117 - 471N)k + 39 - 156N = 39(-4N + 1)(1 + 3k) - 3kN,$$

as desired (compare the leftmost and the rightmost expressions). \square

Lemma 9 *If $d(v) \geq 7$ then $ch(v) \geq 0$.*

Proof. Indeed,

$$ch(v) \geq d(v) - 4 - \frac{f_3^*(v)}{3} + \frac{4N - 1}{n^*} \geq d(v) - 4 - \frac{d(v)}{3} + \frac{4N - 1}{n^*} \geq \frac{2}{3} + \frac{4N - 1}{n^*},$$

and we are done by Lemma 8. \square

If $v \in V$ and $ch(v) < 0$, then v is called *poor*. Put

$$\xi = \frac{-4N + 1}{n^*}$$

and

$$\varepsilon = \frac{1}{39}.$$

By Lemmas 4 and 5, $\xi \leq \varepsilon$, and each poor vertex v satisfies $3 \leq d(v) \leq 6$.

Our rules of redistribution of charge are:

R1. Each ≥ 14 -vertex gives $\frac{2}{3} + \varepsilon$ to each incident $\geq 4^*$ -face and $\frac{1}{3} + \varepsilon$ to each incident 3^* -face.

R2. Each poor vertex v receives $\frac{2}{3} + \varepsilon$

- from each incident $\geq 4^*$ -face f such that there is a ≥ 14 -vertex z in $\partial(f)$ not joined to v along $\partial(f)$ by a path of 2-vertices, and
- from each incident $\geq 13^*$ -face f .

R3. If a 3^* -face f is incident with a ≥ 14 -vertex, then f gives $\frac{1}{6} + \frac{\varepsilon}{2}$ to every incident poor vertex.

Lemma 10 *If v is poor, then v receives $\frac{1}{6} + \frac{\varepsilon}{2}$ by R3 from each of every two consecutive 3^* -faces at v .*

Proof. We first prove that if a 3^* -face f is incident with a 2-vertex u , then the ≥ 3 -vertex which is opposite to u in $\partial(f)$ is in fact a ≥ 14 -vertex.

Let $\partial(f) = xx_1x_2 \dots x_{k(x)}yy_1y_2 \dots y_{k(y)}zz_1z_2 \dots z_{k(z)}$, where all x_i 's, y_i 's, and z_i 's are all 2-vertices, while x , y , and z are ≥ 3 -vertices. Suppose that $k(x) \geq 1$, i.e., x_1 exists. Denote by f_{xy} the face lying on the other side of path $xx_1x_2 \dots x_{k(x)}xy$ than f (perhaps $f_{xy} = f$). The faces f_{yz} and f_{zx} are defined similarly.

Contract the edge xx_1 transfer a good coloring of the obtained map to $P - x_1$. If we color x_1 with a color that does not appear on the boundaries of f and the faces incident with z , then the obtained coloring is good. This is impossible only if $d(z) \geq 14$.

To complete the proof of our Lemma, observe that by the second statement of Lemma 3, at least one of every two consecutive 3^* -faces, f_1 and f_2 , at v is incident with a 2-vertex, z . If z is incident both

with f_1 and with f_2 , then we are done by R3 and the statement just proved. Otherwise, z is opposite to the ≥ 3 -vertex, w , which is incident both with f_1 and with f_2 , because by the above statement, z cannot be opposite to the poor vertex v which has degree at most six due to Lemma 5. It follows that $d(w) \geq 14$, and we again use R3. \square

Lemma 11 *If $d(v) = 3$ then v is incident with at least two faces each of which gives $\frac{2}{3} + \varepsilon$ to v by R2.*

Proof. Let v be incident with paths $vx_1x_2 \dots x_{k(x)}x$, $vy_1y_2 \dots y_{k(y)}y$, and $vz_1z_2 \dots z_{k(z)}z$, where all x_i 's, y_i , and z_i 's are 2-vertices while x , y , and z are all > 2 -vertices. Let $\partial(f_1) = \dots xx_{k(x)} \dots x_2x_1vy_1y_2 \dots y_{k(y)}y$, $\partial(f_2) = \dots yy_{k(y)} \dots y_2y_1vz_1z_2 \dots z_{k(z)}z$, and $\partial(f_3) = \dots zz_{k(z)} \dots z_2z_1vx_1x_2 \dots x_{k(x)}x$.

Suppose the contrary, namely, by symmetry, that neither f_1 nor f_2 gives $\frac{2}{3} + \varepsilon$ to v . Then by R2 each of f_1 and f_2 is a $\leq 12^*$ -face, and each vertex in $\partial(f_1) \setminus \{x, y\}$ and in $\partial(f_2) \setminus \{y, z\}$ has degree at most 13.

Contract the path $vy_1y_2 \dots y_{k(y)}y$ to a vertex $v*y$. Transfer a good coloring of the obtained map to P . Color $y_1, y_2, \dots, y_{k(y)}$ and v pairwise differently using colors that do not appear on the boundaries of the faces f_1, f_2, f_3 , and those at most $2 \times 9 \times (13 - 1)$ faces that are incident with > 2 -vertices in $\partial(f_1) \setminus \{x, y\} \cup \partial(f_2) \setminus \{y, z\}$. The number of restrictions is less than $3k + 2 \times 9 \times (13 - 1)k < c_N k + d_N$, and no nontrivial unicolored edges or bicolored cycles appear. \square

Lemma 12 *If $d(v) = 4$ then v is incident with at least one face giving $\frac{2}{3} + \varepsilon$ to v by R2.*

Proof. Let v be incident with paths $vx_1^i x_2^i \dots x_{k(x^i)}^i x^i$ in a cyclic order, where $0 \leq i \leq 3$, all x_j^i 's are 2-vertices, and all x^i 's are > 2 -vertices.

Let the face f^i have

$$\partial(f^i) = x^i x_{k(x^i)}^i \dots x_2^i x_1^i v x_1^{i+1} x_2^{i+1} \dots x_{k(x^{i+1})}^{i+1} x^{i+1} \dots,$$

where upper indices are taken modulo 4.

Suppose that none of f^i gives $\frac{2}{3} + \varepsilon$ to v . Then by R2 each f^i is a $\leq 12^*$ -face, and each vertex in $\partial(f^i) \setminus \{x^i, x^{i+1}\}$ has degree at most 13.

Add the edge $x^i x^{i+1}$ into f^i whenever $0 \leq i \leq 3$, unless such an edge already exists in $\partial(f^i)$. Delete v and transfer a good coloring of the obtained map to P . Color v using a color that does not appear on the boundaries of the faces incident with vertices in $\partial(f^i) \setminus \{x^i, x^{i+1}\}$ whenever $0 \leq i \leq 3$. The number of restrictions is less than $4 \times 9 \times 13k \leq c_N k + d_N$, and it is not hard to see that no nontrivial unicolored edges or bicolored cycles appear. \square

Lemma 13 *If $v \in V_{3+}$, then $ch^*(v) \geq 0$.*

Proof. First suppose v is poor, i.e., has $ch(v) < 0$. Then, by Lemma 9, $d(v) < 7$. If $d(v) = 3$ then $ch(v) = -1 - \xi$, and we are done due to Lemma 11, because $2 \times (\frac{2}{3} + \varepsilon) > 1 + \xi$.

If $d(v) = 4$ then $ch(v) = -\xi$. By Lemma 12, v gets at least $\frac{2}{3} + \varepsilon$ from an incident $\geq 4^*$ -face. If v is incident with at most two 3^* -faces, then $ch^*(v) \geq \varepsilon - \xi \geq 0$. Otherwise, by Lemma 10, v gets at least $2(\frac{1}{6} + \frac{\varepsilon}{2})$ from the three incident 3^* -faces, and $ch^*(v) \geq 0$.

If $d(v) = 5$ then $ch(v) = 1 - \xi$, and we are done if v is incident with at most two 3^* -faces. If there are r such faces at v , where $3 \leq r \leq 5$, then Lemma 10 provides v with at least $2(\frac{1}{6} + \frac{\varepsilon}{2})$ if $r = 3$ and at least $4(\frac{1}{6} + \frac{\varepsilon}{2})$ if $r \geq 4$, whence $ch^*(v) \geq 0$.

If $d(v) = 6$ then $ch(v) = 2 - \xi$, so that we are done if v is incident with at most five 3^* -faces. Otherwise, by Lemma 10, v gets at least $3 \times (\frac{1}{6} + \frac{\varepsilon}{2})$ from the six incident 3^* -faces, and $ch^*(v) \geq 0$.

Now let v not be poor. If it does not give anything to neighbor vertices by R1 or R2, then $ch^*(v) = ch(v) \geq 0$. Otherwise, $d(v) \geq 14$. Then v makes at most $d(v)$ transfers by R1. It follows that

$$ch^*(v) = d(v) - 4 - \xi - d(v)\left(\frac{2}{3} + \varepsilon\right),$$

whence $ch^*(v) \geq 0$, because $\xi \leq \varepsilon$. □

Lemma 14 *If $f \in F$, then $ch^*(f) \geq 0$.*

Proof. A 3*-face does not participate in discharging if all its incident vertices have degree at most 13. Otherwise, it receives at least $\frac{2}{3} + \varepsilon$ by R1 and gives at most $2\left(\frac{1}{3} + \frac{\varepsilon}{2}\right)$ to the incident poor vertices. Hence $ch^*(f) \geq 0$ in both cases.

Now suppose $f \in F_{4+}^*$. If there are at least two vertices of degree ≥ 14 incident with f , then

$$ch^*(f) \geq s^*(f) - 4 + 2\left(\frac{2}{3} + \varepsilon\right) - (s^*(f) - 2)\left(\frac{2}{3} + \varepsilon\right) = (s^*(f) - 4)\left(\frac{1}{3} - \varepsilon\right) \geq 0.$$

If there is only one ≥ 14 -vertex, z , at f , then

$$ch^*(f) \geq s^*(f) - 4 + \frac{2}{3} + \varepsilon - (s^*(f) - 3)\left(\frac{2}{3} + \varepsilon\right) \geq 0,$$

because f gives $\frac{2}{3} + \varepsilon$ to at most $s^*(f) - 3$ poor vertices by R2: nothing is given to z itself and to its left and right neighbours of degree ≥ 3 along $\partial(f)$.

Suppose there is no vertex of degree ≥ 14 at f . If $s^*(f) \leq 12$, then $ch^*(f) = ch(f) = s^*(f) - 4 \geq 0$, because f does not participate in discharging. Finally, if $s^*(f) \geq 13$ then by R2,

$$ch^*(f) \geq s^*(f) - 4 - s^*(f)\left(\frac{2}{3} + \varepsilon\right) = s^*(f)\left(\frac{1}{3} - \varepsilon\right) - 4 \geq 0,$$

because $\varepsilon = \frac{1}{39}$. □

The lemmas above imply together that $ch^*(x) \geq 0$ for every $x \in V_{3+} \cup F$. This contradiction with (5) completes the proof.

References

- [1] K.Appel, W.Haken, The solution of the four-color-map problem. *Scientific American*, 237, No.4, (1977), 108-121.
- [2] P.J.Heawood, Map-color theorem, *Quart. J. Math.*, 24, (1890), 332-338.
- [3] O.V.Borodin, Solution of Ringel's problems on the vertex-face coloring of plane graphs and on the coloring of 1-planar graphs. *Diskret. Analiz, Novosibirsk*, 41 (1984) 12-26 (in Russian).
- [4] G.Ringel, Ein Sechsfarbenproblem auf der Kugel, *Abh. Math. Sem. Univ. Hamburg*. 1965. V. 29. P. 107-117.
- [5] H.Schumacher, Ein 7-Farbensatz 1-einbettbarer Graphen auf der projektiven Ebene, *Abh. Math. Sem. Univ. Hamburg*, 54 (1984), 5-14.
- [6] G.Ringel, A nine color theorem for the torus and the Klein bottle, *The Theory and Applications of Graphs (Kalamazoo, Mich., 1980)*. New York: Willey, (1981), 507-515.
- [7] O.Ore, M.D.Plummer, Cyclic coloration of plane graphs, *Recent Progress in Combinatorics*. New York: Academic Press, (1969), 287-293.

- [8] O.V.Borodin, D.Sanders and Y.Zhao, On cyclic colorings and their generalizations, *Discrete Math.* (accepted).
- [9] O.V.Borodin, On acyclic colorings of planar graphs. *Discrete Math.* 25, (1979) 198-223.
- [10] M.Albertson and D.Berman, An acyclic analogue to Heawood's theorem. *Glasgow Math. J.*, 19, (1977) 169-174.
- [11] N.Alon, B.Mohar, D.P.Sanders, On acyclic colorings of graphs on surfaces. *Isr. J. Math.* 94 (1996), 273-283.
- [12] N.Alon, T.H.Marshall, Homomorphisms of edge-colored graphs and Coxeter groups *J. Algebraic Combinatorics* 2 (1998), 277-289.
- [13] B.Grünbaum, Acyclic colorings of planar graphs, *Israel J. Math.* 1973. V. 14, N. 3. P. 390-408.
- [14] T. R. Jensen and B. Toft, *Graph coloring problems*, Wiley Interscience (1995).
- [15] J. Nešetřil, A. Raspaud, Colored Homomorphisms of Colored Mixed Graphs, *KAM Series NO. 98-376*, Dept. of Applied Math. Charles University, Prague (Czech Republic); to appear in *J. Combinatorial Theory, Ser. B.*
- [16] A. Raspaud and E. Sopena, Good and semi-strong colorings of oriented planar graphs, *Inform. Processing Letters* 51 (1994), 171-174.
- [17] I.Algor and N.Alon, The star arboricity of graphs *Discrete Math.* 75 (1989), 11-22.
- [18] O.V.Borodin, A.V.Kostochka, A.Raspaud, E.Sopena, On acyclic 20-colorability of 1-planar graphs (in preparation).