

# THERE EXIST ORIENTED PLANAR GRAPHS WITH ORIENTED CHROMATIC NUMBER AT LEAST SIXTEEN

Éric SOPENA

LaBRI, Université Bordeaux 1, 351, cours de la Libération, 33405 Talence, France

**Abstract.** We prove that there exist oriented planar graphs with oriented chromatic number at least 16. Using a result of Raspaud and Sopena [*Good and semi-strong colorings of oriented planar graphs, Inf. Processing Letters 51, 171–174, 1994*], this gives that the oriented chromatic number of the family of oriented planar graphs lies between 16 and 80.

**Keywords.** combinatorial problems, oriented graph colouring, graph homomorphism.

## 1 Introduction

We consider oriented graphs, that is digraphs having no loops and no opposite arcs. If  $G$  is an oriented graph, we denote by  $V(G)$  its set of vertices and by  $E(G)$  its set of arcs. If  $xy$  is an arc in  $E(G)$ , we say that  $y$  is a *successor* of  $x$  and that  $x$  is a *predecessor* of  $y$ .

An *oriented  $k$ -colouring* of an oriented graph  $G$  is a mapping  $c : V(G) \rightarrow \{1, 2, \dots, k\}$  such that (i) if  $xy \in E(G)$  then  $c(x) \neq c(y)$  and (ii) if  $xy, zt \in E(G)$  then  $c(x) = c(t) \implies c(y) \neq c(z)$ . With every oriented  $k$ -colouring  $c$  of  $G$  one can associate a digraph  $H_c$ , called the *colour-graph* of  $c$ , with vertex set  $V(H_c) = \{c(x), x \in V(G)\}$  and arc set  $E(H_c) = \{c(x)c(y), xy \in E(G)\}$ . Thanks to conditions (i) and (ii),  $H_c$  is an oriented graph. The oriented  $k$ -colouring  $c$  can then be viewed as a homomorphism (that is an arc-preserving vertex mapping) from  $G$  to  $H_c$ . Similarly, every homomorphism of  $G$  to an oriented graph  $H$  on  $k$  vertices can be viewed as an oriented  $k$ -colouring of  $G$ , using the vertices of  $H$  as colours. Oriented colourings have been introduced in [5] and studied by several authors (see [7] for a general overview). In particular, various problems related to planar graphs are discussed in [1], [2] and [4].

The *oriented chromatic number*  $\vec{\chi}(G)$  of an oriented graph  $G$  is defined as the smallest  $k$  such that  $G$  admits an oriented  $k$ -colouring or, equivalently, as the smallest order of an oriented graph  $H$  such that  $G$  admits a homomorphism to  $H$ . The oriented chromatic number  $\vec{\chi}(\mathcal{F})$  of a family  $\mathcal{F}$  of oriented graphs is then defined as the largest oriented chromatic number among the oriented chromatic numbers of its members ( $\vec{\chi}(\mathcal{F})$  can thus be infinite). Let us denote by  $\mathcal{P}$  the family of oriented planar graphs. It has been proved in [6] that  $\vec{\chi}(\mathcal{P}) \leq 80$ .

From the definition of oriented  $k$ -colourings, we get that if  $xyz$  is a directed 2-path in  $G$  ( $xy, yz \in E(G)$ ) then  $c(x) \neq c(y) \neq c(z) \neq c(x)$  for every oriented  $k$ -colouring  $c$  of  $G$ . In other words, any two vertices that are linked in  $G$  by a directed path of length 1 or 2 must be assigned distinct colours. Let  $G_1$  be the oriented graph having one vertex and no arc and  $G_i$ ,  $i \geq 2$ , be the oriented graph obtained by taking two disjoint copies of  $G_{i-1}$  and a new vertex  $x_i$  and adding all the arcs from the vertices of the first copy towards  $x_i$  and all the arcs from  $x_i$  towards the vertices of the second copy (see Figure 1). By construction, and considering the observation above, we get  $\vec{\chi}(G_i) = 2^i - 1$  for every  $i \geq 1$ . Note in particular that the graph  $G_3$  is outerplanar and that the graph  $G_4$  is thus planar, so that  $\vec{\chi}(\mathcal{P}) \geq 15$ . In section 2, we shall prove the following improvement of this lower bound:




Figure 1: A sequence of oriented graphs  $G_i$ ,  $i \geq 1$ , with  $\vec{\chi}(G_i) = 2^i - 1$

**Theorem 1**  $\vec{\chi}(\mathcal{P}) \geq 16$ .

Let us say that an oriented graph  $U$  is  $\mathcal{F}$ -universal for some family  $\mathcal{F}$  of oriented graphs if every oriented graph  $F \in \mathcal{F}$  admits a homomorphism to  $U$ . If the family  $\mathcal{F}$  has oriented chromatic number at most  $k$ , a  $\mathcal{F}$ -universal oriented graph on  $k$  vertices does not necessarily exist. (Consider for instance the family  $\mathcal{G}_k$  of all oriented graphs on  $k$  vertices; the oriented chromatic number of  $\mathcal{G}_k$  is clearly at most  $k$  but there exists no  $\mathcal{G}_k$ -universal oriented graph on  $k$  vertices since such a graph would contain all tournaments on  $k$  vertices as subgraphs). However, if we consider the family  $\mathcal{P}$  of oriented planar graphs such a universal graph exists:

**Proposition 2** For every  $k > 0$ , if  $\vec{\chi}(\mathcal{P}) \leq k$  then there exists a  $\mathcal{P}$ -universal graph on  $k$  vertices.

**Proof.** Suppose to the contrary that for every oriented graph  $U$  on  $k$  vertices there exists some oriented planar graph  $P_U$  which does not map homomorphically to  $U$ . Denote by  $\ell$  the number of oriented graphs on  $k$  vertices and let  $\mathcal{G}_k = \{U_1, U_2, \dots, U_\ell\}$ . Consider the oriented graph  $P$  obtained by taking one copy of each graph  $P_{U_1}, P_{U_2}, \dots, P_{U_\ell}$  and gluing them by identifying one vertex of each of them into a unique vertex. The graph  $P$  thus obtained is clearly planar and therefore there exists some oriented graph  $U_i$  in  $\mathcal{G}_k$  such that  $P \rightarrow U_i$ . Considering the restriction to  $P_{U_i}$  of such a homomorphism, we get a homomorphism from  $P_{U_i}$  to  $U_i$  which contradicts our assumption.  $\square$

## 2 Minimal $\mathcal{P}$ -universal graphs

In this section we prove Theorem 1 using a series of Lemmas. For every oriented graph  $G$  and every vertex  $x \in V(G)$  we denote by  $\Gamma_G^+(x)$  the subgraph of  $G$  induced by the successors of  $x$  and by  $\Gamma_G^-(x)$  the subgraph of  $G$  induced by the predecessors of  $x$ . For any two oriented graphs  $G_1$  and  $G_2$ , we denote by  $G_1 \cap G_2$  the graph given by  $V(G_1 \cap G_2) = V(G_1) \cap V(G_2)$  and  $E(G_1 \cap G_2) = E(G_1) \cap E(G_2)$ .

Let  $U$  be a minimal  $\mathcal{P}$ -universal oriented graph (by minimal, we mean here that no proper subgraph of  $U$  is  $\mathcal{P}$ -universal).

**Lemma 3** There exists an oriented planar graph  $P^*$  such that for every homomorphism  $\varphi: P^* \rightarrow U$  and every arc  $uv \in E(U)$  there exists an arc  $xy \in E(P^*)$  such that  $\varphi(x) = u$  and  $\varphi(y) = v$ .

**Proof.** Let  $uv$  be any arc in  $E(U)$ . Since  $U$  is minimal, there exists an oriented planar graph  $P_{uv}$  such that every homomorphism from  $P_{uv}$  to  $U$  has to use the arc  $uv$  (otherwise, the arc  $uv$  could be deleted from  $U$ ). We now construct the oriented graph  $P^*$  as follows: take one copy of  $P_{uv}$  for every  $uv \in E(U)$  and glue them together by choosing one vertex in each copy and identifying them into a unique vertex. The graph  $P^*$  thus obtained is clearly planar and every homomorphism from  $P^*$  to  $U$  induces a homomorphism from  $P_{uv}$  to  $U$  for every  $uv \in E(U)$ . Therefore, every such homomorphism has to use all the arcs of  $U$  and we are done.  $\square$

**Lemma 4** For every vertex  $u \in V(U)$ ,  $|V(\Gamma_U^+(u))| \geq 7$  and  $|V(\Gamma_U^-(u))| \geq 7$ .

**Proof.** Consider the oriented graph  $P$  obtained from  $P^*$  by adding, for every vertex  $x \in V(P^*)$ , two copies of the oriented graph  $G_3$  (given in Figure 1) together with arcs from  $x$  towards every vertex of the first copy and arcs from every vertex of the second copy towards  $x$ . Since  $G_3$  is outerplanar, the oriented graph  $P$  thus obtained is planar. Every homomorphism from  $P$  to  $U$  has to use every arc, and thus every vertex, of  $U$  since  $P^*$  is a subgraph of  $P$ . Considering the fact that  $\bar{\chi}(G_3) = 7$  we get the desired result.  $\square$

Lemma 4 shows in particular that if  $U$  is an oriented graph on 15 vertices then  $U$  is a tournament and every vertex  $u \in V(U)$  has exactly 7 predecessors and 7 successors.

**Lemma 5** *If  $U$  is a tournament on 15 vertices then for every arc  $uv \in E(U)$ , for every  $A \in \{\Gamma_U^+(u), \Gamma_U^-(u)\}$ ,  $B \in \{\Gamma_U^+(v), \Gamma_U^-(v)\}$ ,  $|V(A \cap B)| \geq 3$ . Moreover, if  $|V(A \cap B)| = 3$  then the subgraph  $A \cap B$  is the directed cycle on three vertices.*

**Proof.** Let  $P_4$  be the directed path on 4 vertices. Every oriented colouring of  $P_4$  has to use at least 3 colours and, if  $c$  is an oriented 3-colouring of  $P_4$ , the colour-graph of  $c$  has to be the directed cycle on 3 vertices.

Let now  $P$  be the oriented graph obtained from  $P^*$  by adding, for every arc  $xy \in E(P^*)$ , four copies of  $P_4$ , say  $P_4^1, P_4^2, P_4^3$  and  $P_4^4$  with arcs from  $x$  towards every vertex of  $P_4^i$  if  $i \in \{1, 2\}$ , arcs from every vertex of  $P_4^i$  towards  $x$  if  $i \in \{3, 4\}$ , arcs from  $y$  towards every vertex of  $P_4^i$  if  $i \in \{1, 3\}$ , and arcs from every vertex of  $P_4^i$  towards  $y$  if  $i \in \{2, 4\}$ . The graph  $P$  thus obtained is clearly planar and, since every homomorphism from  $P$  to  $U$  has to use all the arcs of  $U$ , we get the desired result.  $\square$

Lemma 5 shows in particular that if  $U$  is a tournament on 15 vertices then for any two vertices  $u$  and  $v$  in  $V(U)$ , one of the four subgraphs  $\Gamma_U^+(u) \cap \Gamma_U^+(v)$ ,  $\Gamma_U^+(u) \cap \Gamma_U^-(v)$ ,  $\Gamma_U^-(u) \cap \Gamma_U^+(v)$  and  $\Gamma_U^-(u) \cap \Gamma_U^-(v)$  has exactly four vertices while the three others are the directed cycle on three vertices.

**Lemma 6** *If  $U$  is a tournament on 15 vertices then every three vertices  $u, v, w \in V(U)$  have a common successor.*

**Proof.** We have two cases to consider depending on whether  $u, v, w$  induce a transitive tournament or a directed cycle.

**Case 1.**  $u, v, w$  induce a transitive tournament.

Without loss of generality, we may assume that  $uv, uw, vw \in E(U)$ . If  $\Gamma_U^+(u) \cap \Gamma_U^+(v)$  is the directed cycle on 3 vertices (Lemma 5) we are done since  $w \in V(\Gamma_U^+(u) \cap \Gamma_U^+(v))$  and the successor of  $w$  in this cycle is also a successor of  $u$  and  $v$ .


Suppose now that  $\Gamma_U^+(u) \cap \Gamma_U^+(v)$  has 4 vertices. If  $w$  has a successor in  $\Gamma_U^+(u) \cap \Gamma_U^+(v)$  then this successor is again also a successor of  $u$  and  $v$ . If  $w$  has no successor in  $\Gamma_U^+(u) \cap \Gamma_U^+(v)$  let us count the predecessors of  $w$  in  $U$ :  $w$  has 3 predecessors in  $\Gamma_U^+(u) \cap \Gamma_U^+(v)$ , 3 predecessors in  $\Gamma_U^-(u) \cap \Gamma_U^-(w)$ , which, including  $u$  and  $v$ , gives 8 predecessors. This contradicts Lemma 4.

**Case 2.**  $u, v, w$  induce a directed cycle.

Assume to the contrary that  $u, v$  and  $w$  have no common successor in  $U$ . Considering the structure of the graph  $U$  (see Figure 2), we will show that we necessary get a contradiction to Lemma 4.

Let us denote by  $U_{uv}$  (resp.  $U_{vw}, U_{wu}$ ) the set  $V(\Gamma_U^+(u) \cap \Gamma_U^+(v))$  (resp.  $V(\Gamma_U^+(v) \cap \Gamma_U^+(w)), V(\Gamma_U^+(w) \cap \Gamma_U^+(u))$ ). These three sets are clearly disjoint. By Lemma 5, we know that each of these sets contains at least 3 vertices and, by our assumption, that  $w$  is a successor of  $U_{uv}$ ,  $u$  is a successor of  $U_{vw}$  and  $v$  is a successor of  $U_{wu}$ .

Let us now count the successors of  $u$ :  $v$  is a successor of  $u$ , as well as the vertices of  $U_{uv}$  and  $U_{wu}$ . Therefore,  $U_{uv}$  and  $U_{wu}$  necessarily contain exactly 3 vertices each, otherwise Lemma 4 would be contradicted. Considering in a similar way the successors of  $v$ , or  $w$ , we get that the third set  $U_{vw}$  must also contain exactly 3 vertices. By Lemma 5 we thus get that each of these three sets induces

Figure 2: The structure of the graph  $U$ 

a directed cycle on 3 vertices. Let then  $U_{uv}$  be the directed cycle  $a_1a_2a_3$ ,  $U_{vw}$  be the directed cycle  $b_1b_2b_3$  and  $U_{wu}$  be the directed cycle  $c_1c_2c_3$ .

Let us denote by  $t_1$ ,  $t_2$  and  $t_3$  the three remaining vertices in  $U$ . Since  $u$ ,  $v$  and  $w$  have already 7 successors, the vertices  $t_1$ ,  $t_2$  and  $t_3$  are necessarily predecessors of  $u$ ,  $v$  and  $w$  (otherwise Lemma 4 would be contradicted).

Let us finally consider the common successors of  $u$  and any  $a_i$ ,  $1 \leq i \leq 3$ . We know by Lemma 5 that there are at least 3 such common successors. One of them is  $a_{i+1 \pmod{3}}$  and the at least two others necessarily belong to  $U_{wu}$ . Similarly, considering the common successors of  $u$  and any  $c_i$ ,  $1 \leq i \leq 3$ , we get that each  $c_i$  has at least two successors in  $U_{uv}$ . The number of arcs between  $U_{uv}$  and  $U_{wu}$  is therefore at least 12, a contradiction since each of these sets has only 3 vertices.  $\square$

From Lemmas 4, 5 and 6, we get that if  $U$  is a tournament on 15 vertices then every vertex has 7 successors, every two vertices have at least 3 common successors and every three vertices have at least one common successor. The proof of Theorem 1 follows from the fact that such a tournament cannot exist, as shown by the following result of Brown and Reid (considering the case  $n = 1$ ):

**Theorem 7 ([3])** *There exists no tournament on  $8n + 7$  vertices such that every vertex has at least  $4n + 3$  successors, every two vertices have at least  $2n + 1$  common successors and every three vertices have at least  $n$  common successors.*

## References

- [1] Borodin, O.V., Kostochka, A.V., Nešetřil, J., Raspaud, A., Sopena, E.: On universal graphs for planar oriented graphs of a given girth, *Discrete Math.* **188** (1998), 73–85.
- [2] Borodin, O.V., Kostochka, A.V., Nešetřil, J., Raspaud, A., Sopena, E.: On the maximum average degree and the oriented chromatic number of a graph, *Discrete Math.* **206** (1999), 77–89.
- [3] Brown, E, Reid, K.B.: Doubly regular tournaments are equivalent to skew Hadamard matrices, *J. Combin. Theory* **12** (1972), 332–338.
- [4] Nešetřil, J., Raspaud, A., Sopena, E.: Colorings and girth of oriented planar graphs, *Discrete Math.* **165/166** (1997), 519–530.
- [5] Raspaud, A., Sopena, E.: Good and semi-strong colorings of oriented planar graphs, *Inf. Processing Letters* **51** (1994), 171–174.
- [6] Sopena, E.: The chromatic number of oriented graphs, *J. Graph Theory* **25** (1997), 191–205.

- [7] Sopena, E.: Oriented graph coloring, *Discrete Math.* **229(1-3)** (2001), 359–369.