

Asymptotics of meanders with fixed number of bigons

Vincent Delecroix (joint with É. Goujard, P. Zograf et A. Zorich)


CNRS, LaBRI, Bordeaux

Warwick, march 2018

What is a meander

Definition


A meander is a homotopy class of pairs of transverse smooth curves (γ_1, γ_2) on the sphere.



What is a meander

Definition


A meander is a homotopy class of pairs of transverse smooth curves (γ_1, γ_2) on the sphere.



Following the literature, we will have a marked point on the blue curve so that intersections can be numbered (and avoid complications with automorphisms).


Faces and bigons

A meander cut the plane into faces with an even number of sides $2j$. We call $j - 2$ the *degree*.




Faces and bigons

A meander cut the plane into faces with an even number of sides $2j$. We call $j - 2$ the *degree*.



Faces and bigons


A meander cut the plane into faces with an even number of sides $2j$. We call $j - 2$ the *degree*.



Sum of degrees = -4.

Faces and bigons

A meander cut the plane into faces with an even number of sides $2j$. We call $j - 2$ the *degree*.



Sum of degrees = -4. The *bigons* are the faces of degree -1.

Some numbers

Some numbers

- $M_{n,k}$ = number of meanders with $2n$ crossings and k bigons

Some numbers

- $M_{n,k}$ = number of meanders with $2n$ crossings and k bigons
- $M_n = \sum_k M_{n,k}$ = number of meanders with $2n$ crossings

Some numbers

- $M_{n,k}$ = number of meanders with $2n$ crossings and k bigons
- $M_n = \sum_k M_{n,k}$ = number of meanders with $2n$ crossings

n	$M_{n,4}$	$M_{n,5}$	$M_{n,6}$	$M_{n,7}$	$M_{n,8}$	$M_{n,9}$	$M_{n,10}$	M_n
2	2	0	0	0	0	0	0	2
3	6	0	2	0	0	0	0	8
4	8	16	16	0	2	0	0	42
5	20	40	110	60	30	0	2	262
6	12	168	416	576	462	144	48	1828
7	42	280	1470	3276	4228	2884	1288	13820
8	32	544	4128	13632	26424	31200	22096	110954
9	54	1152	9102	45468	130410	220488	243990	933458

Conjectural asymptotics for M_n

M_n = number of meanders with $2n$ crossings

Conjectural asymptotics for M_n

M_n = number of meanders with $2n$ crossings

Conjecture (Di Francesco-Golinelli-Gutter (2000))

$$M_n \sim C R^n n^{-\alpha}$$

where $C > 0$, $R \simeq 12.2628$ and $\alpha = \frac{29 + \sqrt{145}}{12}$.

Conjectural asymptotics for M_n

M_n = number of meanders with $2n$ crossings

Conjecture (Di Francesco-Golinelli-Guitter (2000))

$$M_n \sim C R^n n^{-\alpha}$$

where $C > 0$, $R \simeq 12.2628$ and $\alpha = \frac{29 + \sqrt{145}}{12}$.

Extensive numerical experiments and extrapolations by Jensen and Guttmann (2000) tend to confirm the conjecture.

Average order of $M_{n,k}$

$M_{n,k}$ = number of meanders with $2n$ crossings and k bigons

Average order of $M_{n,k}$

$M_{n,k}$ = number of meanders with $2n$ crossings and k bigons

Theorem (Delecroix-Goujard-Zograf-Zorich)

For all k we have the following asymptotics

$$\sum_{n \leq x} M_{n,k} \sim \frac{2}{k!(k-4)!} \left(\frac{2}{\pi^2}\right)^{k-3} \binom{2k-4}{k-2}^2 \frac{x^{2k-5}}{2k-5}.$$

Average order of $M_{n,k}$

$M_{n,k}$ = number of meanders with $2n$ crossings and k bigons

Theorem (Delecroix-Goujard-Zograf-Zorich)


For all k we have the following asymptotics

$$\sum_{n \leq x} M_{n,k} \sim \frac{2}{k!(k-4)!} \left(\frac{2}{\pi^2}\right)^{k-3} \binom{2k-4}{k-2}^2 \frac{x^{2k-5}}{2k-5}.$$

Examples


$$\sum_{n \leq x} M_{n,4} \sim \frac{6}{\pi^2} \frac{x^3}{3} \quad \sum_{n \leq x} M_{n,5} \sim \frac{80}{3\pi^4} \frac{x^5}{5} \quad \sum_{n \leq x} M_{n,6} \sim \frac{490}{9\pi^6} \frac{x^7}{7}$$

Meanders as square tiled surfaces of genus 0




A surface in $\mathcal{Q}(1, -1^5)$.

Meanders as square tiled surfaces of genus 0



A surface in $\mathcal{Q}(1, -1^5)$. The **horizontal curves** and **vertical curves** dual to the squares gives a pair of transverse multicurve (γ_1, γ_2) on the sphere.

Meanders as square tiled surfaces of genus 0



A surface in $Q(1, -1^5)$. The **horizontal curves** and **vertical curves** dual to the squares gives a pair of transverse multicurve (γ_1, γ_2) on the sphere. In the example: γ_1 has three components while γ_2 has one.

Meanders as square tiled surfaces of genus 0

Let $\mathcal{Q}_{0,k}$ denote the moduli space of quadratic differentials on the sphere with k poles. We have

$$\mathcal{Q}_{0,4} = \mathcal{Q}(-1^4) \quad \mathcal{Q}_{0,5} = \mathcal{Q}(1, -1^5) \quad \mathcal{Q}_{0,6} = \mathcal{Q}(1^2, -1^6) \cup \mathcal{Q}(2, -1^6)$$

Meanders as square tiled surfaces of genus 0

Let $\mathcal{Q}_{0,k}$ denote the moduli space of quadratic differentials on the sphere with k poles. We have

$$\mathcal{Q}_{0,4} = \mathcal{Q}(-1^4) \quad \mathcal{Q}_{0,5} = \mathcal{Q}(1, -1^5) \quad \mathcal{Q}_{0,6} = \mathcal{Q}(1^2, -1^6) \cup \mathcal{Q}(2, -1^6)$$


Lemma

We have


$$M_{n,k} = n \sum_S \frac{1}{|\text{Aut}(S)|}$$

where the sum is over the square tiled surfaces in $\mathcal{Q}_{0,k}$ whose horizontal and vertical directions are made of a single cylinder of height 1.


Meanders as square tiled surfaces of genus 0 (proof)




Meanders as square tiled surfaces of genus 0 (proof)



Meanders as square tiled surfaces of genus 0 (proof)



Meanders as square tiled surfaces of genus 0 (proof)



Reformulation of the asymptotics theorem

Lemma

We have

$$M_{n,k} = n \sum_S \frac{1}{|\text{Aut}(S)|}$$

where the sum is over the square tiled surfaces in $\mathcal{Q}_{0,k}$ whose horizontal and vertical directions are made of a single cylinder of height 1.

Theorem (Delecroix-Goujard-Zograf-Zorich)

Let $\mathcal{C}_{\mathbb{Z}}^h$ (respectively $\mathcal{C}_{\mathbb{Z}}^v$) denote the set of square tiled surfaces in $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$ that are made of a single horizontal cylinder (resp. vertical cylinder) of height one. Then

$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v : \text{Area}(S) \leq x\} \sim \frac{2}{k!(k-4)!} \left(\frac{2}{\pi^2}\right)^{k-3} \binom{2k-4}{k-2}^2 \frac{x^{2k-6}}{2k-6}.$$

Reformulation of the asymptotics theorem

Theorem (Delecroix-Goujard-Zograf-Zorich)

Let $\mathcal{C}_{\mathbb{Z}}^h$ (respectively $\mathcal{C}_{\mathbb{Z}}^v$) denote the set of square tiled surfaces in $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$ that are made of a single horizontal cylinder (resp. vertical cylinder) of height one. Then

$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v : \text{Area}(S) \leq x\} \sim \frac{2}{k!(k-4)!} \left(\frac{2}{\pi^2}\right)^{k-3} \binom{2k-4}{k-2}^2 \frac{x^{2k-6}}{2k-6}.$$

Proof steps

Reformulation of the asymptotics theorem

Theorem (Delecroix-Goujard-Zograf-Zorich)

Let $\mathcal{C}_{\mathbb{Z}}^h$ (respectively $\mathcal{C}_{\mathbb{Z}}^v$) denote the set of square tiled surfaces in $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$ that are made of a single horizontal cylinder (resp. vertical cylinder) of height one. Then

$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v : \text{Area}(S) \leq x\} \sim \frac{2}{k!(k-4)!} \left(\frac{2}{\pi^2}\right)^{k-3} \binom{2k-4}{k-2}^2 \frac{x^{2k-6}}{2k-6}.$$

Proof steps

1

$$c(\mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v) = \frac{c(\mathcal{C}_{\mathbb{Z}}^h) c(\mathcal{C}_{\mathbb{Z}}^v)}{c(\mathcal{C}_{\mathbb{Z}})}$$

where $c(\mathcal{D})$ stands for the constant so that

$$\#\{S \in \mathcal{D} : \text{Area}(S) \leq x\} \sim c(\mathcal{D}) \frac{x^{2k-6}}{2k-6}.$$

Reformulation of the asymptotics theorem

Theorem (Delecroix-Goujard-Zograf-Zorich)

Let $\mathcal{C}_{\mathbb{Z}}^h$ (respectively $\mathcal{C}_{\mathbb{Z}}^v$) denote the set of square tiled surfaces in $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$ that are made of a single horizontal cylinder (resp. vertical cylinder) of height one. Then

$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v : \text{Area}(S) \leq x\} \sim \frac{2}{k!(k-4)!} \left(\frac{2}{\pi^2}\right)^{k-3} \binom{2k-4}{k-2}^2 \frac{x^{2k-6}}{2k-6}.$$

Proof steps

1

$$c(\mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v) = \frac{c(\mathcal{C}_{\mathbb{Z}}^h) c(\mathcal{C}_{\mathbb{Z}}^v)}{c(\mathcal{C}_{\mathbb{Z}})}$$

where $c(\mathcal{D})$ stands for the constant so that

$$\#\{S \in \mathcal{D} : \text{Area}(S) \leq x\} \sim c(\mathcal{D}) \frac{x^{2k-6}}{2k-6}.$$

2 compute the terms in rhs: $c(\mathcal{C}_{\mathbb{Z}}^h)$, $c(\mathcal{C}_{\mathbb{Z}}^v)$ and $c(\mathcal{C}_{\mathbb{Z}})$.

The case $k = 4$

For $\mathcal{C} = \mathcal{Q}(-1^4)$ we have

$$\#\{S \in \mathcal{C}_{\mathbb{Z}} : \text{Area}(S) = n\} = \sigma(n)$$

$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h : \text{Area}(S) = n\} = n$$

$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v : \text{Area}(S) = n\} = \frac{M_{n,4}}{n} = \phi(n)$$

where ϕ is the Euler totient function and $\sigma(n) = \sum_{d|n} d$.

The case $k = 4$

For $\mathcal{C} = \mathcal{Q}(-1^4)$ we have

$$\#\{S \in \mathcal{C}_{\mathbb{Z}} : \text{Area}(S) = n\} = \sigma(n)$$

$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h : \text{Area}(S) = n\} = n$$


$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v : \text{Area}(S) = n\} = \frac{M_{n,4}}{n} = \phi(n)$$

where ϕ is the Euler totient function and $\sigma(n) = \sum_{d|n} d$. We have


$$\sum_{n \leq x} \phi(n) \sim \frac{6}{\pi^2} x^2 \quad \text{and} \quad \sum_{n \leq x} \sigma(n) \sim \frac{\pi^2}{6} x^2.$$

$$\sigma(n)/n$$

$$\sigma(n)/n = \# \text{ "area } n \text{ s.t.s. in } \mathcal{Q}(-1^4)"/n$$



$$\phi(n)/n$$



The case $k = 5$

For $\mathcal{C} = \mathcal{Q}(1, -1^5)$ we have

$$\#\{S \in \mathcal{C}_{\mathbb{Z}} : \text{Area}(S) = n\} = \frac{3}{8} (\sigma_3(n) - (2n - 1)\sigma(n))$$

$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h : \text{Area}(S) = n\} = \frac{n(n-1)(n-2)}{6}$$

$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v : \text{Area}(S) = n\} = \frac{M_{n,5}}{n} = ??.$$

where $\sigma_3(n) = \sum_{d|n} d^3$.

The case $k = 5$

For $\mathcal{C} = \mathcal{Q}(1, -1^5)$ we have

$$\#\{S \in \mathcal{C}_{\mathbb{Z}} : \text{Area}(S) = n\} = \frac{3}{8} (\sigma_3(n) - (2n-1)\sigma(n))$$


$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h : \text{Area}(S) = n\} = \frac{n(n-1)(n-2)}{6}$$

$$\#\{S \in \mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v : \text{Area}(S) = n\} = \frac{M_{n,5}}{n} = ??.$$


where $\sigma_3(n) = \sum_{d|n} d^3$. For the general $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$ the counting functions

- 1 for $\mathcal{C}_{\mathbb{Z}}$ is expressible in terms of σ ("quasimodularity" by Eskin, Okounkov, Pandharipande, Engels)
- 2 for $\mathcal{C}_{\mathbb{Z}}^h$ is a rational multiple of $\binom{n-k+4}{k-3}$
- 3 is not known for the meanders $(\mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v)$!

"area n s.t.s. in $Q(1, -1^5)$ " / n^3



$$n \mapsto M_{n,5}/n^4$$




Existence of densities

Given a connected component of stratum \mathcal{C} , our aim is now to prove that $c(\mathcal{C}_{\mathbb{Z}})$, $c(\mathcal{C}_{\mathbb{Z}}^h)$, $c(\mathcal{C}_{\mathbb{Z}}^v)$ and $c(\mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v)$ exist and satisfy

$$c(\mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v) = \frac{c(\mathcal{C}_{\mathbb{Z}}^h) c(\mathcal{C}_{\mathbb{Z}}^v)}{c(\mathcal{C}_{\mathbb{Z}})}.$$


Equidistribution in \mathbb{Z}^2

Let $\Lambda = \mathbb{Z}^2$ be the set of integral vectors.



Equidistribution in \mathbb{Z}^2

Let $\Lambda = \mathbb{Z}^2$ be the set of integral vectors.




Theorem

For any relatively compact open set $U \subset \mathbb{R}^2$ we have as $\epsilon \rightarrow 0$ the following asymptotics

$$\#(U \cap (\epsilon\Lambda)) \sim \frac{\text{Vol}(U)}{\epsilon^2}.$$


Equidistribution in \mathbb{Z}^2 (bis)

Let $\Lambda' \subset \mathbb{Z}^2$ be the set of integral vectors with relatively prime entries.



Equidistribution in \mathbb{Z}^2 (bis)

Let $\Lambda' \subset \mathbb{Z}^2$ be the set of integral vectors with relatively prime entries.



Theorem

For any relatively compact open set $U \subset \mathbb{R}^2$ we have as $\epsilon \rightarrow 0$ the following asymptotics

$$\#(U \cap (\epsilon\Lambda')) \sim \frac{6}{\pi^2} \frac{\text{Vol}(U)}{\epsilon^2}.$$

Equidistribution of square tiled surfaces

Let \mathcal{D} be a subset of the set of square tiled surfaces in a component of stratum \mathcal{C} . We say that \mathcal{D} has *uniform density in \mathcal{C}* if there exists a number $\delta(\mathcal{D})$ so that for any open set $U \subset \mathcal{C}$ we have as $\epsilon \rightarrow 0$ the following equivalent

$$\#(U \cap (\epsilon\mathcal{D})) \sim \delta(\mathcal{D}) \frac{\mu_{MV}(U)}{\epsilon^d}$$

where μ_{MV} is the Masur-Veech measure on \mathcal{C} .

Equidistribution of square tiled surfaces

Let \mathcal{D} be a subset of the set of square tiled surfaces in a component of stratum \mathcal{C} . We say that \mathcal{D} has *uniform density in \mathcal{C}* if there exists a number $\delta(\mathcal{D})$ so that for any open set $U \subset \mathcal{C}$ we have as $\epsilon \rightarrow 0$ the following equivalent

$$\#(U \cap (\epsilon\mathcal{D})) \sim \delta(\mathcal{D}) \frac{\mu_{MV}(U)}{\epsilon^d}$$

where μ_{MV} is the Masur-Veech measure on \mathcal{C} .

- 1 By construction of μ_{MV} , the set of all square tiled surfaces has uniform density $\delta = 1$.

Equidistribution of square tiled surfaces

Let \mathcal{D} be a subset of the set of square tiled surfaces in a component of stratum \mathcal{C} . We say that \mathcal{D} has *uniform density in \mathcal{C}* if there exists a number $\delta(\mathcal{D})$ so that for any open set $U \subset \mathcal{C}$ we have as $\epsilon \rightarrow 0$ the following equivalent

$$\#(U \cap (\epsilon\mathcal{D})) \sim \delta(\mathcal{D}) \frac{\mu_{MV}(U)}{\epsilon^d}$$

where μ_{MV} is the Masur-Veech measure on \mathcal{C} .

- 1 By construction of μ_{MV} , the set of all square tiled surfaces has uniform density $\delta = 1$.
- 2 The definition of square tiled surfaces and densities make sense for linear $GL_2(\mathbb{R})$ -invariant manifolds defined over \mathbb{Q} .

Equidistribution of square tiled surfaces

Let \mathcal{D} be a subset of the set of square tiled surfaces in a component of stratum \mathcal{C} . We say that \mathcal{D} has *uniform density in \mathcal{C}* if there exists a number $\delta(\mathcal{D})$ so that for any open set $U \subset \mathcal{C}$ we have as $\epsilon \rightarrow 0$ the following equivalent

$$\#(U \cap (\epsilon\mathcal{D})) \sim \delta(\mathcal{D}) \frac{\mu_{MV}(U)}{\epsilon^d}$$

where μ_{MV} is the Masur-Veech measure on \mathcal{C} .

- 1 By construction of μ_{MV} , the set of all square tiled surfaces has uniform density $\delta = 1$.
- 2 The definition of square tiled surfaces and densities make sense for linear $GL_2(\mathbb{R})$ -invariant manifolds defined over \mathbb{Q} .
- 3 c and δ are equivalent up to the total mass of the unit hyperboloid

$$c(\mathcal{D}) = \delta(\mathcal{D}) \mu_{MV}^{(1)}(\mathcal{C}^{(1)}).$$

Equidistribution of square tiled surfaces

Theorem (Delecroix-Goujard-Zograf-Zorich)

Let \mathcal{C} be a component of stratum of quadratic differentials. Let $\mathcal{C}_{\mathbb{Z}}$ be the square tiled surfaces in \mathcal{C} and $\mathcal{C}_{\mathbb{Z}}^h$ (respectively $\mathcal{C}_{\mathbb{Z}}^v$) be the s.t.s with a single horizontal cylinder with height 1 (resp. vertical cylinder with height 1). Then $\mathcal{C}_{\mathbb{Z}}^h$, $\mathcal{C}_{\mathbb{Z}}^v$ and $\mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v$ have uniform densities and

$$\delta(\mathcal{C}_h \cap \mathcal{C}_v) = \delta(\mathcal{C}_h) \delta(\mathcal{C}_v) = \delta(\mathcal{C}_h)^2.$$

Sketch of proof of the equidistribution result

Our approach follows closely the steps of Mirzakhani for counting simple closed geodesics in hyperbolic surfaces.

Sketch of proof of the equidistribution result

Our approach follows closely the steps of Mirzakhani for counting simple closed geodesics in hyperbolic surfaces.

step 1: density exists as a whole for $\mathcal{C}_{\mathbb{Z}}$, $\mathcal{C}_{\mathbb{Z}}^h$ and $\mathcal{C}_{\mathbb{Z}}^v$

$$\#\{S \in \mathcal{D} : \text{Area}(S) \leq x\} \sim c(\mathcal{D}) \frac{x^d}{2d}.$$

for some (more or less) explicit $c(\mathcal{D}) > 0$.

Sketch of proof of the equidistribution result

Our approach follows closely the steps of Mirzakhani for counting simple closed geodesics in hyperbolic surfaces.

step 1: density exists as a whole for $\mathcal{C}_{\mathbb{Z}}$, $\mathcal{C}_{\mathbb{Z}}^h$ and $\mathcal{C}_{\mathbb{Z}}^v$

$$\#\{S \in \mathcal{D} : \text{Area}(S) \leq x\} \sim c(\mathcal{D}) \frac{x^d}{2d}.$$

for some (more or less) explicit $c(\mathcal{D}) > 0$.

step 2: from density to uniform density via ergodicity of $\begin{pmatrix} 1 & \mathbb{Z} \\ 0 & 1 \end{pmatrix}$ or $\begin{pmatrix} 1 & 0 \\ \mathbb{Z} & 1 \end{pmatrix}$.

Sketch of proof of the equidistribution result

Our approach follows closely the steps of Mirzakhani for counting simple closed geodesics in hyperbolic surfaces.

step 1: density exists as a whole for $\mathcal{C}_{\mathbb{Z}}$, $\mathcal{C}_{\mathbb{Z}}^h$ and $\mathcal{C}_{\mathbb{Z}}^v$

$$\#\{S \in \mathcal{D} : \text{Area}(S) \leq x\} \sim c(\mathcal{D}) \frac{x^d}{2d}.$$

for some (more or less) explicit $c(\mathcal{D}) > 0$.

step 2: from density to uniform density via ergodicity of $\begin{pmatrix} 1 & \mathbb{Z} \\ 0 & 1 \end{pmatrix}$ or $\begin{pmatrix} 1 & 0 \\ \mathbb{Z} & 1 \end{pmatrix}$.

step 3: the intersection $\mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v$ satisfies $\delta(\mathcal{C}_{\mathbb{Z}}^h \cap \mathcal{C}_{\mathbb{Z}}^v) = \delta(\mathcal{C}_{\mathbb{Z}}^h) \delta(\mathcal{C}_{\mathbb{Z}}^v)$ because of the product structure of \mathcal{C} .

Explicit numbers in genus 0

Theorem

Let $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$ and $\mathcal{C}_{\mathbb{Z}}^h$ be the set of square tiled surfaces in \mathcal{C} with a single horizontal cylinder of height. Then

① (Athreya-Eskin-Zorich)

$$c(\mathcal{C}_{\mathbb{Z}}) = \mu_{MV}^{(1)}(\mathcal{Q}(1^{k-4}, -1^k)) = \frac{\pi^{2k-6}}{2^{k-5}} = \frac{1}{4} \left(\frac{\pi^2}{2} \right)^{k-3}$$

② (Delecroix-Goujard-Zograf-Zorich)

$$c(\mathcal{C}_{\mathbb{Z}}^h) = 2 \binom{2k-4}{k-2}.$$

Explicit numbers in genus 0

Theorem

Let $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$ and $\mathcal{C}_{\mathbb{Z}}^h$ be the set of square tiled surfaces in \mathcal{C} with a single horizontal cylinder of height. Then

① (Athreya-Eskin-Zorich)

$$c(\mathcal{C}_{\mathbb{Z}}) = \mu_{MV}^{(1)}(\mathcal{Q}(1^{k-4}, -1^k)) = \frac{\pi^{2k-6}}{2^{k-5}} = \frac{1}{4} \left(\frac{\pi^2}{2} \right)^{k-3}$$

② (Delecroix-Goujard-Zograf-Zorich)

$$c(\mathcal{C}_{\mathbb{Z}}^h) = 2 \binom{2k-4}{k-2}.$$

$$\sum_{n \leq x} M_{n,k} \sim \frac{2}{k!(k-4)!} \left(\frac{2}{\pi^2} \right)^{k-3} \binom{2k-4}{k-2}^2 \frac{x^{2k-5}}{2k-5}.$$

The computation of $\mathcal{C}_{\mathbb{Z}}$ (Athreya-Eskin-Zorich)

As usual $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$.

The computation of $\mathcal{C}_{\mathbb{Z}}$ (Athreya-Eskin-Zorich)

As usual $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$.

The formula $c(\mathcal{C}_{\mathbb{Z}}) = \mu_{MV}^{(1)}(\mathcal{Q}(1^{k-4}, -1^k)) = \frac{1}{4} \left(\frac{\pi^2}{2}\right)^{k-3}$ was a conjecture of Kontsevich.

The computation of $\mathcal{C}_{\mathbb{Z}}$ (Athreya-Eskin-Zorich)

As usual $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$.

The formula $c(\mathcal{C}_{\mathbb{Z}}) = \mu_{MV}^{(1)}(\mathcal{Q}(1^{k-4}, -1^k)) = \frac{1}{4} \left(\frac{\pi^2}{2}\right)^{k-3}$ was a conjecture of Kontsevich.

The generating series for the square tiled surfaces $\mathcal{C}_{\mathbb{Z}}$ is quasimodular, though it seems hard to obtain the formula with this approach.

The computation of $\mathcal{C}_{\mathbb{Z}}$ (Athreya-Eskin-Zorich)

As usual $\mathcal{C} = \mathcal{Q}(1^{k-4}, -1^k)$.


The formula $c(\mathcal{C}_{\mathbb{Z}}) = \mu_{MV}^{(1)}(\mathcal{Q}(1^{k-4}, -1^k)) = \frac{1}{4} \left(\frac{\pi^2}{2}\right)^{k-3}$ was a conjecture of Kontsevich.

The generating series for the square tiled surfaces $\mathcal{C}_{\mathbb{Z}}$ is quasimodular, though it seems hard to obtain the formula with this approach.

The proof of Athreya-Eskin-Zorich is quite convoluted and uses the formula for the sum of Lyapunov exponents Λ^+ (Eskin-Kontsevich-Zorich) and the fact that on the sphere $\Lambda^+ = 0$ (bundle has dimension 0).

The computation of $\mathcal{C}_{\mathbb{Z}}^h$ (Delecroix-Goujard-Zograf-Zorich)


Let us consider the case of $\mathcal{Q}(1, -1^5)$.



$$\{(\lambda_A, \lambda_B, \lambda_C, \lambda_D, \tau) \in \mathbb{Z}_{>0} : \lambda_A = \lambda_B + \lambda_C + \lambda_D \text{ and } 0 \leq \tau < \lambda_A\}$$

The computation of $\mathcal{C}_{\mathbb{Z}}^h$ (Delecroix-Goujard-Zograf-Zorich)

Let us consider the case of $\mathcal{Q}(1, -1^5)$.




$$\{(\lambda_A, \lambda_B, \lambda_C, \lambda_D, \tau) \in \mathbb{Z}_{>0} : \lambda_A = \lambda_B + \lambda_C + \lambda_D \text{ and } 0 \leq \tau < \lambda_A\}$$

For fixed area n we got $\frac{1}{3} \binom{n}{2} n$ such surfaces.

The computation of $\mathcal{C}_{\mathbb{Z}}^h$ (Delecroix-Goujard-Zograf-Zorich)

Let us consider the case of $\mathcal{Q}(1, -1^5)$.



$$\{(\lambda_A, \lambda_B, \lambda_C, \lambda_D, \tau) \in \mathbb{Z}_{>0} : \lambda_A = \lambda_B + \lambda_C + \lambda_D \text{ and } 0 \leq \tau < \lambda_A\}$$

For fixed area n we got $\frac{1}{3} \binom{n}{2} n$ such surfaces.

In general we have to make a count over all separatrix diagrams that are counted by trees.

References

- S. K. Lando, A. K. Zvonkin *Plane and projective meanders* Theoret. Comput. Sci. 117 (1993).
- P. Di Francesco, O. Golinelli, E. Guitter
 - ▶ *Meanders: a direct enumeration approach* Nuclear Phys. B 482 (1996).
 - ▶ *Meander, folding, and arch statistics* Combinatorics and physics, Math. Comput. Modelling 26 (1997).
 - ▶ *Meanders and the Temperley-Lieb algebra* Comm. Math. Phys. 186 (1997).
 - ▶ *Meanders: exact asymptotics* Nuclear Phys. B 570 (2000).
- I. Jensen, Iwan *A transfer matrix approach to the enumeration of plane meanders* J. Phys. A 33 (2000).
- I. Jensen, A. J. Guttmann *Critical exponents of plane meanders* J. Phys. A 33 (2000).
- V. Delecroix, É. Goujard, P. Zograf A. Zorich *Enumeration of meanders and Masur-Veech volumes* <https://arxiv.org/abs/1705.05190>