

Algebraic extensions of finitely generated subgroups of free groups¹

Pascal Weil

LaBRI, Université Bordeaux-1 and CNRS

CSL, Lisboa, July 2005

¹Joint work with E. Ventura (Barcelona) and A. Myasnikov (Montréal)

Subgroups of free groups

Congruent subgroups

Takahasi's theorem

Representing subgroups of free groups

A 3-line proof

Algebraic extensions of a subgroup

What they are

About terminology

Composition of extensions

The semilattice of algebraic extensions

Elementary algebraic extensions

Some applications

Malnormality, purity, etc

Profinite topologies

Equations

Free groups

A an alphabet; $F(A)$, the free group over A , is the set of *reduced* words over $A \cup A^{-1}$, without factors aa^{-1} or $a^{-1}a$

Free groups

A an alphabet; $F(A)$, the free group over A , is the set of *reduced* words over $A \cup A^{-1}$, without factors aa^{-1} or $a^{-1}a$

- ▶ $F(A)$ has several bases, but all have the same cardinality, the *rank* of $F(A)$

Free groups

A an alphabet; $F(A)$, the free group over A , is the set of *reduced* words over $A \cup A^{-1}$, without factors aa^{-1} or $a^{-1}a$

- ▶ $F(A)$ has several bases, but all have the same cardinality, the *rank* of $F(A)$
- ▶ Subgroups of free groups are free, but

Free groups

A an alphabet; $F(A)$, the free group over A , is the set of *reduced* words over $A \cup A^{-1}$, without factors aa^{-1} or $a^{-1}a$

- ▶ $F(A)$ has several bases, but all have the same cardinality, the *rank* of $F(A)$
- ▶ Subgroups of free groups are free, but
- ▶ F_2 contains F_n for every $n \geq 1$, and also F_∞

Free groups

A an alphabet; $F(A)$, the free group over A , is the set of *reduced* words over $A \cup A^{-1}$, without factors aa^{-1} or $a^{-1}a$

- ▶ $F(A)$ has several bases, but all have the same cardinality, the *rank* of $F(A)$
- ▶ Subgroups of free groups are free, but
- ▶ F_2 contains F_n for every $n \geq 1$, and also F_∞
- ▶ $a_n \mapsto a^n b^n$ is injective from F_∞ to F_2

A general problem

How does a (finitely generated) free group sit as a subgroup in another?

A general problem

How does a (finitely generated) free group sit as a subgroup in another?

If A is a basis of F and $B \subseteq A$, $\langle B \rangle$ is a *free factor* of F , $\langle B \rangle \leq_{\text{ff}} F$

A general problem

How does a (finitely generated) free group sit as a subgroup in another?

If A is a basis of F and $B \subseteq A$, $\langle B \rangle$ is a *free factor* of F , $\langle B \rangle \leq_{\text{ff}} F$

▶ In $F = F(a, b, c)$, $H = \langle a, b \rangle \leq_{\text{ff}} F$

A general problem

How does a (finitely generated) free group sit as a subgroup in another?

If A is a basis of F and $B \subseteq A$, $\langle B \rangle$ is a *free factor* of F , $\langle B \rangle \leq_{\text{ff}} F$

- ▶ In $F = F(a, b, c)$, $H = \langle a, b \rangle \leq_{\text{ff}} F$
- ▶ but also $K = \langle ab^2, bac \rangle \leq_{\text{ff}} F$ since $\{ab^2, bac, b\}$ is a basis of F

A general problem

How does a (finitely generated) free group sit as a subgroup in another?

If A is a basis of F and $B \subseteq A$, $\langle B \rangle$ is a *free factor* of F , $\langle B \rangle \leq_{\text{ff}} F$

- ▶ In $F = F(a, b, c)$, $H = \langle a, b \rangle \leq_{\text{ff}} F$
- ▶ but also $K = \langle ab^2, bac \rangle \leq_{\text{ff}} F$ since $\{ab^2, bac, b\}$ is a basis of F
- ▶ H and K both have rank 2 and if we define $\varphi: F \rightarrow F$ by $\varphi(a) = ab^2$, $\varphi(b) = bac$ and $\varphi(c) = b$, then $\varphi \in \text{Aut}(F)$ and $\varphi(H) = K$

A general problem

How does a (finitely generated) free group sit as a subgroup in another?

If A is a basis of F and $B \subseteq A$, $\langle B \rangle$ is a *free factor* of F , $\langle B \rangle \leq_{\text{ff}} F$

- ▶ In $F = F(a, b, c)$, $H = \langle a, b \rangle \leq_{\text{ff}} F$
- ▶ but also $K = \langle ab^2, bac \rangle \leq_{\text{ff}} F$ since $\{ab^2, bac, b\}$ is a basis of F
- ▶ H and K both have rank 2 and if we define $\varphi: F \rightarrow F$ by $\varphi(a) = ab^2$, $\varphi(b) = bac$ and $\varphi(c) = b$, then $\varphi \in \text{Aut}(F)$ and $\varphi(H) = K$

In that case, say that H and K are *congruent*

A general problem

How does a (finitely generated) free group sit as a subgroup in another?

If A is a basis of F and $B \subseteq A$, $\langle B \rangle$ is a *free factor* of F , $\langle B \rangle \leq_{\text{ff}} F$

- ▶ In $F = F(a, b, c)$, $H = \langle a, b \rangle \leq_{\text{ff}} F$
- ▶ but also $K = \langle ab^2, bac \rangle \leq_{\text{ff}} F$ since $\{ab^2, bac, b\}$ is a basis of F
- ▶ H and K both have rank 2 and if we define $\varphi: F \rightarrow F$ by $\varphi(a) = ab^2$, $\varphi(b) = bac$ and $\varphi(c) = b$, then $\varphi \in \text{Aut}(F)$ and $\varphi(H) = K$

In that case, say that H and K are *congruent*

Thus: *two free factors of F are congruent if and only if they have the same rank*

For more general subgroups of F , the equality of ranks is necessary but not sufficient

For more general subgroups of F , the equality of ranks is necessary but not sufficient

- ▶ They must have the same index, and all kinds of other invariants must match

For more general subgroups of F , the equality of ranks is necessary but not sufficient

- ▶ They must have the same index, and all kinds of other invariants must match
- ▶ For instance, if H and K are congruent, H and K must have congruent radical closures \sqrt{H} and \sqrt{K} , where

$$\sqrt{H} = \langle x \in F \mid x^n \in H \text{ for some } n \in \mathbb{N} \rangle$$

For more general subgroups of F , the equality of ranks is necessary but not sufficient

- ▶ They must have the same index, and all kinds of other invariants must match
- ▶ For instance, if H and K are congruent, H and K must have congruent radical closures \sqrt{H} and \sqrt{K} , where

$$\sqrt{H} = \langle x \in F \mid x^n \in H \text{ for some } n \in \mathbb{N} \rangle$$

- ▶ There exists a nice Whitehead-style algorithm to decide whether two finitely generated subgroups of F are congruent (Gersten)

For more general subgroups of F , the equality of ranks is necessary but not sufficient

- ▶ They must have the same index, and all kinds of other invariants must match
- ▶ For instance, if H and K are congruent, H and K must have congruent radical closures \sqrt{H} and \sqrt{K} , where

$$\sqrt{H} = \langle x \in F \mid x^n \in H \text{ for some } n \in \mathbb{N} \rangle$$

- ▶ There exists a nice Whitehead-style algorithm to decide whether two finitely generated subgroups of F are congruent (Gersten)

but this algorithm is not completely satisfactory: if H and K are not congruent, it does not always tell you why – we would like a classification of the congruence classes of subgroups; ideally, a computable list of invariants of a subgroup H , characterizing its congruence class

We don't solve this problem, but discuss one such invariant, the *semilattice of algebraic extensions of H* , which

We don't solve this problem, but discuss one such invariant, the *semilattice of algebraic extensions of H* , which

- ▶ must be preserved by the congruence relation

We don't solve this problem, but discuss one such invariant, the *semilattice of algebraic extensions of H* , which

- ▶ must be preserved by the congruence relation
- ▶ is computable

We don't solve this problem, but discuss one such invariant, the *semilattice of algebraic extensions of H* , which

- ▶ must be preserved by the congruence relation
- ▶ is computable
- ▶ describes the way H sits in larger subgroups of F

We don't solve this problem, but discuss one such invariant, the *semilattice of algebraic extensions of H* , which

- ▶ must be preserved by the congruence relation
- ▶ is computable
- ▶ describes the way H sits in larger subgroups of F
- ▶ is related with Takahasi's theorem (1951)

Takahasi's theorem

Theorem

Let $H \leq_{fg} F$ (H is a finitely generated subgroup of F).

Takahasi's theorem

Theorem

Let $H \leq_{fg} F$ (H is a finitely generated subgroup of F).

There exist H_1, \dots, H_n , finitely many subgroups such that

Takahasi's theorem

Theorem

Let $H \leq_{fg} F$ (H is a finitely generated subgroup of F).

There exist H_1, \dots, H_n , finitely many subgroups such that

If $H \leq K \leq F$, then $H \leq H_i \leq_{ff} K$ for some $1 \leq i \leq n$

Takahasi's theorem

Theorem

Let $H \leq_{fg} F$ (H is a finitely generated subgroup of F).

There exist H_1, \dots, H_n , finitely many subgroups such that

If $H \leq K \leq F$, then $H \leq H_i \leq_{ff} K$ for some $1 \leq i \leq n$

- ▶ Many fun applications: as an exercise, show that if $(H_i)_{i \geq 0}$ is a chain of subgroups of F with bounded rank (there exists n such that $\text{rk}(H_i) \leq n$ for each i), then (H_i) is eventually constant

Takahasi's theorem

Theorem

Let $H \leq_{fg} F$ (H is a finitely generated subgroup of F).

There exist H_1, \dots, H_n , finitely many subgroups such that

If $H \leq K \leq F$, then $H \leq H_i \leq_{ff} K$ for some $1 \leq i \leq n$

- ▶ Many fun applications: as an exercise, show that if $(H_i)_{i \geq 0}$ is a chain of subgroups of F with bounded rank (there exists n such that $\text{rk}(H_i) \leq n$ for each i), then (H_i) is eventually constant
- ▶ Such a family H_1, \dots, H_n (a Takahasi family for H), can be effectively computed

Takahasi's theorem

Theorem

Let $H \leq_{fg} F$ (H is a finitely generated subgroup of F).

There exist H_1, \dots, H_n , finitely many subgroups such that

If $H \leq K \leq F$, then $H \leq H_i \leq_{ff} K$ for some $1 \leq i \leq n$


- ▶ Many fun applications: as an exercise, show that if $(H_i)_{i \geq 0}$ is a chain of subgroups of F with bounded rank (there exists n such that $\text{rk}(H_i) \leq n$ for each i), then (H_i) is eventually constant
- ▶ Such a family H_1, \dots, H_n (a Takahasi family for H), can be effectively computed
- ▶ so let me digress on the representation of finitely generated subgroups

Representing a finitely generated subgroup

$$H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$$


Representing a finitely generated subgroup

$$H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$$




Representing a finitely generated subgroup

$$H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$$




Representing a finitely generated subgroup

$$H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$$




Representing a finitely generated subgroup

$$H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$$




Representing a finitely generated subgroup

$$H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$$



Representing a finitely generated subgroup


$$H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$$



- ▶ a deterministic, co-deterministic automaton $\Gamma_A(H)$, with a distinguished vertex, and every non-distinguished vertex has valence at least 2, denoted $\Gamma_A(H)$


Representing a finitely generated subgroup

$$H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$$




- ▶ a deterministic, co-deterministic automaton $\Gamma_A(H)$, with a distinguished vertex, and every non-distinguished vertex has valence at least 2, denoted $\Gamma_A(H)$
- ▶ $\Gamma_A(H)$ depends on H , not on the given generators of H – but depends on A !

Spanning tree, rank and basis




Spanning tree, rank and basis



Choose a **spanning tree**


Spanning tree, rank and basis



Choose a **spanning tree**

For each edge of $\Gamma_A(H)$ not in the spanning tree, construct a basis element for H

Spanning tree, rank and basis




Choose a **spanning tree**

For each edge of $\Gamma_A(H)$ not in the spanning tree, construct a basis element for H

► b^2

Spanning tree, rank and basis




Choose a **spanning tree**

For each edge of $\Gamma_A(H)$ not in the spanning tree, construct a basis element for H

- ▶ b^2
- ▶ ab^2a^{-1}

Spanning tree, rank and basis




Choose a **spanning tree**

For each edge of $\Gamma_A(H)$ not in the spanning tree, construct a basis element for H

- ▶ b^2
- ▶ ab^2a^{-1}
- ▶ $ab^{-1}ab^{-1}$

Spanning tree, rank and basis



Choose a **spanning tree**

For each edge of $\Gamma_A(H)$ not in the spanning tree, construct a basis element for H

- ▶ b^2
- ▶ ab^2a^{-1}
- ▶ $ab^{-1}ab^{-1}$



A spanning tree has $|V| - 1$ edges, so $\text{rk}(H) = |E| - |V| + 1$

Morphisms of automata

Consider $H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$ and $K = \langle b, aba, ab^{-1}a, a^4 \rangle$
Note that $H \leq K$



Morphisms of automata

Consider $H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$ and $K = \langle b, aba, ab^{-1}a, a^4 \rangle$
Note that $H \leq K$



Morphisms of automata

Consider $H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$ and $K = \langle b, aba, ab^{-1}a, a^4 \rangle$
 Note that $H \leq K$





There is a graph morphism $\varphi_{H,K}: \Gamma_A(H) \rightarrow \Gamma_A(K)$

► $1, 3 \mapsto 1, 2 \mapsto 2, 4 \mapsto 4$

Morphisms of automata



Consider $H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$ and $K = \langle b, aba, ab^{-1}a, a^4 \rangle$
 Note that $H \leq K$



There is a graph morphism $\varphi_{H,K}: \Gamma_A(H) \rightarrow \Gamma_A(K)$ if and only if $H \leq K$

Morphisms of automata

Consider $H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$ and $K = \langle b, aba, ab^{-1}a, a^4 \rangle$
 Note that $H \leq K$





There is a graph morphism $\varphi_{H,K}: \Gamma_A(H) \rightarrow \Gamma_A(K)$ if and only if $H \leq K$

- ▶ If $\varphi_{H,K}$ is one-to-one, then $H \leq_{\text{ff}} K$

Morphisms of automata

Consider $H = \langle abab^{-1}, abab, ab^{-1}ab \rangle$ and $K = \langle b, aba, ab^{-1}a, a^4 \rangle$
 Note that $H \leq K$



There is a graph morphism $\varphi_{H,K}: \Gamma_A(H) \rightarrow \Gamma_A(K)$ if and only if $H \leq K$

- ▶ If $\varphi_{H,K}$ is one-to-one, then $H \leq_{\text{ff}} K$
- ▶ If $\varphi_{H,K}$ is onto, we say that K is an *overgroup* of H (a finite number of them, which depends on A)

Proof of Takahasi's theorem

Theorem

Let $H \leq_{fg} F$. There exist H_1, \dots, H_n , finitely many subgroups such that: if $H \leq K \leq F$, then $H \leq H_i \leq_{ff} K$ for some $1 \leq i \leq n$

Proof of Takahasi's theorem

Theorem

Let $H \leq_{fg} F$. There exist H_1, \dots, H_n , finitely many subgroups such that: if $H \leq K \leq F$, then $H \leq H_i \leq_{ff} K$ for some $1 \leq i \leq n$

If $H \leq K$, then there exists $\varphi_{H,K}: \Gamma_A(H) \rightarrow \Gamma_A(K)$. Let L such that $\Gamma_A(L) = \varphi_{H,K}(\Gamma_A(H))$. Then L is an overgroup of H , and $L \leq_{ff} K$.

Proof of Takahasi's theorem

Theorem

Let $H \leq_{fg} F$. There exist H_1, \dots, H_n , finitely many subgroups such that: if $H \leq K \leq F$, then $H \leq H_i \leq_{ff} K$ for some $1 \leq i \leq n$

If $H \leq K$, then there exists $\varphi_{H,K}: \Gamma_A(H) \rightarrow \Gamma_A(K)$. Let L such that $\Gamma_A(L) = \varphi_{H,K}(\Gamma_A(H))$. Then L is an overgroup of H , and $L \leq_{ff} K$.

- ▶ That's it! The set of overgroups of H is a Takahasi family for H : if $H \leq K$, then K is a free extension of an overgroup of H – besides, H has finitely many overgroups, and they are computable.

Proof of Takahasi's theorem

Theorem

Let $H \leq_{fg} F$. There exist H_1, \dots, H_n , finitely many subgroups such that: if $H \leq K \leq F$, then $H \leq H_i \leq_{ff} K$ for some $1 \leq i \leq n$

If $H \leq K$, then there exists $\varphi_{H,K}: \Gamma_A(H) \rightarrow \Gamma_A(K)$. Let L such that $\Gamma_A(L) = \varphi_{H,K}(\Gamma_A(H))$. Then L is an overgroup of H , and $L \leq_{ff} K$.

- ▶ That's it! The set of overgroups of H is a Takahasi family for H : if $H \leq K$, then K is a free extension of an overgroup of H – besides, H has finitely many overgroups, and they are computable.
- ▶ But it is not a canonical family

Subgroups of free groups

Congruent subgroups

Takahasi's theorem

Representing subgroups of free groups

A 3-line proof

Algebraic extensions of a subgroup

What they are

About terminology

Composition of extensions

The semilattice of algebraic extensions

Elementary algebraic extensions

Some applications

Malnormality, purity, etc

Profinite topologies

Equations

Algebraic extensions

Let $H \leq K \leq F$. Say that K is an *algebraic extension* of H ($H \leq_{\text{alg}} K$) if no proper free factor of K contains H , that is,
 $H \leq L \leq_{\text{ff}} K \implies L = K$

Algebraic extensions

Let $H \leq K \leq F$. Say that K is an *algebraic extension* of H ($H \leq_{\text{alg}} K$) if no proper free factor of K contains H , that is,
 $H \leq L \leq_{\text{ff}} K \implies L = K$

Let $\text{AE}(H)$ be the set of algebraic extensions of H

Algebraic extensions

Let $H \leq K \leq F$. Say that K is an *algebraic extension* of H ($H \leq_{\text{alg}} K$) if no proper free factor of K contains H , that is,
 $H \leq L \leq_{\text{ff}} K \implies L = K$

Let $\text{AE}(H)$ be the set of algebraic extensions of H

Proposition $\text{AE}(H)$ is the least Takahasi family for H . In fact, if \mathcal{L} is a Takahasi family for H , then $\text{AE}(H)$ is the set of \leq_{ff} -minimal elements of \mathcal{L} .

Algebraic extensions

Let $H \leq K \leq F$. Say that K is an *algebraic extension* of H ($H \leq_{\text{alg}} K$) if no proper free factor of K contains H , that is,
 $H \leq L \leq_{\text{ff}} K \implies L = K$

Let $\text{AE}(H)$ be the set of algebraic extensions of H

Proposition $\text{AE}(H)$ is the least Takahasi family for H . In fact, if \mathcal{L} is a Takahasi family for H , then $\text{AE}(H)$ is the set of \leq_{ff} -minimal elements of \mathcal{L} .

- ▶ If $H \leq_{\text{alg}} K$, there exists $L \in \mathcal{L}$ such that $H \leq L \leq_{\text{ff}} K$, so $L = K$, and $\text{AE}(H) \subseteq \mathcal{L}$

Algebraic extensions

Let $H \leq K \leq F$. Say that K is an *algebraic extension* of H ($H \leq_{\text{alg}} K$) if no proper free factor of K contains H , that is,
 $H \leq L \leq_{\text{ff}} K \implies L = K$

Let $\text{AE}(H)$ be the set of algebraic extensions of H

Proposition $\text{AE}(H)$ is the least Takahasi family for H . In fact, if \mathcal{L} is a Takahasi family for H , then $\text{AE}(H)$ is the set of \leq_{ff} -minimal elements of \mathcal{L} .

- ▶ If $H \leq_{\text{alg}} K$, there exists $L \in \mathcal{L}$ such that $H \leq L \leq_{\text{ff}} K$, so $L = K$, and $\text{AE}(H) \subseteq \mathcal{L}$
- ▶ If K is \leq_{ff} -minimal in \mathcal{L} and $H \leq L \leq_{\text{ff}} K$, then $H \leq M \leq_{\text{ff}} L$ for some $M \in \mathcal{L}$, so $M \leq_{\text{ff}} K$, and by minimality, $M = L = K$, that is, $K \in \text{AE}(H)$

Examples

▶ $\langle x^n \rangle \leq_{\text{alg}} \langle x \rangle$

Examples

- ▶ $\langle x^n \rangle \leq_{\text{alg}} \langle x \rangle$
- ▶ If $H \leq F_2$ and H is non-cyclic, then $H \leq_{\text{alg}} F_2$

Examples

- ▶ $\langle x^n \rangle \leq_{\text{alg}} \langle x \rangle$
- ▶ If $H \leq F_2$ and H is non-cyclic, then $H \leq_{\text{alg}} F_2$
- ▶ $\langle aba^{-1}b^{-1} \rangle \leq_{\text{alg}} F_2$ (more generally, $\langle x \rangle$ where x is not a power of a primitive element)

Examples

- ▶ $\langle x^n \rangle \leq_{\text{alg}} \langle x \rangle$
- ▶ If $H \leq F_2$ and H is non-cyclic, then $H \leq_{\text{alg}} F_2$
- ▶ $\langle aba^{-1}b^{-1} \rangle \leq_{\text{alg}} F_2$ (more generally, $\langle x \rangle$ where x is not a power of a primitive element)
- ▶ If $H \leq_{\text{fi}} F$, then every K such that $H \leq K \leq F$ is an algebraic extension

Examples

- ▶ $\langle x^n \rangle \leq_{\text{alg}} \langle x \rangle$
- ▶ If $H \leq F_2$ and H is non-cyclic, then $H \leq_{\text{alg}} F_2$
- ▶ $\langle aba^{-1}b^{-1} \rangle \leq_{\text{alg}} F_2$ (more generally, $\langle x \rangle$ where x is not a power of a primitive element)
- ▶ If $H \leq_{\text{fi}} F$, then every K such that $H \leq K \leq F$ is an algebraic extension
- ▶ If $H \leq_{\text{ff}} F$, then $\text{AE}(H) = \{H\}$. For instance, no non-trivial overgroup of $\langle ab, aca \rangle$ is an algebraic extension

Computing the algebraic extensions

- ▶ Note that $AE(H)$ is finite.

Computing the algebraic extensions

- ▶ Note that $AE(H)$ is finite.
- ▶ To compute $AE(H)$, first compute the overgroups of H

Computing the algebraic extensions

- ▶ Note that $AE(H)$ is finite.
- ▶ To compute $AE(H)$, first compute the overgroups of H
- ▶ then decide which are the \leq_{ff} -minimal elements. . .

Computing the algebraic extensions

- ▶ Note that $AE(H)$ is finite.
- ▶ To compute $AE(H)$, first compute the overgroups of H
- ▶ then decide which are the \leq_{ff} -minimal elements. . .
- ▶ . . . using the Whitehead algorithm, which determines whether a subgroup is a free factor in another

About terminology

- ▶ Let $H \leq K \leq F$ and $x \in K$. Say x is K -algebraic over H if every free factor of K containing H , also contains x :

About terminology

- ▶ Let $H \leq K \leq F$ and $x \in K$. Say x is K -algebraic over H if every free factor of K containing H , also contains x :
$$H \leq L \leq_{\text{ff}} K \implies x \in L$$

About terminology

- ▶ Let $H \leq K \leq F$ and $x \in K$. Say x is K -algebraic over H if every free factor of K containing H , also contains x :
$$H \leq L \leq_{\text{ff}} K \implies x \in L$$
- ▶ Otherwise, x is K -transcendental over H . In fact, this is if and only if $\langle H, x \rangle = H * \langle x \rangle$

About terminology

- ▶ Let $H \leq K \leq F$ and $x \in K$. Say x is K -algebraic over H if every free factor of K containing H , also contains x :
$$H \leq L \leq_{\text{ff}} K \implies x \in L$$
- ▶ Otherwise, x is K -transcendental over H . In fact, this is if and only if $\langle H, x \rangle = H * \langle x \rangle$
- ▶ Every element of $K \setminus H$ is K -transcendental over H iff $H \leq_{\text{ff}} K$.

About terminology

- ▶ Let $H \leq K \leq F$ and $x \in K$. Say x is K -algebraic over H if every free factor of K containing H , also contains x :
$$H \leq L \leq_{\text{ff}} K \implies x \in L$$
- ▶ Otherwise, x is K -transcendental over H . In fact, this is if and only if $\langle H, x \rangle = H * \langle x \rangle$
- ▶ Every element of $K \setminus H$ is K -transcendental over H iff $H \leq_{\text{ff}} K$.
- ▶ Let $H \leq K$. Then $H \leq_{\text{alg}} K$ iff every element of K is K -algebraic over H

About terminology

- ▶ Let $H \leq K \leq F$ and $x \in K$. Say x is K -algebraic over H if every free factor of K containing H , also contains x :
$$H \leq L \leq_{\text{ff}} K \implies x \in L$$
- ▶ Otherwise, x is K -transcendental over H . In fact, this is if and only if $\langle H, x \rangle = H * \langle x \rangle$
- ▶ Every element of $K \setminus H$ is K -transcendental over H iff $H \leq_{\text{ff}} K$.
- ▶ Let $H \leq K$. Then $H \leq_{\text{alg}} K$ iff every element of K is K -algebraic over H

Note that the property of being K -algebraic over H , depends on K

Composition of extensions

Let $H \leq K$, and let $H \leq K_i \leq K$, $i = 1, 2$

(i) If $H \leq_{\text{alg}} K_1 \leq_{\text{alg}} K$ then $H \leq_{\text{alg}} K$.

Composition of extensions

Let $H \leq K$, and let $H \leq K_i \leq K$, $i = 1, 2$

(i) If $H \leq_{\text{alg}} K_1 \leq_{\text{alg}} K$ then $H \leq_{\text{alg}} K$.

(i') If $H \leq_{\text{ff}} K_1 \leq_{\text{ff}} K$ then $H \leq_{\text{ff}} K$.

Composition of extensions

Let $H \leq K$, and let $H \leq K_i \leq K$, $i = 1, 2$

- (i) If $H \leq_{\text{alg}} K_1 \leq_{\text{alg}} K$ then $H \leq_{\text{alg}} K$.
- (i') If $H \leq_{\text{ff}} K_1 \leq_{\text{ff}} K$ then $H \leq_{\text{ff}} K$.
- (ii) If $H \leq_{\text{alg}} K$ then $K_1 \leq_{\text{alg}} K$, while $H \leq K_1$ need not be algebraic.

Composition of extensions

Let $H \leq K$, and let $H \leq K_i \leq K$, $i = 1, 2$

- (i) If $H \leq_{\text{alg}} K_1 \leq_{\text{alg}} K$ then $H \leq_{\text{alg}} K$.
- (i') If $H \leq_{\text{ff}} K_1 \leq_{\text{ff}} K$ then $H \leq_{\text{ff}} K$.
- (ii) If $H \leq_{\text{alg}} K$ then $K_1 \leq_{\text{alg}} K$, while $H \leq K_1$ need not be algebraic.
- (ii') If $H \leq_{\text{ff}} K$ then $H \leq_{\text{ff}} K_1$, while $K_1 \leq K$ need not be purely transcendental.

Composition of extensions

Let $H \leq K$, and let $H \leq K_i \leq K$, $i = 1, 2$

- (i) If $H \leq_{\text{alg}} K_1 \leq_{\text{alg}} K$ then $H \leq_{\text{alg}} K$.
- (i') If $H \leq_{\text{ff}} K_1 \leq_{\text{ff}} K$ then $H \leq_{\text{ff}} K$.
- (ii) If $H \leq_{\text{alg}} K$ then $K_1 \leq_{\text{alg}} K$, while $H \leq K_1$ need not be algebraic.
- (ii') If $H \leq_{\text{ff}} K$ then $H \leq_{\text{ff}} K_1$, while $K_1 \leq K$ need not be purely transcendental.
- (iii) If $H \leq_{\text{alg}} K_1$ and $H \leq_{\text{alg}} K_2$ then $H \leq_{\text{alg}} \langle K_1 \cup K_2 \rangle$, while $H \leq K_1 \cap K_2$ need not be algebraic.

Composition of extensions

Let $H \leq K$, and let $H \leq K_i \leq K$, $i = 1, 2$

- (i) If $H \leq_{\text{alg}} K_1 \leq_{\text{alg}} K$ then $H \leq_{\text{alg}} K$.
- (i') If $H \leq_{\text{ff}} K_1 \leq_{\text{ff}} K$ then $H \leq_{\text{ff}} K$.
- (ii) If $H \leq_{\text{alg}} K$ then $K_1 \leq_{\text{alg}} K$, while $H \leq K_1$ need not be algebraic.
- (ii') If $H \leq_{\text{ff}} K$ then $H \leq_{\text{ff}} K_1$, while $K_1 \leq K$ need not be purely transcendental.
- (iii) If $H \leq_{\text{alg}} K_1$ and $H \leq_{\text{alg}} K_2$ then $H \leq_{\text{alg}} \langle K_1 \cup K_2 \rangle$, while $H \leq K_1 \cap K_2$ need not be algebraic.
- (iii') If $H \leq_{\text{ff}} K_1$ and $H \leq_{\text{ff}} K_2$ then $H \leq_{\text{ff}} K_1 \cap K_2$, while $H \leq \langle K_1 \cup K_2 \rangle$ need not be purely transcendental.

The semilattice of algebraic extensions

- ▶ It follows that $AE(H)$ is a semilattice

The semilattice of algebraic extensions

- ▶ It follows that $AE(H)$ is a semilattice

More precisely, if $H \leq K$, there exists a unique L such that $H \leq_{\text{alg}} L \leq_{\text{ff}} K$, and L is

The semilattice of algebraic extensions

- ▶ It follows that $AE(H)$ is a semilattice

More precisely, if $H \leq K$, there exists a unique L such that $H \leq_{\text{alg}} L \leq_{\text{ff}} K$, and L is

- ▶ the intersection of the free factors of K containing H (and hence the least free factor of K containing H)

The semilattice of algebraic extensions

- ▶ It follows that $AE(H)$ is a semilattice

More precisely, if $H \leq K$, there exists a unique L such that $H \leq_{\text{alg}} L \leq_{\text{ff}} K$, and L is

- ▶ the intersection of the free factors of K containing H (and hence the least free factor of K containing H)
- ▶ the greatest algebraic extension of H in K

The semilattice of algebraic extensions

- ▶ It follows that $AE(H)$ is a semilattice

More precisely, if $H \leq K$, there exists a unique L such that $H \leq_{\text{alg}} L \leq_{\text{ff}} K$, and L is

- ▶ the intersection of the free factors of K containing H (and hence the least free factor of K containing H)
- ▶ the greatest algebraic extension of H in K
- ▶ the set of elements, that are K -algebraic over H

The semilattice of algebraic extensions

- ▶ It follows that $\text{AE}(H)$ is a semilattice

More precisely, if $H \leq K$, there exists a unique L such that $H \leq_{\text{alg}} L \leq_{\text{ff}} K$, and L is

- ▶ the intersection of the free factors of K containing H (and hence the least free factor of K containing H)
- ▶ the greatest algebraic extension of H in K
- ▶ the set of elements, that are K -algebraic over H

This extension L is called *the algebraic closure of H in K*

Elementary algebraic extensions

elementary extension $H \leq \langle H, x \rangle$

Elementary algebraic extensions

An elementary extension $H \leq \langle H, x \rangle$ is either algebraic or free, depending on the rank:

Elementary algebraic extensions

An elementary extension $H \leq \langle H, x \rangle$ is either algebraic or free, depending on the rank: $H \leq_{\text{ff}} \langle H, x \rangle$ iff $\text{rk}(\langle H, x \rangle) = \text{rk}(H) + 1$, and $H \leq_{\text{alg}} \langle H, x \rangle$ iff $\text{rk}(\langle H, x \rangle) \leq \text{rk}(H)$

Elementary algebraic extensions

An elementary extension $H \leq \langle H, x \rangle$ is either algebraic or free, depending on the rank: $H \leq_{\text{ff}} \langle H, x \rangle$ iff $\text{rk}(\langle H, x \rangle) = \text{rk}(H) + 1$, and $H \leq_{\text{alg}} \langle H, x \rangle$ iff $\text{rk}(\langle H, x \rangle) \leq \text{rk}(H)$

Say that the extension $H \leq K$ is elementary-algebraic, $H \leq_{\text{ealg}} K$, if it is a composition of algebraic elementary extensions

$$H \leq_{\text{alg}} H_1 \leq_{\text{alg}} \cdots \leq_{\text{alg}} H_k = K$$

Elementary algebraic extensions

An elementary extension $H \leq \langle H, x \rangle$ is either algebraic or free, depending on the rank: $H \leq_{\text{ff}} \langle H, x \rangle$ iff $\text{rk}(\langle H, x \rangle) = \text{rk}(H) + 1$, and $H \leq_{\text{alg}} \langle H, x \rangle$ iff $\text{rk}(\langle H, x \rangle) \leq \text{rk}(H)$

Say that the extension $H \leq K$ is elementary-algebraic, $H \leq_{\text{ealg}} K$, if it is a composition of algebraic elementary extensions

$$H \leq_{\text{alg}} H_1 \leq_{\text{alg}} \cdots \leq_{\text{alg}} H_k = K$$

▶ $H \leq_{\text{ealg}} K \implies \text{rk}(K) \leq \text{rk}(H)$

Elementary algebraic extensions

An elementary extension $H \leq \langle H, x \rangle$ is either algebraic or free, depending on the rank: $H \leq_{\text{ff}} \langle H, x \rangle$ iff $\text{rk}(\langle H, x \rangle) = \text{rk}(H) + 1$, and $H \leq_{\text{alg}} \langle H, x \rangle$ iff $\text{rk}(\langle H, x \rangle) \leq \text{rk}(H)$

Say that the extension $H \leq K$ is elementary-algebraic, $H \leq_{\text{ealg}} K$, if it is a composition of algebraic elementary extensions

$$H \leq_{\text{alg}} H_1 \leq_{\text{alg}} \cdots \leq_{\text{alg}} H_k = K$$

- ▶ $H \leq_{\text{ealg}} K \implies \text{rk}(K) \leq \text{rk}(H)$
- ▶ Not every algebraic extension is elementary algebraic; e.g. $\langle aba^{-1}b^{-1} \rangle \leq_{\text{alg}} F_2$ but $\text{rk}(\langle aba^{-1}b^{-1}, x \rangle) = 2$ for each $x \notin \langle aba^{-1}b^{-1} \rangle$

Elementary algebraic extensions

An elementary extension $H \leq \langle H, x \rangle$ is either algebraic or free, depending on the rank: $H \leq_{\text{ff}} \langle H, x \rangle$ iff $\text{rk}(\langle H, x \rangle) = \text{rk}(H) + 1$, and $H \leq_{\text{alg}} \langle H, x \rangle$ iff $\text{rk}(\langle H, x \rangle) \leq \text{rk}(H)$

Say that the extension $H \leq K$ is elementary-algebraic, $H \leq_{\text{ealg}} K$, if it is a composition of algebraic elementary extensions

$$H \leq_{\text{alg}} H_1 \leq_{\text{alg}} \cdots \leq_{\text{alg}} H_k = K$$

- ▶ $H \leq_{\text{ealg}} K \implies \text{rk}(K) \leq \text{rk}(H)$
- ▶ Not every algebraic extension is elementary algebraic; e.g. $\langle aba^{-1}b^{-1} \rangle \leq_{\text{alg}} F_2$ but $\text{rk}(\langle aba^{-1}b^{-1}, x \rangle) = 2$ for each $x \notin \langle aba^{-1}b^{-1} \rangle$
- ▶ Every $H \leq_{\text{fg}} F$ has an e-algebraic closure, which is computable

Subgroups of free groups

Congruent subgroups

Takahasi's theorem

Representing subgroups of free groups

A 3-line proof

Algebraic extensions of a subgroup

What they are

About terminology

Composition of extensions

The semilattice of algebraic extensions

Elementary algebraic extensions

Some applications

Malnormality, purity, etc

Profinite topologies

Equations

Abstract properties of subgroups

Let \mathcal{P} be a property of subgroups of a free group F , which is satisfied by F

Abstract properties of subgroups

Let \mathcal{P} be a property of subgroups of a free group F , which is satisfied by F

- ▶ If \mathcal{P} is closed under intersection, then every $H \leq F$ admits a \mathcal{P} -closure $\text{cl}_{\mathcal{P}}(H)$

Abstract properties of subgroups

Let \mathcal{P} be a property of subgroups of a free group F , which is satisfied by F

- ▶ If \mathcal{P} is closed under intersection, then every $H \leq F$ admits a \mathcal{P} -closure $\text{cl}_{\mathcal{P}}(H)$
- ▶ If \mathcal{P} is closed under finite intersection and under free factors, then every $H \leq_{\text{fg}} F$ admits a \mathcal{P} -closure, and $H \leq_{\text{alg}} \text{cl}_{\mathcal{P}}(H)$

Abstract properties of subgroups

Let \mathcal{P} be a property of subgroups of a free group F , which is satisfied by F

- ▶ If \mathcal{P} is closed under intersection, then every $H \leq F$ admits a \mathcal{P} -closure $\text{cl}_{\mathcal{P}}(H)$
- ▶ If \mathcal{P} is closed under finite intersection and under free factors, then every $H \leq_{\text{fg}} F$ admits a \mathcal{P} -closure, and $H \leq_{\text{alg}} \text{cl}_{\mathcal{P}}(H)$
- ▶ Proof: Suppose $H \leq K$ and K satisfies \mathcal{P}

Abstract properties of subgroups

Let \mathcal{P} be a property of subgroups of a free group F , which is satisfied by F

- ▶ If \mathcal{P} is closed under intersection, then every $H \leq F$ admits a \mathcal{P} -closure $\text{cl}_{\mathcal{P}}(H)$
- ▶ If \mathcal{P} is closed under finite intersection and under free factors, then every $H \leq_{\text{fg}} F$ admits a \mathcal{P} -closure, and $H \leq_{\text{alg}} \text{cl}_{\mathcal{P}}(H)$
- ▶ Proof: Suppose $H \leq K$ and K satisfies \mathcal{P}
- ▶ Then $H \leq L \leq_{\text{ff}} K$ for some $L \in \text{AE}(H)$

Abstract properties of subgroups

Let \mathcal{P} be a property of subgroups of a free group F , which is satisfied by F

- ▶ If \mathcal{P} is closed under intersection, then every $H \leq F$ admits a \mathcal{P} -closure $\text{cl}_{\mathcal{P}}(H)$
- ▶ If \mathcal{P} is closed under finite intersection and under free factors, then every $H \leq_{\text{fg}} F$ admits a \mathcal{P} -closure, and $H \leq_{\text{alg}} \text{cl}_{\mathcal{P}}(H)$
- ▶ Proof: Suppose $H \leq K$ and K satisfies \mathcal{P}
- ▶ Then $H \leq L \leq_{\text{ff}} K$ for some $L \in \text{AE}(H)$
- ▶ It follows that L satisfies \mathcal{P} , and the intersection of the K containing H and satisfying \mathcal{P} is the finite intersection of the algebraic extensions of H satisfying \mathcal{P}

Abstract properties of subgroups

Let \mathcal{P} be a property of subgroups of a free group F , which is satisfied by F

- ▶ If \mathcal{P} is closed under intersection, then every $H \leq F$ admits a \mathcal{P} -closure $\text{cl}_{\mathcal{P}}(H)$
- ▶ If \mathcal{P} is closed under finite intersection and under free factors, then every $H \leq_{\text{fg}} F$ admits a \mathcal{P} -closure, and $H \leq_{\text{alg}} \text{cl}_{\mathcal{P}}(H)$
- ▶ If in addition \mathcal{P} is decidable, then $\text{cl}_{\mathcal{P}}(H)$ is computable

Examples

Say that H is *malnormal* if $H^g \cap H = 1$ for every $g \in F \setminus H$.

Examples

Say that H is *malnormal* if $H^g \cap H = 1$ for every $g \in F \setminus H$.
Malnormality is closed under intersections and free products,
decidable, and hence

Examples

Say that H is *malnormal* if $H^g \cap H = 1$ for every $g \in F \setminus H$.

Malnormality is closed under intersections and free products, decidable, and hence

every $H \leq_{\text{fg}} F$ admits a malnormal closure, which is an algebraic extension, and is computable

Examples

Say that H is *malnormal* if $H^g \cap H = 1$ for every $g \in F \setminus H$.

Malnormality is closed under intersections and free products, decidable, and hence

every $H \leq_{\text{fg}} F$ admits a malnormal closure, which is an algebraic extension, and is computable

The same holds for

Examples

Say that H is *malnormal* if $H^g \cap H = 1$ for every $g \in F \setminus H$.

Malnormality is closed under intersections and free products, decidable, and hence

every $H \leq_{\text{fg}} F$ admits a malnormal closure, which is an algebraic extension, and is computable

The same holds for

- ▶ the property of being pure: H is pure if $x^n \in H$ and $n \neq 0$ implies $x \in H$

Examples

Say that H is *malnormal* if $H^g \cap H = 1$ for every $g \in F \setminus H$.

Malnormality is closed under intersections and free products, decidable, and hence

every $H \leq_{\text{fg}} F$ admits a malnormal closure, which is an algebraic extension, and is computable

The same holds for

- ▶ the property of being pure: H is pure if $x^n \in H$ and $n \neq 0$ implies $x \in H$
- ▶ the property of being p -pure: H is p -pure if $x^n \in H$, $n \neq 0$ and $(n, p) = 1$ implies $x \in H$

Examples

Say that H is *malnormal* if $H^g \cap H = 1$ for every $g \in F \setminus H$.

Malnormality is closed under intersections and free products, decidable, and hence

every $H \leq_{\text{fg}} F$ admits a malnormal closure, which is an algebraic extension, and is computable

The same holds for

- ▶ the property of being pure: H is pure if $x^n \in H$ and $n \neq 0$ implies $x \in H$
- ▶ the property of being p -pure: H is p -pure if $x^n \in H$, $n \neq 0$ and $(n, p) = 1$ implies $x \in H$
- ▶ the property of being a retract

Closure of a subgroup in a profinite topology

If V is a pseudovariety of finite groups which is extension-closed, the property of being closed in the pro- V topology of the free group is closed under intersection and free factors

Closure of a subgroup in a profinite topology

If V is a pseudovariety of finite groups which is extension-closed, the property of being closed in the pro- V topology of the free group is closed under intersection and free factors
It follows that the pro- V closure of a finitely generated subgroup H is an algebraic extension of H (Margolis, Sapir, W), and in particular that it has finite rank (Ribes, Zaleskiĭ)

Closure of a subgroup in a profinite topology

If V is a pseudovariety of finite groups which is extension-closed, the property of being closed in the pro- V topology of the free group is closed under intersection and free factors
It follows that the pro- V closure of a finitely generated subgroup H is an algebraic extension of H (Margolis, Sapir, W), and in particular that it has finite rank (Ribes, Zalesskiĭ)

- ▶ However, it is not always computable

Closure of a subgroup in a profinite topology

If V is a pseudovariety of finite groups which is extension-closed, the property of being closed in the pro- V topology of the free group is closed under intersection and free factors
It follows that the pro- V closure of a finitely generated subgroup H is an algebraic extension of H (Margolis, Sapir, W), and in particular that it has finite rank (Ribes, Zalesskiĭ)

- ▶ However, it is not always computable
- ▶ Note that $\text{rk}(\text{cl}_V(H)) \leq \text{rk}(H)$, but it is not known whether this is because $H \leq_{\text{ealg}} \text{cl}_V(H)$

Equations over subgroups

- ▶ Let $H \leq F$. An H -equation is a word $e \in H * \langle X \rangle$, where X is a new symbol (the unknown). A solution of H is an element x such that $e(x) = 1$

Equations over subgroups

- ▶ Let $H \leq F$. An H -equation is a word $e \in H * \langle X \rangle$, where X is a new symbol (the unknown). A solution of H is an element x such that $e(x) = 1$
- ▶ Thus x is a solution of some H -equation if and only if the homomorphism $H * \langle X \rangle \rightarrow \langle H, x \rangle$ is not one-to-one, if and only if $\langle H, x \rangle$ has rank at most $\text{rk}(H)$, if and only if $H \leq_{\text{alg}} \langle H, x \rangle$

Equations over subgroups

- ▶ Let $H \leq F$. An H -equation is a word $e \in H * \langle X \rangle$, where X is a new symbol (the unknown). A solution of H is an element x such that $e(x) = 1$
- ▶ Thus x is a solution of some H -equation if and only if the homomorphism $H * \langle X \rangle \rightarrow \langle H, x \rangle$ is not one-to-one, if and only if $\langle H, x \rangle$ has rank at most $\text{rk}(H)$, if and only if $H \leq_{\text{alg}} \langle H, x \rangle$
- ▶ an equation is an element e of the countably generated free group $F(X, Y_1, \dots, Y_n, \dots)$

Equations over subgroups

- ▶ Let $H \leq F$. An H -equation is a word $e \in H * \langle X \rangle$, where X is a new symbol (the unknown). A solution of H is an element x such that $e(x) = 1$
- ▶ Thus x is a solution of some H -equation if and only if the homomorphism $H * \langle X \rangle \rightarrow \langle H, x \rangle$ is not one-to-one, if and only if $\langle H, x \rangle$ has rank at most $\text{rk}(H)$, if and only if $H \leq_{\text{alg}} \langle H, x \rangle$
- ▶ an equation is an element e of the countably generated free group $F(X, Y_1, \dots, Y_n, \dots)$
- ▶ Let \mathcal{E} be a set of equations: say that H is \mathcal{E} -closed if H contains every solution of every particularization over H of an equation in \mathcal{E} .

Examples

- ▶ Let $\mathcal{E}_{mal} = \{X^{-1}Y_1XY_2\}$. A solution of a particularization of this equation over H is an element x such that $H^x \cap H \neq 1$,

Examples

- ▶ Let $\mathcal{E}_{mal} = \{X^{-1}Y_1XY_2\}$. A solution of a particularization of this equation over H is an element x such that $H^x \cap H \neq 1$,
- ▶ so H is \mathcal{E}_{mal} -closed if and only if H is malnormal.

Examples

- ▶ Let $\mathcal{E}_{mal} = \{X^{-1}Y_1XY_2\}$. A solution of a particularization of this equation over H is an element x such that $H^x \cap H \neq 1$,
- ▶ so H is \mathcal{E}_{mal} -closed if and only if H is malnormal.
- ▶ Similarly, let $\mathcal{E}_{pure} = \{X^nY_1 \mid n \geq 2\}$. Then H is \mathcal{E}_{pure} -closed if and only if H is pure.

Let \mathcal{E} be a set of equations

- ▶ The property of being \mathcal{E} -closed is closed under intersections and free products,

Let \mathcal{E} be a set of equations

- ▶ The property of being \mathcal{E} -closed is closed under intersections and free products,
- ▶ so H admits an \mathcal{E} -closure, and $H \leq_{\text{alg}} \text{cl}_{\mathcal{E}}(H)$

Let \mathcal{E} be a set of equations

- ▶ The property of being \mathcal{E} -closed is closed under intersections and free products,
- ▶ so H admits an \mathcal{E} -closure, and $H \leq_{\text{alg}} \text{cl}_{\mathcal{E}}(H)$
- ▶ In addition, $H \leq_{\text{ealg}} \text{cl}_{\mathcal{E}}(H)$, so $\text{rk}(\text{cl}_{\mathcal{E}}(H)) \leq \text{rk}(H)$

Let \mathcal{E} be a set of equations

- ▶ The property of being \mathcal{E} -closed is closed under intersections and free products,
- ▶ so H admits an \mathcal{E} -closure, and $H \leq_{\text{alg}} \text{cl}_{\mathcal{E}}(H)$
- ▶ In addition, $H \leq_{\text{ealg}} \text{cl}_{\mathcal{E}}(H)$, so $\text{rk}(\text{cl}_{\mathcal{E}}(H)) \leq \text{rk}(H)$

Thus the malnormal closure of $H \leq_{\text{fg}} F$ has rank at most $\text{rk}(H)$, as well as the pure closure. . .