

ASEP with open boundaries and Koornwinder polynomials


Luigi Cantini

Journées de combinatoire de Bordeaux
27 January 2016



Introduction


Particles propagating under the effect of an **external field** and interacting with two **reservoirs** at different chemical potential ρ_L, ρ_R



No detailed balance: **Macroscopic particle current**


Introduction

Particles propagating under the effect of an **external field** and interacting with two **reservoirs** at different chemical potential ρ_L, ρ_R






No detailed balance: **Macroscopic particle current**

The Asymmetric Simple Exclusion Process (ASEP)




- ▶ One dimensional lattice
- ▶ **Exclusion:** at most one particle per site
- ▶ **Asymmetric:** jump rate to the right $t^{1/2}$, to the left $t^{-1/2}$
- ▶ Particles enter with rate α from left, with rate δ from right
- ▶ Particles leave with rate γ from left, with rate β from right

Applications




Impurities

Here we want to take into account the presence of impurities that are **NOT EXCHANGED** with the reservoirs



Two species ASEP



The impurity (*second-class particle*) has the same dynamics as a normal (*first-class particle*), while the first-class particles treat it as a hole.

It is convenient to think at first class, second class particles and empty sites as *three kinds of particles*



Master equation and Markov generator

Probabilities of the configurations $\{w\}$ evolve under a Master equation

$$\frac{d}{dt}P_w(t) = \sum_{w' \neq w} \mathcal{M}(w' \rightarrow w)P_{w'}(t) - \sum_{w' \neq w} \mathcal{M}(w \rightarrow w')P_w(t)$$

Using a vector representation for the probabilities

$$\mathcal{P}_{N,m}(t) = \sum_{w \in \mathcal{Q}(N,m)} P_w(t)$$

Number of impurities

Length of the chain

The Master equation reads

$$\frac{d}{dt}\mathcal{P}(t) = \mathcal{M}\mathcal{P}(t)$$

Where the Markov generator is given by the sum of local terms

$$\mathcal{M} = \sum_{i=1}^{N-1} \mathbf{e}_i + \mathbf{f}_1 + \mathbf{f}_N.$$

Master equation and Markov generator

Probabilities of the configurations $\{w\}$ evolve under a Master equation

$$\frac{d}{dt}P_w(t) = \sum_{w' \neq w} \mathcal{M}(w' \rightarrow w)P_{w'}(t) - \sum_{w' \neq w} \mathcal{M}(w \rightarrow w')P_w(t)$$

Using a vector representation for the probabilities

$$\mathcal{P}_{N,m}(t) = \sum_{w \in \mathcal{Q}(N,m)} P_w(t)w$$

Number of impurities

Length of the chain

The Master equation reads

$$\frac{d}{dt}\mathcal{P}(t) = \mathcal{M}\mathcal{P}(t)$$

Where the Markov generator is given by the sum of local terms

$$\mathcal{M} = \sum_{i=1}^{N-1} \mathbf{e}_i + \mathbf{f}_1 + \mathbf{f}_N.$$

Master equation and Markov generator

Probabilities of the configurations $\{w\}$ evolve under a Master equation

$$\frac{d}{dt}P_w(t) = \sum_{w' \neq w} \mathcal{M}(w' \rightarrow w)P_{w'}(t) - \sum_{w' \neq w} \mathcal{M}(w \rightarrow w')P_w(t)$$

Using a vector representation for the probabilities

$$\mathcal{P}_{N,m}(t) = \sum_{w \in \mathcal{Q}(N,m)} P_w(t)w$$

Number of impurities

Length of the chain

The Master equation reads

$$\frac{d}{dt}\mathcal{P}(t) = \mathcal{M}\mathcal{P}(t)$$

Where the Markov generator is given by the sum of local terms

$$\mathcal{M} = \sum_{i=1}^{N-1} \mathbf{e}_i + \mathbf{f}_1 + \mathbf{f}_N.$$

Master equation and Markov generator

Probabilities of the configurations $\{w\}$ evolve under a Master equation

$$\frac{d}{dt}P_w(t) = \sum_{w' \neq w} \mathcal{M}(w' \rightarrow w)P_{w'}(t) - \sum_{w' \neq w} \mathcal{M}(w \rightarrow w')P_w(t)$$

Using a vector representation for the probabilities

$$\mathcal{P}_{N,m}(t) = \sum_{w \in \mathcal{Q}(N,m)} P_w(t)w$$

Number of impurities

Length of the chain

The Master equation reads

$$\frac{d}{dt}\mathcal{P}(t) = \mathcal{M}\mathcal{P}(t)$$

Where the Markov generator is given by the sum of local terms

$$\mathcal{M} = \sum_{i=1}^{N-1} \mathbf{e}_i + \mathbf{f}_1 + \mathbf{f}_N.$$

Master equation and Markov generator

Probabilities of the configurations $\{w\}$ evolve under a Master equation

$$\frac{d}{dt}P_w(t) = \sum_{w' \neq w} \mathcal{M}(w' \rightarrow w)P_{w'}(t) - \sum_{w' \neq w} \mathcal{M}(w \rightarrow w')P_w(t)$$

Using a vector representation for the probabilities

$$\mathcal{P}_{N,m}(t) = \sum_{w \in \mathcal{Q}(N,m)} P_w(t)w$$

Number of impurities

Length of the chain

The Master equation reads

$$\frac{d}{dt}\mathcal{P}(t) = \mathcal{M}\mathcal{P}(t)$$

Where the Markov generator is given by the sum of local terms

$$\mathcal{M} = \sum_{i=1}^{N-1} \mathbf{e}_i + \mathbf{f}_1 + \mathbf{f}_N.$$

Stationary measure

In this talk I'll focus on the *Stationary measure*

$$\mathcal{MP}_{N,m} = 0$$

- ▶ Usually dealt with by the Matrix Product Ansatz [Derrida, Evans, Hakim, Pasquier]
- ▶ Rich combinatorics: rhombic alternative and rhombic staircase tableaux [Corteel, Williams, Mandelshtam, Viennot, ...]
- ▶ Boundary induced phase transitions [Krug, ...]
- ▶ Here I'll describe an algebraic approach based on:

Exchange/reflection equations.

Stationary measure

In this talk I'll focus on the *Stationary measure*

$$\mathcal{M}P_{N,m} = 0$$

- ▶ Usually dealt with by the Matrix Product Ansatz [Derrida, Evans, Hakim, Pasquier]
- ▶ Rich combinatorics: rhombic alternative and rhombic staircase tableaux [Corteel, Williams, Mandelshtam, Viennot, ...]
- ▶ Boundary induced phase transitions [Krug, ...]
- ▶ Here I'll describe an algebraic approach based on:

Exchange/reflection equations.

Stationary measure

In this talk I'll focus on the *Stationary measure*

$$\mathcal{MP}_{N,m} = 0$$

- ▶ Usually dealt with by the Matrix Product Ansatz [Derrida, Evans, Hakim, Pasquier]
- ▶ Rich combinatorics: rhombic alternative and rhombic staircase tableaux [Corteel, Williams, Mandelshtam, Viennot, ...]
- ▶ Boundary induced phase transitions [Krug, ...]
- ▶ Here I'll describe an algebraic approach based on:

Exchange/reflection equations.

\hat{C}_N Hecke Algebra

Define operators T_0, T_1, \dots, T_N as

$$T_0 = \alpha^{-\frac{1}{2}} \delta^{-\frac{1}{2}} \mathbf{f}_1 + \alpha^{\frac{1}{2}} \delta^{-\frac{1}{2}} \mathbf{1}$$

$$T_N = \beta^{-\frac{1}{2}} \gamma^{-\frac{1}{2}} \mathbf{f}_N + \beta^{\frac{1}{2}} \gamma^{-\frac{1}{2}} \mathbf{1}$$

and for $1 \leq i \leq N-1$

$$T_i = \mathbf{e}_i + t^{-\frac{1}{2}} \mathbf{1}$$

They satisfy the commutation relations of the generators of the affine Hecke algebra \hat{C}_N

$$T_i - T_i^{-1} = t_i^{\frac{1}{2}} - t_i^{-\frac{1}{2}}$$

$$T_i T_j = T_j T_i \quad \text{if } |i-j| > 1$$

$$T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1} \quad \text{if } i \neq 0, N-1$$

$$T_0 T_1 T_0 T_1 = T_1 T_0 T_1 T_0$$

$$T_N T_{N-1} T_N T_{N-1} = T_{N-1} T_N T_{N-1} T_N$$

with $t_0^{\frac{1}{2}} = \alpha^{\frac{1}{2}} \delta^{-\frac{1}{2}}$, $t_N^{\frac{1}{2}} = \beta^{\frac{1}{2}} \gamma^{-\frac{1}{2}}$ and $t_i = t$ for $1 \leq i \leq N-1$.

Integrability

The \check{R} matrices

$$\check{R}_i(z) = \mathbf{1} + \frac{z-1}{t^{\frac{1}{2}}z - t^{-\frac{1}{2}}} \mathbf{e}_i.$$

The boundary scattering matrices

$$K_1(z|a, b) = \mathbf{1} + \frac{(z^2 - 1)}{(z-a)(z-b)} \delta^{-1} \mathbf{f}_1(\delta, \alpha),$$

$$K_N(z|c, d) = \mathbf{1} + \frac{(1 - z^2)}{(cz-1)(dz-1)} \gamma^{-1} \mathbf{f}_N(\beta, \gamma).$$

Where we assume $t \neq 1$ and parametrize the boundary rates as

$$\alpha = \frac{(t^{\frac{1}{2}} - t^{-\frac{1}{2}})ab}{(a-1)(b-1)}, \quad \delta = \frac{t^{-\frac{1}{2}} - t^{\frac{1}{2}}}{(a-1)(b-1)}$$
$$\beta = \frac{(t^{\frac{1}{2}} - t^{-\frac{1}{2}})cd}{(c-1)(d-1)}, \quad \gamma = \frac{t^{-\frac{1}{2}} - t^{\frac{1}{2}}}{(c-1)(d-1)}$$

Integrability

The \check{R} matrices

$$\check{R}_i(z) = \mathbf{1} + \frac{z-1}{t^{\frac{1}{2}}z - t^{-\frac{1}{2}}} \mathbf{e}_i.$$

The boundary scattering matrices

$$K_1(z|a, b) = \mathbf{1} + \frac{(z^2 - 1)}{(z - a)(z - b)} \delta^{-1} \mathbf{f}_1(\delta, \alpha),$$

$$K_N(z|c, d) = \mathbf{1} + \frac{(1 - z^2)}{(cz - 1)(dz - 1)} \gamma^{-1} \mathbf{f}_N(\beta, \gamma).$$

Where we assume $t \neq 1$ and parametrize the boundary rates as

$$\alpha = \frac{(t^{\frac{1}{2}} - t^{-\frac{1}{2}})ab}{(a-1)(b-1)}, \quad \delta = \frac{t^{-\frac{1}{2}} - t^{\frac{1}{2}}}{(a-1)(b-1)}$$
$$\beta = \frac{(t^{\frac{1}{2}} - t^{-\frac{1}{2}})cd}{(c-1)(d-1)}, \quad \gamma = \frac{t^{-\frac{1}{2}} - t^{\frac{1}{2}}}{(c-1)(d-1)}$$

Stationary measure: exchange/reflection equations

We introduce a vector

$$\Psi_{N,m}(\mathbf{z}) = \sum_{w \in \mathcal{Q}(N,m)} \psi_w(\mathbf{z}) w$$

solution of the following exchange/reflection equations

$$\check{R}_i(z_i z_{i+1}^{-1}) \Psi_{N,m}(\mathbf{z}) = \mathbf{s}_i \Psi_{N,m}(\mathbf{z})$$

$$K_1(z_1) \Psi_{N,m}(\mathbf{z}) = \mathbf{s}_0 \Psi_{N,m}(\mathbf{z})$$

$$K_N(z_N) \Psi_{N,m}(\mathbf{z}) = \mathbf{s}_N \Psi_{N,m}(\mathbf{z}).$$

where

$$\mathbf{s}_i f(\dots, z_i, z_{i+1}, \dots) = f(\dots, z_{i+1}, z_i, \dots)$$

$$\mathbf{s}_0 f(z_1, \dots) = f(z_1^{-1}, \dots)$$

$$\mathbf{s}_N f(\dots, z_N) = f(\dots, z_N^{-1})$$

Stationary measure: exchange/reflection equations

Consistency of the exchange/reflection equations is ensured by the Yang-Baxter equations

$$\check{R}_i(yz^{-1})\check{R}_{i+1}(xz^{-1})\check{R}_i(xy^{-1}) = \check{R}_{i+1}(xy^{-1})\check{R}_i(xz^{-1})\check{R}_{i+1}(yz^{-1})$$

$$\begin{aligned} \check{R}_1(xy^{-1})K_1(y)\check{R}_1(x^{-1}y^{-1})K_1(x) = \\ K_1(x)\check{R}_1(x^{-1}y^{-1})K_1(y)\check{R}_1(xy^{-1}), \end{aligned}$$

$$\begin{aligned} \check{R}_{N-1}(xy^{-1})K_N(x)\check{R}_{N-1}(xy)K_N(y) = \\ K_N(y)\check{R}_{N-1}(xy)K_N(x)\check{R}_{N-1}(xy^{-1}), \end{aligned}$$

and the so called unitarity condition

$$\check{R}_i(z)\check{R}_i(z^{-1}) = \mathbf{1}.$$

$$K_1(x)K_1(x^{-1}) = \mathbf{1}, \quad K_N(x)K_N(x^{-1}) = \mathbf{1}.$$

Stationary measure: exchange/reflection equations

Theorem

- ▶ The exchange equations have unique solution (up to multiplication by a function invariant under the action of $\mathbf{s}_0, \mathbf{s}_i, \mathbf{s}_N$)
- ▶ Under specialization $z_i = 1$ we get

$$\Psi_{N,m}(\mathbf{1}) \propto \mathcal{P}_{N,m}$$

Stationary measure: exchange/reflection equations

Theorem

- ▶ The exchange equations have unique solution (up to multiplication by a function invariant under the action of $\mathbf{s}_0, \mathbf{s}_i, \mathbf{s}_N$)
- ▶ Under specialization $z_i = 1$ we get

$$\Psi_{N,m}(\mathbf{1}) \propto \mathcal{P}_{N,m}$$

Exchange/reflection equations in components

$$\psi_{\dots, \underbrace{\bullet, \bullet}_{i, i+1}, \dots}(\mathbf{z}) = \mathbf{s}_i \psi_{\dots, \underbrace{\bullet, \bullet}_{i, i+1}, \dots}(\mathbf{z})$$

$$\psi_{*, \dots}(\mathbf{z}) = \mathbf{s}_0 \psi_{*, \dots}(\mathbf{z})$$

$$\psi_{\dots, *}(\mathbf{z}) = \mathbf{s}_N \psi_{\dots, *}(\mathbf{z})$$

$$\psi_{\dots, \underbrace{\bullet, \circ}_{i, i+1}, \dots}(\mathbf{z}) = t^{\frac{1}{2}} \widehat{T}_i \psi_{\dots, \underbrace{\circ, \bullet}_{i, i+1}, \dots}(\mathbf{z})$$

$$\psi_{\bullet, \dots}(\mathbf{z}) = t_0^{\frac{1}{2}} \widehat{T}_0 \psi_{\circ, \dots}(\mathbf{z})$$

$$\psi_{\dots, \bullet}(\mathbf{z}) = t_N^{\frac{1}{2}} \widehat{T}_N \psi_{\dots, \circ}(\mathbf{z})$$

Affine Hecke Again: Noumi representation

Noumi introduced a representation of $\widehat{\mathcal{C}}_N$ depending on 6 parameters a, b, c, d, t, q , acting on $\mathbb{C}[z_1^{\pm 1}, \dots, z_N^{\pm 1}]$

$$\widehat{T}_i = t^{\frac{1}{2}} - (t^{\frac{1}{2}} z_i - t^{-\frac{1}{2}} z_{i+1}) \partial_i \quad (1)$$

$$\widehat{T}_0 = t_0^{\frac{1}{2}} - t_0^{-\frac{1}{2}} \frac{(z_1 - a)(z_1 - b)}{z_1} \partial_0 \quad (2)$$

$$\widehat{T}_N = t_N^{\frac{1}{2}} - t_N^{-\frac{1}{2}} \frac{(cz_N - 1)(dz_N - 1)}{z_N} \partial_N, \quad (3)$$

where $t_0 = -q^{-1}ab$ $t_N = -cd$ and

$$\partial_i = \frac{1 - \mathbf{s}_i}{z_i - z_{i+1}}, \quad \partial_0 = \frac{1 - \mathbf{s}_0}{z_1 - qz_1^{-1}}, \quad \partial_N = \frac{1 - \mathbf{s}_N}{z_N - z_N^{-1}}.$$

Where $\mathbf{s}_i, \mathbf{s}_N$ are as before but

$$\mathbf{s}_0 f(z_1, \dots) = f(qz_1^{-1}, \dots)$$

For ASEP $q = 1$

Affine Hecke Again: Noumi representation

Noumi introduced a representation of $\widehat{\mathcal{C}}_N$ depending on 6 parameters a, b, c, d, t, q , acting on $\mathbb{C}[z_1^{\pm 1}, \dots, z_N^{\pm 1}]$

$$\widehat{T}_i = t^{\frac{1}{2}} - (t^{\frac{1}{2}} z_i - t^{-\frac{1}{2}} z_{i+1}) \partial_i \quad (1)$$

$$\widehat{T}_0 = t_0^{\frac{1}{2}} - t_0^{-\frac{1}{2}} \frac{(z_1 - a)(z_1 - b)}{z_1} \partial_0 \quad (2)$$

$$\widehat{T}_N = t_N^{\frac{1}{2}} - t_N^{-\frac{1}{2}} \frac{(cz_N - 1)(dz_N - 1)}{z_N} \partial_N, \quad (3)$$

where $t_0 = -q^{-1}ab$ $t_N = -cd$ and

$$\partial_i = \frac{1 - \mathbf{s}_i}{z_i - z_{i+1}}, \quad \partial_0 = \frac{1 - \mathbf{s}_0}{z_1 - qz_1^{-1}}, \quad \partial_N = \frac{1 - \mathbf{s}_N}{z_N - z_N^{-1}}.$$

Where $\mathbf{s}_i, \mathbf{s}_N$ are as before but

$$\mathbf{s}_0 f(z_1, \dots) = f(qz_1^{-1}, \dots)$$

For ASEP $q = 1$

The commutative sub-algebra \mathcal{Y}_N generated by elements $Y_1^{\pm 1}, \dots, Y_N^{\pm 1}$
[Lusztig]

$$Y_i = (T_i \dots T_{N-1})(T_N \dots T_0)(T_1^{-1} \dots T_{i-1}^{-1}).$$

Its common eigenfunctions are the *non-symmetric Koornwinder polynomials* $E_\alpha(\mathbf{z})$ ($\alpha \in \mathbb{Z}^N$)

$$E_\alpha(\mathbf{z}) = \mathbf{z}^\alpha + \sum_{\mathbf{z}^\beta \prec \mathbf{z}^\alpha} c_\beta \mathbf{z}^\beta,$$

$$\hat{Y}_i E_\alpha(\mathbf{z}) = \omega_i(\alpha) E_\alpha(\mathbf{z}).$$

By using the exchange/reflection relations it is easy to show that

$$\psi_{\underbrace{\circ \dots \circ}_{N-m} \underbrace{* \dots *}_m}(\mathbf{z}) = E_{\underbrace{-1 \dots -1}_{N-m} \underbrace{0 \dots 0}_m}(\mathbf{z})$$

(Actually $E_{\underbrace{-1 \dots -1}_{N-m} \underbrace{0 \dots 0}_m}(\mathbf{z})$ doesn't depend on q).

The commutative sub-algebra \mathcal{Y}_N generated by elements $Y_1^{\pm 1}, \dots, Y_N^{\pm 1}$
[Lusztig]

$$Y_i = (T_i \dots T_{N-1})(T_N \dots T_0)(T_1^{-1} \dots T_{i-1}^{-1}).$$

Its common eigenfunctions are the *non-symmetric Koornwinder polynomials* $E_\alpha(\mathbf{z})$ ($\alpha \in \mathbb{Z}^N$)

$$E_\alpha(\mathbf{z}) = \mathbf{z}^\alpha + \sum_{\mathbf{z}^\beta \prec \mathbf{z}^\alpha} c_\beta \mathbf{z}^\beta,$$

$$\hat{Y}_i E_\alpha(\mathbf{z}) = \omega_i(\alpha) E_\alpha(\mathbf{z}).$$

By using the exchange/reflection relations it is easy to show that

$$\psi_{\underbrace{\circ \dots \circ}_{N-m} \underbrace{* \dots *}_m}(\mathbf{z}) = E_{\underbrace{-1 \dots -1}_{N-m} \underbrace{0 \dots 0}_m}(\mathbf{z})$$

(Actually $E_{\underbrace{-1 \dots -1}_{N-m} \underbrace{0 \dots 0}_m}(\mathbf{z})$ doesn't depend on q).

Symmetric Macdonald-Koornwinder polynomials [Koornwinder]

Koornwinder q -difference operator

$$\mathcal{D}_{q,t} = \sum_{i=1}^N \Phi_i(z_i)(T_{q,z_i} - 1) + \Phi_i(z_i^{-1})(T_{q,z_i}^{-1} - 1)$$

where T_{q,z_i} is the i -th q -shift operator

$$T_{q,z_i} f(z_1, \dots, z_i, \dots, z_N) = f(z_1, \dots, qz_i, \dots, z_N)$$

and

$$\Phi_i(z) = \frac{(1-az)(1-bz)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)} \prod_{\substack{j=1 \\ j \neq i}}^N \frac{(1-tzz_j)(1-tzz_j^{-1})}{(1-zz_j)(1-zz_j^{-1})}$$

Symmetric Macdonald-Koornwinder polynomials

The symmetric Macdonald-Koornwinder polynomials $P_\lambda(\mathbf{z})$

- ▶ Laurent polynomials in N variables
- ▶ Labeled by a partition λ , coefficient of \mathbf{z}^λ in $P_\lambda(\mathbf{z})$ is 1
- ▶ Eigenfunctions of $\mathcal{D}_{q,t}$

$$\mathcal{D}_{q,t}P_\lambda(\mathbf{z}) = d_\lambda P_\lambda(\mathbf{z})$$

$$d_\lambda = \sum_{i=1}^N [q^{-1}abcdt^{2n-i-1}(q^{\lambda_i} - 1) + t^{i-1}(q^{-\lambda_i} - 1)]$$

- ▶ They are multivariables generalization of the Askey-Wilson polynomials

$$P_{\{m\}}(\mathbf{z}) \propto p_m(x; a, b, c, d|q).$$

Symmetric Macdonald-Koornwinder polynomials

The symmetric Macdonald-Koornwinder polynomials $P_\lambda(\mathbf{z})$

- ▶ Laurent polynomials in N variables
- ▶ Labeled by a partition λ , coefficient of \mathbf{z}^λ in $P_\lambda(\mathbf{z})$ is 1
- ▶ Eigenfunctions of $\mathcal{D}_{q,t}$

$$\mathcal{D}_{q,t}P_\lambda(\mathbf{z}) = d_\lambda P_\lambda(\mathbf{z})$$

$$d_\lambda = \sum_{i=1}^N [q^{-1}abcdt^{2n-i-1}(q^{\lambda_i} - 1) + t^{i-1}(q^{-\lambda_i} - 1)]$$

- ▶ They are multivariables generalization of the Askey-Wilson polynomials

$$P_{\{m\}}(\mathbf{z}) \propto p_m(x; a, b, c, d|q).$$

Symmetric Macdonald-Koornwinder polynomials

The symmetric Macdonald-Koornwinder polynomials $P_\lambda(\mathbf{z})$

- ▶ Laurent polynomials in N variables
- ▶ Labeled by a partition λ , coefficient of \mathbf{z}^λ in $P_\lambda(\mathbf{z})$ is 1
- ▶ Eigenfunctions of $\mathcal{D}_{q,t}$

$$\mathcal{D}_{q,t}P_\lambda(\mathbf{z}) = d_\lambda P_\lambda(\mathbf{z})$$

$$d_\lambda = \sum_{i=1}^N [q^{-1}abcdt^{2n-i-1}(q^{\lambda_i} - 1) + t^{i-1}(q^{-\lambda_i} - 1)]$$

- ▶ They are multivariables generalization of the Askey-Wilson polynomials

$$P_{\{m\}}(\mathbf{z}) \propto p_m(x; a, b, c, d|q).$$

Symmetric Macdonald-Koornwinder polynomials

► Orthogonality

$$\langle P_\lambda, P_\mu \rangle = 0 \quad \text{if} \quad \lambda \neq \mu.$$

$$\langle p, q \rangle := \oint_{\mathcal{C}} \prod_{j=1}^N \frac{dz_j}{4\pi iz_j} \Delta_N(t, q; \mathbf{z}) p(\mathbf{z}) q(\mathbf{z}^{-1})$$

where the contour of integration \mathcal{C} encircles the poles in aq^k, bq^k, cq^k, dq^k ($k \in \mathbb{Z}_+$) and excludes all the others.

$$\Delta_N(t, q; \mathbf{z}) = \prod_{\substack{1 \leq i < j \leq N \\ \epsilon_1, \epsilon_2 = \pm 1}} \frac{(z_i^{\epsilon_1} z_j^{\epsilon_2}; q)_\infty}{(tz_i^{\epsilon_1} z_j^{\epsilon_2}; q)_\infty} \prod_{i=1}^N \frac{(z_i^{2\epsilon}; q)_\infty}{(az_i^\epsilon, bz_i^\epsilon, cz_i^\epsilon, dz_i^\epsilon; q)_\infty},$$

Weighted partition function

Let $\bullet(w)$ be the number of first class particles in configuration w . We define the *weighted partition function* as

$$Z_N(\xi; \mathbf{z}; a, b, c, d) := \sum_{w \in \mathcal{Q}(N, m)} \xi^{2 \bullet(w)} \psi_w(\mathbf{z}).$$

Theorem

The generating function for the number of particles in the system is given by

$$Z_{N, m}(\xi; \mathbf{z}; a, b, c, d) = \xi^{N-m} P_{1^{N-m} 0^m}(\xi \mathbf{z} | \xi \mathbf{z}; a_\xi, b_\xi, c_\xi, d_\xi)$$

with $a_\xi = \xi a$, $b_\xi = \xi b$, $c_\xi = \xi^{-1} c$, $d_\xi = \xi^{-1} d$.

Remark: the Macdonald-Koornwinder polynomials $P_{1^{N-m} 0^m}(\mathbf{z})$ associated to single column diagrams $\lambda = 1^{N-m} 0^m$ are independent of the parameter q .

Weighted partition function

Let $\bullet(w)$ be the number of first class particles in configuration w . We define the *weighted partition function* as

$$Z_N(\xi; \mathbf{z}; a, b, c, d) := \sum_{w \in \mathcal{Q}(N, m)} \xi^{2\bullet(w)} \psi_w(\mathbf{z}).$$

Theorem

The generating function for the number of particles in the system is given by

$$Z_{N, m}(\xi; \mathbf{z}; a, b, c, d) = \xi^{N-m} P_{1^{N-m} 0^m}(\xi \mathbf{z} | \xi \mathbf{z}; a_\xi, b_\xi, c_\xi, d_\xi)$$

with $a_\xi = \xi a$, $b_\xi = \xi b$, $c_\xi = \xi^{-1} c$, $d_\xi = \xi^{-1} d$.

Remark: the Macdonald-Koornwinder polynomials $P_{1^{N-m} 0^m}(\mathbf{z})$ associated to single column diagrams $\lambda = 1^{N-m} 0^m$ are independent of the parameter q .

Current and density

We define an inhomogenous version of the current of particles

$$J_{N,m}(\mathbf{z}) := \frac{1}{Z_{N,m}(\xi = 1; \mathbf{z})} \sum_{w \in \mathcal{Q}(N,m)} \alpha \psi_{\circ w}(\mathbf{z}) - \gamma \psi_{\bullet w}(\mathbf{z}),$$

which under specialization $\mathbf{z} = \mathbf{1}$ reduces to the homogeneous case,

$$\langle J_{N,m} \rangle = J_{N,m}(\mathbf{1}).$$

Using the recursion relations we get

$$J_{N,m}(\mathbf{z}) = -\gamma \frac{(a - z_1)(b - z_1)}{z_1} \frac{Z_{N-1,m}(\mathbf{z} \setminus z_1)}{Z_{N,m}(\mathbf{z})}.$$

Current and density

We define an inhomogenous version of the current of particles

$$J_{N,m}(\mathbf{z}) := \frac{1}{Z_{N,m}(\xi = 1; \mathbf{z})} \sum_{w \in \mathcal{Q}(N,m)} \alpha \psi_{\circ w}(\mathbf{z}) - \gamma \psi_{\bullet w}(\mathbf{z}),$$

which under specialization $\mathbf{z} = \mathbf{1}$ reduces to the homogeneous case,

$$\langle J_{N,m} \rangle = J_{N,m}(\mathbf{1}).$$

Using the recursion relations we get

$$J_{N,m}(\mathbf{z}) = -\gamma \frac{(a - z_1)(b - z_1)}{z_1} \frac{Z_{N-1,m}(\mathbf{z} \setminus z_1)}{Z_{N,m}(\mathbf{z})}.$$

Current and density

- ▶ After specialization $\mathbf{z} = \mathbf{1}$, we obtain

$$\langle J_{N,m} \rangle = (t^{\frac{1}{2}} - t^{-\frac{1}{2}}) \frac{Z_{N-1,m}(\mathbf{1})}{Z_{N,m}(\mathbf{1})}.$$

- ▶ For the density of first class particles we have

$$\langle \rho_{N,m}^\bullet \rangle = \left(\frac{1}{2N} \frac{\partial}{\partial \xi} \log Z_{N,m}(\xi; \mathbf{1}) \right) \Big|_{\xi=0},$$

- ▶ In order to determine their asymptotic behavior $N \rightarrow \infty$, $\rho_* = m/N$, we use an integral representation of the Macdonald-Koornwinder polynomials.

Current and density

- ▶ After specialization $\mathbf{z} = \mathbf{1}$, we obtain

$$\langle J_{N,m} \rangle = (t^{\frac{1}{2}} - t^{-\frac{1}{2}}) \frac{Z_{N-1,m}(\mathbf{1})}{Z_{N,m}(\mathbf{1})}.$$

- ▶ For the density of first class particles we have

$$\langle \rho_{N,m}^\bullet \rangle = \left(\frac{1}{2N} \frac{\partial}{\partial \xi} \log Z_{N,m}(\xi; \mathbf{1}) \right) \Big|_{\xi=0},$$

- ▶ In order to determine their asymptotic behavior $N \rightarrow \infty$, $\rho_* = m/N$, we use an integral representation of the Macdonald-Koornwinder polynomials.

Current and density

- ▶ After specialization $\mathbf{z} = \mathbf{1}$, we obtain

$$\langle J_{N,m} \rangle = (t^{\frac{1}{2}} - t^{-\frac{1}{2}}) \frac{Z_{N-1,m}(\mathbf{1})}{Z_{N,m}(\mathbf{1})}.$$

- ▶ For the density of first class particles we have

$$\langle \rho_{N,m}^\bullet \rangle = \left(\frac{1}{2N} \frac{\partial}{\partial \xi} \log Z_{N,m}(\xi; \mathbf{1}) \right) \Big|_{\xi=0},$$

- ▶ In order to determine their asymptotic behavior $N \rightarrow \infty$, $\rho_* = m/N$, we use an integral representation of the Macdonald-Koornwinder polynomials.

Integral representation

Using the Cauchy identity for Macdonald-Koornwinder polynomials [Mimachi] and assuming $t < 1$, we can write

$$Z_{N,m}(\xi; \mathbf{z}) = r_m^{-1}(a_\xi, b_\xi, c_\xi, d_\xi | t) \times \xi^{N-m} \oint_C \frac{dx}{4\pi i x} \Pi(\xi \mathbf{z}, x) w(x; a_\xi, b_\xi, c_\xi, d_\xi | t) p_m(x; a_\xi, b_\xi, c_\xi, d_\xi | t)$$

$p_m(x; a, b, c, d | t)$ is the m -th Askey-Wilson polynomial of base t in the variable $\frac{x+x^{-1}}{2}$,

$$w(x; a, b, c, d | t) = \frac{(x^2, x^{-2}; t)_\infty}{(ax, ax^{-1}, bx, bx^{-1}, cx, cx^{-1}, dx, dx^{-1}; t)_\infty},$$

$$\Pi(\mathbf{z}, \mathbf{x}) = \prod_{1 \leq i \leq N} (z_i + z_i^{-1} - x - x^{-1})$$

$$r_m(a, b, c, d | t) = \frac{(abcdt^{2m}; t)_\infty}{(t^{m+1}, abt^m, act^m, adt^m, bct^m, bdt^m, cdt^m; t)_\infty}.$$

Integral representation

Using the Cauchy identity for Macdonald-Koornwinder polynomials [Mimachi] and assuming $t < 1$, we can write

$$Z_{N,m}(\xi; \mathbf{z}) = r_m^{-1}(a_\xi, b_\xi, c_\xi, d_\xi | t) \times \\ \xi^{N-m} \oint_{\mathcal{C}} \frac{dx}{4\pi ix} \Pi(\xi \mathbf{z}, x) w(x; a_\xi, b_\xi, c_\xi, d_\xi | t) p_m(x; a_\xi, b_\xi, c_\xi, d_\xi | t)$$

Remarks


- ▶ This formula at $\mathbf{z} = \mathbf{1}$ generalizes the result of Uchiyama, Sasamoto and Wadati obtained for $m = 0$.
- ▶ At $\mathbf{z} = \mathbf{1}$ improves a much more complicated formula obtained by Uchiyama.
- ▶ Comparison with results of Corteel, Mandelshtam and Williams: combinatorial representations of P_w in terms of Rhombic Staircase Tableaux?

Phase diagram

$$x_0 < a, c : J = \frac{(t^{\frac{1}{2}} - t^{-\frac{1}{2}})(1 - \rho_*^2)}{4}, \quad \rho^\bullet = \frac{1 - \rho_*}{2}$$

$$a < x_0, c : J = \frac{a(t^{-\frac{1}{2}} - t^{\frac{1}{2}})}{(1 - a)^2}, \quad \rho^\bullet = \frac{a}{a - 1} - \rho_* > \frac{1 - \rho_*}{2}$$

$$c < x_0, a : J = \frac{c(t^{-\frac{1}{2}} - t^{\frac{1}{2}})}{(1 - c)^2}, \quad \rho^\bullet = \frac{1}{1 - c} < \frac{1 - \rho_*}{2}$$



Conclusion

- ▶ The approach to the study of the stationary measure through exchange relations can be applied to any Yang-Baxter integrable stochastic process.
- ▶ Relations with multivariate orthogonal polynomials.
- ▶ Work backward: find Matrix Product representations of Macdonald-Koornwinder polynomials.
- ▶ Work with several species of particles \longrightarrow general Koornwinder polynomials at $q = 1$.

Conclusion

- ▶ The approach to the study of the stationary measure through exchange relations can be applied to any Yang-Baxter integrable stochastic process.
- ▶ Relations with multivariate orthogonal polynomials.
- ▶ Work backward: find Matrix Product representations of Macdonald-Koornwinder polynomials.
- ▶ Work with several species of particles \longrightarrow general Koornwinder polynomials at $q = 1$.

Conclusion

- ▶ The approach to the study of the stationary measure through exchange relations can be applied to any Yang-Baxter integrable stochastic process.
- ▶ Relations with multivariate orthogonal polynomials.
- ▶ Work backward: find Matrix Product representations of Macdonald-Koornwinder polynomials.
- ▶ Work with several species of particles \longrightarrow general Koornwinder polynomials at $q = 1$.