


An elementary solution of Gessel's walks in the quadrant

Mireille Bousquet-Mélou, CNRS, LaBRI, Bordeaux




Walks in the quadrant

Let \mathcal{S} be a finite subset of \mathbb{Z}^2 (set of **steps**) and $p_0 \in \mathbb{N}^2$ (starting point).

Questions


- What is the number $a(n)$ of n -step walks starting at p_0 and contained in the non-negative quadrant \mathbb{N}^2 ?
- For $(i, j) \in \mathbb{N}^2$, what is the number $a(i, j; n)$ of such walks that end at (i, j) ?

Example. $\mathcal{S} = \{10, \bar{1}0, 1\bar{1}, \bar{1}1\}$, $p_0 = (0, 0)$.




Example [Gouyou-Beauchamps 86], [mbm-Mishna 10]

Take $\mathcal{S} = \{10, \bar{1}0, 1\bar{1}, \bar{1}\bar{1}\}$ and $p_0 = (0, 0)$



Example [Gouyou-Beauchamps 86], [mbm-Mishna 10]

Take $\mathcal{S} = \{10, \bar{1}0, 1\bar{1}, \bar{1}\bar{1}\}$ and $p_0 = (0, 0)$



Nice numbers

If $n = 2m + \delta$, with $\delta \in \{0, 1\}$,

$$a(n) = \frac{n!(n+1)!}{m!(m+1)!(m+\delta)!(m+\delta+1)!}$$

Moreover, if $n = 2m + i$,

$$a(i, j; n) = \frac{(i+1)(j+1)(i+j+2)(i+2j+3)n!(n+2)!}{(m-j)!(m+1)!(m+i+2)!(m+i+j+3)!}$$

Many contributions

Adan, Banderier, Bernardi, Bostan, Cori, Denisov, Duchon, Dulucq, Fayolle, Gessel, Fisher, Flajolet, Gouyou-Beauchamps, Guttmann, Guy, Janse van Rensburg, Johnson, Kauers, Koutschan, Krattenthaler, Kurkova, Kreweras, van Leeuwarden, MacMahon, Melczer, Mishna, Niederhausen, Petkovšek, Prellberg, Raschel, Rechnitzer, Sagan, Salvy, Viennot, Wachtel, Wilf, Yeats, Zeilberger...

etc.

Generating functions

- Our original question:

$$a(n) = ? \quad a(i, j; n) = ?$$

Generating functions

- Our original question:

$$a(n) = ? \quad a(i, j; n) = ?$$

- Generating functions and their nature

$$A(t) = \sum_{n \geq 0} a(n)t^n, \quad A(x, y; t) = \sum_{i, j, n} a(i, j; n)x^i y^j t^n$$

Generating functions

- Our original question:

$$a(n) = ? \quad a(i, j; n) = ?$$

- Generating functions and their nature

$$A(t) = \sum_{n \geq 0} a(n)t^n, \quad A(x, y; t) = \sum_{i, j, n} a(i, j; n)x^i y^j t^n$$

Remarks

- $A(1, 1; t) = A(t)$
- $A(0, 0; t)$ counts walks ending at $(0, 0)$
- $A(x, 0; t)$ counts walks ending on the x -axis

Generating functions

- Our original question:

$$a(n) = ? \quad a(i, j; n) = ?$$

- Generating functions and their nature

$$A(t) = \sum_{n \geq 0} a(n)t^n, \quad A(x, y; t) = \sum_{i, j, n} a(i, j; n)x^i y^j t^n$$

Can one express these series? Are they rational? algebraic? D-finite?

Remarks

- $A(1, 1; t) = A(t)$
- $A(0, 0; t)$ counts walks ending at $(0, 0)$
- $A(x, 0; t)$ counts walks ending on the x -axis

A hierarchy of formal power series

- The formal power series $A(t)$ is **rational** if it can be written

$$A(t) = P(t)/Q(t)$$

where $P(t)$ and $Q(t)$ are polynomials in t .

- The formal power series $A(t)$ is **algebraic** (over $\mathbb{Q}(t)$) if it satisfies a (non-trivial) polynomial equation:

$$P(t, A(t)) = 0.$$

- The formal power series $A(t)$ is **D-finite (holonomic)** if it satisfies a (non-trivial) linear differential equation with polynomial coefficients:

$$P_k(t)A^{(k)}(t) + \cdots + P_0(t)A(t) = 0.$$

- Nice closure properties + asymptotics of the coefficients
- Extension to several variables (D-finite: **one DE per variable**)

Walks in the full space


Rational series

A rational generating function:

$$a(n) = |\mathcal{S}|^n \Leftrightarrow A(t) = \sum_{n \geq 0} a(n)t^n = \frac{1}{1 - |\mathcal{S}|t}$$

More generally:

$$A(x, y; t) = \frac{1}{1 - t \sum_{(i,j) \in \mathcal{S}} x^i y^j}.$$




Walks in a rational half-space

Algebraic series

The associated generating function is algebraic, given by an explicit system of polynomial equations.


[Gessel 80], [Duchon 00], [mbm-Petkovšek 00]...



Walks in the quadrant

- Start from $p_0 = (0, 0)$



$$A(x, y; t) = \sum_{i, j, n \geq 0} a(i, j; n) x^i y^j t^n = ?$$




Walks in the quadrant

- Start from $p_0 = (0, 0)$


$$A(x, y; t) = \sum_{i, j, n \geq 0} a(i, j; n) x^i y^j t^n = ?$$



D-finite



D-finite





D-finite


Walks in the quadrant

- Start from $p_0 = (0, 0)$


$$A(x, y; t) = \sum_{i, j, n \geq 0} a(i, j; n) x^i y^j t^n = ?$$




D-finite



D-finite



D-finite





Algebraic


Walks in the quadrant

- Start from $p_0 = (0, 0)$


$$A(x, y; t) = \sum_{i, j, n \geq 0} a(i, j; n) x^i y^j t^n = ?$$




D-finite




D-finite



D-finite



Algebraic





Non-D-finite


Walks in the quadrant

- Start from $p_0 = (0, 0)$


$$A(x, y; t) = \sum_{i, j, n \geq 0} a(i, j; n) x^i y^j t^n = ?$$




D-finite




D-finite




D-finite




Algebraic



Non-D-finite



Non-D-finite




D-finite

Quadrant walks with *small* steps: classification


- $\mathcal{S} \subset \{\bar{1}, 0, 1\} \setminus \{00\} \Rightarrow 2^8 = 256$ step sets (or: **models**)

Quadrant walks with *small* steps: classification

- $\mathcal{S} \subset \{\bar{1}, 0, 1\} \setminus \{00\} \Rightarrow 2^8 = 256$ step sets (or: **models**)
- However, some models are equivalent:
 - to a model of walks in the full or half-plane (\Rightarrow algebraic)



- to another model in the collection (diagonal symmetry)




\approx



- One is left with **79 interesting distinct models**.


Non-singular



Singular



Non-singular



Singular



Classification of quadrant walks with small steps

Theorem

The series $A(x, y; t)$ is D-finite iff the associated group G is finite.
It is algebraic iff, in addition, the orbit sum is zero.

[mbm-Mishna 10], [Bostan-Kauers 10]


[Kurkova-Raschel 12]

[Mishna-Rechnitzer 07], [Melczer-Mishna 13]


D-finite

non-singular non-D-finite


singular non-D-finite




Classification of quadrant walks with small steps




When the orbit sum is zero: the four algebraic models




Kreweras



reverse Kreweras



double Kreweras




Gessel

- Kreweras walks


- ▶ expression for $a(i, 0; n)$ [Kreweras 65], simplified by [Niederhausen 83]
- ▶ $A(x, y; t)$ is algebraic [Gessel 86, mbm 02/05]

- Unified approach for the Kreweras trilogy [mbm-Mishna 10]


When the orbit sum is zero: the four algebraic models




Kreweras



reverse Kreweras



double Kreweras



Gessel

- Kreweras walks

- ▶ expression for $a(i, 0; n)$ [Kreweras 65], simplified by [Niederhausen 83]
- ▶ $A(x, y; t)$ is algebraic [Gessel 86, mbm 02/05]

- Unified approach for the Kreweras trilogy [mbm-Mishna 10]

- Gessel walks:

- ▶ conjecture for $a(0, 0; n)$ [Gessel \simeq 00]
- ▶ proof of this conjecture [Kauers, Koutschan & Zeilberger 08]
- ▶ $A(t)$ and $A(x, y; t)$ are algebraic! [Bostan & Kauers 09a]
- ▶ new proof via complex analysis [Bostan, Kurkova & Raschel 13]
- ▶ an elementary and constructive proof [mbm 15]



Theorem

- The number of walks of length $n = 3m + 2i$ ending at $(i, 0)$ is

$$a(i, 0; n) = \frac{4^m(2i+1)}{(m+i+1)(2m+2i+1)} \binom{2i}{i} \binom{3m+2i}{m}$$

[Kreweras 65]

- The generating function $A(x, y; t)$ is algebraic [Gessel 86].

For instance, the series $Q(t) := A(0, 0; t^{1/3})$ has degree 3:

$$54t - 1 + (1 - 72t)Q(t) + 16tQ(t)^2 + 64t^2Q(t)^3 = 0$$



Theorem

- The number of walks of length $n = 2m$ ending at $(0, 0)$ is

$$a(0, 0; n) = 16^m \frac{(5/6)_m (1/2)_m}{(5/3)_m (2)_m},$$

where $(a)_m = a(a+1) \cdots (a+m-1)$ is the ascending factorial.

[Kauers, Koutschan, Zeilberger 08]

- The generating function $A(x, y; t)$ is algebraic.

[Bostan, Kauers 09]


For instance, the series $Q(t) := A(0, 0; \sqrt{t})$ has degree 8:

$$\begin{aligned} & 27t^7 Q^8 + 108t^6 Q^7 + 189t^5 Q^6 + 189t^4 Q^5 - 9t^3(32t^2 + 28t - 13)Q^4 \\ & - 9t^2(64t^2 + 56t - 5)Q^3 - 2t(256t^3 - 312t^2 + 156t - 5)Q^2 \\ & - (32t - 1)(16t^2 - 28t + 1)Q - 256t^3 - 576t^2 - 48t + 1 = 0. \end{aligned}$$

I. A solution of...

I. A solution of Kreweras' model

[mbm 02]



I. A solution of Kreweras' model

[mbm 02]

1. functional equation for $A(x, y; t)$
2. canceling the kernel: the roots Y_0 and Y_1
3. the group of the walk
4. symmetric functions of Y_0 and Y_1
5. Brown's quadratic method (1969)




1. A functional equation




Set step: $\mathcal{S} = \{11, \bar{1}0, 0\bar{1}\}$. Denote $\bar{x} = 1/x$ and $\bar{y} = 1/y$.


$$A(x, y; t) \equiv A(x, y) = 1 + t(xy + \bar{x} + \bar{y})A(x, y) - t\bar{x}A(0, y) - t\bar{y}A(x, 0)$$



\ominus



\ominus



1. A functional equation



Set step: $\mathcal{S} = \{11, \bar{1}0, 0\bar{1}\}$. Denote $\bar{x} = 1/x$ and $\bar{y} = 1/y$.

$$A(x, y; t) \equiv A(x, y) = 1 + t(xy + \bar{x} + \bar{y})A(x, y) - t\bar{x}A(0, y) - t\bar{y}A(x, 0)$$

or

$$(1 - t(xy + \bar{x} + \bar{y}))A(x, y) = 1 - t\bar{x}A(0, y) - t\bar{y}A(x, 0),$$

1. A functional equation



Set step: $\mathcal{S} = \{11, \bar{1}0, 0\bar{1}\}$. Denote $\bar{x} = 1/x$ and $\bar{y} = 1/y$.

$$A(x, y; t) \equiv A(x, y) = 1 + t(xy + \bar{x} + \bar{y})A(x, y) - t\bar{x}A(0, y) - t\bar{y}A(x, 0)$$

or

$$(1 - t(xy + \bar{x} + \bar{y}))A(x, y) = 1 - t\bar{x}A(0, y) - t\bar{y}A(x, 0),$$

or

$$(1 - t(xy + \bar{x} + \bar{y}))xyA(x, y) = xy - tyA(0, y) - txA(x, 0)$$

- The polynomial $1 - t(xy + \bar{x} + \bar{y})$ is the **kernel** of this equation
- The equation is linear, with two **catalytic** variables x and y (tautological at $x = 0$ or $y = 0$)

2. Canceling the kernel



- The equation:

$$(1 - t(xy + \bar{x} + \bar{y}))_{xy}A(x, y) = xy - t_{y}A(0, y) - t_{x}A(x, 0)$$

2. Canceling the kernel



- The equation:

$$(1 - t(xy + \bar{x} + \bar{y}))xyA(x, y) = xy - tyA(0, y) - txA(x, 0)$$

- The kernel $(1 - t(xy + \bar{x} + \bar{y}))$, as a polynomial in y , has two roots:

$$Y_0(x) = \frac{1 - t\bar{x} - \sqrt{(1 - t\bar{x})^2 - 4t^2x}}{2tx} = t + \bar{x}t^2 + O(t^3),$$

$$Y_1(x) = \frac{1 - t\bar{x} + \sqrt{(1 - t\bar{x})^2 - 4t^2x}}{2tx} = \frac{\bar{x}}{t} - \bar{x}^2 - t - \bar{x}t^2 + O(t^3).$$

2. Canceling the kernel



- The equation:

$$(1 - t(xy + \bar{x} + \bar{y}))xyA(x, y) = xy - tyA(0, y) - txA(x, 0)$$

- The kernel $(1 - t(xy + \bar{x} + \bar{y}))$, as a polynomial in y , has two roots:

$$Y_0(x) = \frac{1 - t\bar{x} - \sqrt{(1 - t\bar{x})^2 - 4t^2x}}{2tx} = t + \bar{x}t^2 + O(t^3),$$

$$Y_1(x) = \frac{1 - t\bar{x} + \sqrt{(1 - t\bar{x})^2 - 4t^2x}}{2tx} = \frac{\bar{x}}{t} - \bar{x}^2 - t - \bar{x}t^2 + O(t^3).$$

- Specializing y to Y_0 in the equation gives:

$$R(x) + R(Y_0) = xY_0$$

with $R(x) = txA(x, 0) = txA(0, x)$ (symmetry).

This equation characterizes $R(x)$ but... why algebraicity?

3. The group of the model



The kernel

$$K(x, y) = 1 - t(xy + \bar{x} + \bar{y})$$

is left unchanged by the rational transformations

$$\Phi : (x, y) \mapsto (\bar{x}\bar{y}, y) \quad \text{and} \quad \Psi : (x, y) \mapsto (x, \bar{x}\bar{y}).$$

3. The group of the model



The kernel

$$K(x, y) = 1 - t(xy + \bar{x} + \bar{y})$$

is left unchanged by the rational transformations


$$\Phi : (x, y) \mapsto (\bar{x}\bar{y}, y) \quad \text{and} \quad \Psi : (x, y) \mapsto (x, \bar{x}\bar{y}).$$

They are involutions, and generate a dihedral group G of order 6:

$$\begin{array}{ccccc} & & \Psi & & \\ & & \text{---} & & \\ & \Phi & (\bar{x}\bar{y}, y) & \text{---} & (\bar{x}\bar{y}, x) & \text{---} & \Phi & & \\ (x, y) & \text{---} & & & & & & & (y, x) \\ & \Psi & (x, \bar{x}\bar{y}) & \text{---} & (y, \bar{x}\bar{y}) & \text{---} & \Psi & & \\ & & & \Phi & & & & & \end{array}$$


Build more pairs that cancel the kernel

- Since $K(x, Y_0) = 0$ and the transformations of G preserve the kernel, any element of the orbit of (x, Y_0) cancels the kernel.



Build more pairs that cancel the kernel

- Since $K(x, Y_0) = 0$ and the transformations of G preserve the kernel, any element of the orbit of (x, Y_0) cancels the kernel.



Gives **two** equations for R :

$$\begin{aligned}R(x) + R(Y_0) &= xY_0 \\R(Y_0) + R(Y_1) &= Y_0Y_1 = \bar{x}\end{aligned}$$

with $R(x) = txA(x, 0)$.

4. Symmetric functions of Y_0 and Y_1

- The elementary symmetric functions of the Y_i are **polynomials in \bar{x}** :

$$Y_0 + Y_1 = \frac{\bar{x}}{t} - \bar{x}^2 \quad \text{and} \quad Y_0 Y_1 = \bar{x}.$$

4. Symmetric functions of Y_0 and Y_1

- The elementary symmetric functions of the Y_i are **polynomials in \bar{x}** :

$$Y_0 + Y_1 = \frac{\bar{x}}{t} - \bar{x}^2 \quad \text{and} \quad Y_0 Y_1 = \bar{x}.$$

- Rewrite the equations for R as

$$R(Y_0) - xY_0 = -R(x)$$

$$R(Y_1) - xY_1 = R(x) + 2\bar{x} - 1/t$$

4. Symmetric functions of Y_0 and Y_1

- The elementary symmetric functions of the Y_i are **polynomials in \bar{x}** :

$$Y_0 + Y_1 = \frac{\bar{x}}{t} - \bar{x}^2 \quad \text{and} \quad Y_0 Y_1 = \bar{x}.$$

- Rewrite the equations for R as

$$R(Y_0) - xY_0 = -R(x)$$

$$R(Y_1) - xY_1 = R(x) + 2\bar{x} - 1/t$$

- The product is symmetric in Y_0 and Y_1 :

$$(R(Y_0) - xY_0)(R(Y_1) - xY_1) = -R(x)(R(x) + 2\bar{x} - 1/t)$$

4. Symmetric functions of Y_0 and Y_1

- The elementary symmetric functions of the Y_i are **polynomials in \bar{x}** :

$$Y_0 + Y_1 = \frac{\bar{x}}{t} - \bar{x}^2 \quad \text{and} \quad Y_0 Y_1 = \bar{x}.$$

- Rewrite the equations for R as

$$R(Y_0) - xY_0 = -R(x)$$

$$R(Y_1) - xY_1 = R(x) + 2\bar{x} - 1/t$$

- The product is symmetric in Y_0 and Y_1 :

$$(R(Y_0) - xY_0)(R(Y_1) - xY_1) = -R(x)(R(x) + 2\bar{x} - 1/t)$$

- Extracting the non-negative powers of x gives:

$$x - 2R'(0) = -R(x)(R(x) + 2\bar{x} - 1/t).$$

No more Y_i ! Equivalently,

$$xtR(x)^2 + (2t - x)R(x) + tx^2 - 2txR'(0) = 0.$$

5. Finish: the quadratic method

The series $R(x) = txA(x, 0)$ satisfies

$$xtR(x)^2 + (2t - x)R(x) + tx^2 - 2txR'(0) = 0.$$

or

$$t^2x^2A(x, 0)^2 + (2t - x)A(x, 0) - 2tA(0, 0) + x = 0.$$

Tautological at $x = 0$: a quadratic equation with one **catalytic variable** x , and one additional unknown one-variable series $A(0, 0)$

⇒ Brown's quadratic method [Brown 69]

5. Finish: the quadratic method

The series $R(x) = txA(x, 0)$ satisfies

$$xtR(x)^2 + (2t - x)R(x) + tx^2 - 2txR'(0) = 0.$$

or

$$t^2x^2A(x, 0)^2 + (2t - x)A(x, 0) - 2tA(0, 0) + x = 0.$$

Tautological at $x = 0$: a quadratic equation with one **catalytic variable** x , and one additional unknown one-variable series $A(0, 0)$

⇒ Brown's quadratic method [Brown 69]

Theorem

Let $X \equiv X(t)$ be the unique series in t defined by

$$X = t(2 + X^3).$$

Then the generating function of Kreweras' walks ending on the x -axis is

$$A(x, 0; t) = \frac{1}{tx} \left(\frac{1}{2t} - \frac{1}{x} - \left(\frac{1}{X} - \frac{1}{x} \right) \sqrt{1 - xX^2} \right).$$

II. A solution of Gessel's model

[mbm 15]

II. A solution of Gessel's model

[mbm 15]

- functional equation for $A(x, y)$
- canceling the kernel: the roots Y_0 and Y_1
- the group of the walk
- symmetric functions of Y_0 and Y_1
- ...
- the *generalized* quadratic method [mbm-Jehanne 06]




1. A functional equation




Set step: $\mathcal{S} = \{11, 10, \bar{1}\bar{1}, 0\bar{1}\}$


$$A(x, y) = 1 + t(xy + x + \bar{x}\bar{y} + \bar{x})A(x, y) \\ - t\bar{x}(1 + \bar{y})A(0, y) - t\bar{x}\bar{y}(A(x, 0) - A(0, 0))$$



⊖



⊖



1. A functional equation



Set step: $\mathcal{S} = \{11, 10, \bar{1}\bar{1}, 0\bar{1}\}$

$$A(x, y) = 1 + t(xy + x + \bar{x}\bar{y} + \bar{x})A(x, y) \\ - t\bar{x}(1 + \bar{y})A(0, y) - t\bar{x}\bar{y}(A(x, 0) - A(0, 0))$$

or

$$(1 - t(xy + x + \bar{x}\bar{y} + \bar{x}))xyA(x, y) = xy - S(y) - R(x)$$

with

$$S(y) = t(1 + y)A(0, y) \quad \text{and} \quad R(x) = t(A(x, 0) - A(0, 0)).$$

- Main difference with K' walks: no x/y symmetry, two bivariate unknown series $R(x)$ and $S(y)$

2. Canceling the kernel



- The equation:

$$(1 - t(xy + x + \bar{x}\bar{y} + \bar{x}))xyA(x, y) = xy - S(y) - R(x)$$

2. Canceling the kernel



- The equation:

$$(1 - t(xy + x + \bar{x}\bar{y} + \bar{x}))xyA(x, y) = xy - S(y) - R(x)$$

- The kernel, as a polynomial in y , has two roots:

$$Y_0(x) = \frac{1 - t(x + \bar{x}) - \sqrt{(1 - t(x + \bar{x}))^2 - 4t^2}}{2tx} = \bar{x}t + \dots$$

$$Y_1(x) = \frac{1 - t(x + \bar{x}) + \sqrt{(1 - t(x + \bar{x}))^2 - 4t^2}}{2tx} = \frac{\bar{x}}{t} - (1 + \bar{x}^2) - \bar{x}t + \dots$$

Observe that $xY_i(x)$ is symmetric in x and \bar{x} .

2. Canceling the kernel



- The equation:

$$(1 - t(xy + x + \bar{x}\bar{y} + \bar{x}))xyA(x, y) = xy - S(y) - R(x)$$

- The kernel, as a polynomial in y , has two roots:

$$Y_0(x) = \frac{1 - t(x + \bar{x}) - \sqrt{(1 - t(x + \bar{x}))^2 - 4t^2}}{2tx} = \bar{x}t + \dots$$

$$Y_1(x) = \frac{1 - t(x + \bar{x}) + \sqrt{(1 - t(x + \bar{x}))^2 - 4t^2}}{2tx} = \frac{\bar{x}}{t} - (1 + \bar{x}^2) - \bar{x}t + \dots$$

Observe that $xY_i(x)$ is symmetric in x and \bar{x} .

- Specializing y to Y_0 in the equation gives:

$$R(x) + S(Y_0) = xY_0$$

3. The group of the model



The kernel

$$K(x, y) = 1 - t(xy + x + \bar{x}\bar{y} + \bar{x})$$

is left unchanged by the rational transformations

$$\Phi : (x, y) \mapsto (\bar{x}\bar{y}, y) \quad \text{and} \quad \Psi : (x, y) \mapsto (x, \bar{x}^2\bar{y}).$$

3. The group of the model




The kernel

$$K(x, y) = 1 - t(xy + x + \bar{x}\bar{y} + \bar{x})$$

is left unchanged by the rational transformations

$$\Phi : (x, y) \mapsto (\bar{x}\bar{y}, y) \quad \text{and} \quad \Psi : (x, y) \mapsto (x, \bar{x}^2\bar{y}).$$

They are involutions, and generate a dihedral group of order 8:




Build more pairs that cancel the kernel

Since $K(x, Y_0) = 0$ and the transformations of G preserve the kernel, any element of the orbit of (x, Y_0) cancels the kernel.

$$\begin{array}{ccccccc} & \Phi & & \Psi & & \Phi & & \Psi \\ & \swarrow & & \xrightarrow{\quad} & & \xrightarrow{\quad} & & \searrow \\ (x, Y_0) & & (xY_1, Y_0) & & (xY_1, x^2 Y_0) & & (\bar{x}, x^2 Y_0) & & (\bar{x}, x^2 Y_1) \\ & \searrow & & \xrightarrow{\quad} & & \xrightarrow{\quad} & & \swarrow \\ & \Psi & & \Phi & & \Psi & & \Phi \end{array}$$


Build more pairs that cancel the kernel

Since $K(x, Y_0) = 0$ and the transformations of G preserve the kernel, any element of the orbit of (x, Y_0) cancels the kernel.



Build more pairs that cancel the kernel

Since $K(x, Y_0) = 0$ and the transformations of G preserve the kernel, any element of the orbit of (x, Y_0) cancels the kernel.



Gives **four** equations for R and S :

$$R(x) + S(Y_0) = xY_0$$

$$R(Y_0) + S(Y_1) = Y_0 Y_1 = \bar{x}$$

$$R(\bar{x}) + S(x^2 Y_0) = xY_0$$

$$R(xY_0) + S(x^2 Y_1) = x^3 Y_0 Y_1 = x.$$

4. Symmetric functions of Y_0 and Y_1

- The elementary symmetric functions of the Y_i are **polynomials in \bar{x}** :

$$Y_0 + Y_1 = -1 + \frac{\bar{x}}{t} - \bar{x}^2 \quad \text{and} \quad Y_0 Y_1 = \bar{x}^2.$$

4. Symmetric functions of Y_0 and Y_1

- The elementary symmetric functions of the Y_i are **polynomials in \bar{x}** :

$$Y_0 + Y_1 = -1 + \frac{\bar{x}}{t} - \bar{x}^2 \quad \text{and} \quad Y_0 Y_1 = \bar{x}^2.$$

- From the four equations:

$$R(x) + S(Y_0) = xY_0$$

$$R(xY_0) + S(Y_1) = Y_0 Y_1 = \bar{x}$$

$$R(\bar{x}) + S(x^2 Y_0) = xY_0$$

$$R(xY_0) + S(x^2 Y_1) = x^3 Y_0 Y_1 = x,$$

one constructs **two** symmetric functions of Y_0 and Y_1 , and extracts non-negative powers of x .

This gives an equation without any Y_i ... but with $R(x)$ and $R(\bar{x})$.

4. Symmetric functions of Y_0 and Y_1 (cont'd)

This gives:

$$R(x)^2 + R(x)R(\bar{x}) + R(\bar{x})^2 + (2\bar{x} - 1/t + x + S(0))R(x) \\ + (2x - 1/t + \bar{x} + S(0))R(\bar{x}) = 2R'(0) - (\bar{x} + x - 1/t)S(0) - 1$$

while for Kreweras' walks:

$$xtR(x)^2 + (2t - x)R(x) + tx^2 - 2txR'(0) = 0.$$

- Main difference: the terms $R(\bar{x})$
- Main difficulty: the hybrid term $R(x)R(\bar{x})$

5. Les identités remarquables à notre secours

The equation reads:

$$R(x)^2 + R(x)R(\bar{x}) + R(\bar{x})^2 + (2\bar{x} - 1/t + x + S(0))R(x) \\ + (2x - 1/t + \bar{x} + S(0))R(\bar{x}) = 2R'(0) - (\bar{x} + x - 1/t)S(0) - 1.$$

But but but...

$$(a^2 + ab + b^2)(a - b) = a^3 - b^3$$

5. Les identités remarquables à notre secours

The equation reads:

$$R(x)^2 + R(x)R(\bar{x}) + R(\bar{x})^2 + (2\bar{x} - 1/t + x + S(0))R(x) \\ + (2x - 1/t + \bar{x} + S(0))R(\bar{x}) = 2R'(0) - (\bar{x} + x - 1/t)S(0) - 1.$$

But but but...

$$(a^2 + ab + b^2)(a - b) = a^3 - b^3$$

⇒ Multiply by $R(x) - R(\bar{x}) + \bar{x} - x$!

5. Les identités remarquables à notre secours

The equation reads:

$$R(x)^2 + R(x)R(\bar{x}) + R(\bar{x})^2 + (2\bar{x} - 1/t + x + S(0))R(x) \\ + (2x - 1/t + \bar{x} + S(0))R(\bar{x}) = 2R'(0) - (\bar{x} + x - 1/t)S(0) - 1.$$

But but but...

$$(a^2 + ab + b^2)(a - b) = a^3 - b^3$$

⇒ Multiply by $R(x) - R(\bar{x}) + \bar{x} - x$! This gives $P(x) = P(\bar{x})$, with

$$P(x) = R(x)^3 + (S(0) + 3\bar{x} - 1/t)R(x)^2 \\ + (2\bar{x}^2 - \bar{x}/t + x/t - x^2 - 2R'(0) + (2\bar{x} - 1/t)S(0)) R(x) \\ - x^2 S(0) + x (2R'(0) + S(0)/t - 1).$$

5. Les identités remarquables à notre secours

We have obtained $P(x) = P(\bar{x})$, with

$$\begin{aligned}P(x) &= R(x)^3 + (S(0) + 3\bar{x} - 1/t)R(x)^2 \\ &+ (2\bar{x}^2 - \bar{x}/t + x/t - x^2 - 2R'(0) + (2\bar{x} - 1/t)S(0)) R(x) \\ &\quad - x^2 S(0) + x (2R'(0) + S(0)/t - 1).\end{aligned}$$

Extract nonnegative powers of x :

$$\begin{aligned}R(x)^3 &+ (S(0) + 3\bar{x} - 1/t)R(x)^2 \\ &+ (2\bar{x}^2 - \bar{x}/t + x/t - x^2 - 2R'(0) + (2\bar{x} - 1/t)S(0)) R(x) \\ &= R''(0) + R'(0)(2S(0) + 2\bar{x} - 1/t) + xS(0)(x - 1/t) + x.\end{aligned}$$

A cubic equation in one catalytic variable x , with three additional unknown one-variable series $S(0)$, $R'(0)$ and $R''(0)$.

6. Finish: the generalized quadratic method

$$\begin{aligned} &R(x)^3 + (S(0) + 3\bar{x} - 1/t)R(x)^2 \\ &+ (2\bar{x}^2 - \bar{x}/t + x/t - x^2 - 2R'(0) + (2\bar{x} - 1/t)S(0)) R(x) \\ &= R''(0) + R'(0)(2S(0) + 2\bar{x} - 1/t) + xS(0)(x - 1/t) + x. \end{aligned}$$

Theorem

Every series $R(x; t) \equiv R(x)$ solution of a (proper) polynomial equation with one catalytic variable x is algebraic. That is, if

$$P(t, x, R(x), A_1, \dots, A_k) = 0$$

for some polynomial P with coefficients in \mathbb{Q} , then $R(x)$ is algebraic over $\mathbb{Q}(t, x)$, and each A_i is algebraic over $\mathbb{Q}(t)$.

[mbm-Jehanne 2006] (after [Knuth 72] and [Brown 65])

6. Finish: the generalized quadratic method

Theorem

The generating function $A(x, y; t)$ is algebraic of degree 72 (explicit equations and rational parametrizations for $A(x, 0)$ and $A(0, y)$)

- In particular, the series $Q(t) := A(0, 0; \sqrt{t})$ has degree 8:

$$\begin{aligned} 27t^7 Q^8 + 108t^6 Q^7 + 189t^5 Q^6 + 189t^4 Q^5 - 9t^3(32t^2 + 28t - 13)Q^4 \\ - 9t^2(64t^2 + 56t - 5)Q^3 - 2t(256t^3 - 312t^2 + 156t - 5)Q^2 \\ - (32t - 1)(16t^2 - 28t + 1)Q - 256t^3 - 576t^2 - 48t + 1 = 0. \end{aligned}$$


- The number of walks of length $n = 2m$ ending at $(0, 0)$ is

$$a(0, 0; 2m) = 16^m \frac{(5/6)_m (1/2)_m}{(5/3)_m (2)_m},$$

where $(a)_m = a(a+1) \cdots (a+m-1)$ is the ascending factorial.


Comments

- other elementary proofs will come for sure
- 3D Kreweras' walks remain mysterious
- works for other algebraic models with repeated steps, proved so far by computer algebra [Kauers, Yatchak 14(a)]




Comments

- other elementary proofs will come for sure
- 3D Kreweras' walks remain mysterious
- works for other algebraic models with repeated steps, proved so far by computer algebra [Kauers, Yatchak 14(a)]




- Open: Find elementary proofs for



Algebraic

Conj. [Kauers, Yatchak 14(a)]

Proof via complex analysis [R. et al.]



D-finite

Proof via computer algebra

[Kauers, Yatchak 14(a)]

Polynomial equations with one catalytic variable

- Assume

$$P(F(x), A_1, \dots, A_k, t, x) = 0$$

where $P(x_0, x_1, \dots, x_k, t, x)$ is a polynomial with coefficients in \mathbb{K} , $F(x) \equiv F(x; t) \in \mathbb{K}[x][[t]]$, and $A_i \in \mathbb{K}[[t]]$ for all i .

Polynomial equations with one catalytic variable

- Assume

$$P(F(x), A_1, \dots, A_k, t, x) = 0$$

where $P(x_0, x_1, \dots, x_k, t, x)$ is a polynomial with coefficients in \mathbb{K} , $F(x) \equiv F(x; t) \in \mathbb{K}[x][[t]]$, and $A_i \in \mathbb{K}[[t]]$ for all i .

- For any series $X \equiv X(t)$ such that
 - the series $F(X) \equiv F(X; t)$ is well-defined
 - $\frac{\partial P}{\partial x_0}(F(X), A_1, \dots, A_k, t, X) = 0$,

one has

$$\frac{\partial P}{\partial x}(F(X), A_1, \dots, A_k, t, X) = 0.$$

Polynomial equations with one catalytic variable

- Assume

$$P(F(x), A_1, \dots, A_k, t, x) = 0$$

where $P(x_0, x_1, \dots, x_k, t, x)$ is a polynomial with coefficients in \mathbb{K} , $F(x) \equiv F(x; t) \in \mathbb{K}[x][[t]]$, and $A_i \in \mathbb{K}[[t]]$ for all i .

- For any series $X \equiv X(t)$ such that
 - the series $F(X) \equiv F(X; t)$ is well-defined
 - $\frac{\partial P}{\partial x_0}(F(X), A_1, \dots, A_k, t, X) = 0$,

one has

$$\frac{\partial P}{\partial x}(F(X), A_1, \dots, A_k, t, X) = 0.$$

Proof: differentiate (1) with respect to x

$$F'(x) \frac{\partial P}{\partial x_0}(F(x), A_1, \dots, A_k, t, x) + \frac{\partial P}{\partial x}(F(x), A_1, \dots, A_k, t, x) = 0.$$

Polynomial equations with one catalytic variable

- Assume there exist k series X_1, \dots, X_k such that

$$\frac{\partial P}{\partial x_0}(F(X_i), A_1, \dots, A_k, t, X_i) = 0.$$

In this case, for each X_i ,

$$\frac{\partial P}{\partial x}(F(X_i), A_1, \dots, A_k, t, X_i) = 0$$

and

$$P(F(X_i), A_1, \dots, A_k, t, X_i) = 0.$$

- This system of $3k$ **polynomial** equations in $3k$ unknowns $A_1, \dots, A_k, X_1, \dots, X_k, F(X_1), \dots, F(X_k)$ may imply (together with the fact that the X_i are distinct) the **algebraicity** of the A_i .



Example. When $\mathcal{S} = \{\bar{1}0, 0\bar{1}, 11\}$, the equation reads

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyA(0, y; t) - txA(x, 0; t).$$

Naïve route: guess and check!

- Guess a polynomial equation Pol satisfied by $A(x, y; t)$
(degrees $[18, 18, 17, 12]$ in x, y, t, A)
- Let $F(x, y; t)$ be the solution of Pol that coincides with $A(x, y; t)$ up to high order (in t)
- Prove that $F(x, y; t)$ is a formal power series in t with **polynomial coefficients in x and y** $\Rightarrow F(x, 0; t)$ and $F(0, y; t)$ are well-defined
- By taking resultants, prove that $F(x, y; t)$ satisfies the above functional equation.



Example. When $\mathcal{S} = \{\bar{1}0, 0\bar{1}, 11\}$, the equation reads

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyA(0, y; t) - txA(x, 0; t).$$

Naïve route: guess and check!

- Guess a polynomial equation Pol satisfied by $A(x, y; t)$
(degrees $[18, 18, 17, 12]$ in x, y, t, A)
- Let $F(x, y; t)$ be the solution of Pol that coincides with $A(x, y; t)$ up to high order (in t)
- Prove that $F(x, y; t)$ is a formal power series in t with **polynomial coefficients in x and y** $\Rightarrow F(x, 0; t)$ and $F(0, y; t)$ are well-defined
- By taking resultants, prove that $F(x, y; t)$ satisfies the above functional equation.



Example. When $\mathcal{S} = \{\bar{1}0, 0\bar{1}, 11\}$, the equation reads

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyA(0, y; t) - txA(x, 0; t).$$

Naïve route: guess and check!

- Guess a polynomial equation Pol satisfied by $A(x, y; t)$
(degrees $[18, 18, 17, 12]$ in x, y, t, A)
- Let $F(x, y; t)$ be the solution of Pol that coincides with $A(x, y; t)$ up to high order (in t)
- Prove that $F(x, y; t)$ is a formal power series in t with **polynomial coefficients in x and y** $\Rightarrow F(x, 0; t)$ and $F(0, y; t)$ are well-defined
- By taking resultants, prove that $F(x, y; t)$ satisfies the above functional equation.



Example. When $\mathcal{S} = \{\bar{1}0, 0\bar{1}, 11\}$, the equation reads

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyA(0, y; t) - txA(x, 0; t).$$

Naïve route: guess and check!

- Guess a polynomial equation Pol satisfied by $A(x, y; t)$
(degrees $[18, 18, 17, 12]$ in x, y, t, A)
- Let $F(x, y; t)$ be the solution of Pol that coincides with $A(x, y; t)$ up to high order (in t)
- Prove that $F(x, y; t)$ is a formal power series in t with **polynomial coefficients in x and y** $\Rightarrow F(x, 0; t)$ and $F(0, y; t)$ are well-defined
- By taking resultants, prove that $F(x, y; t)$ satisfies the above functional equation.



Example. When $\mathcal{S} = \{\bar{1}0, 0\bar{1}, 11\}$, the equation reads

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyA(0, y; t) - txA(x, 0; t).$$

Naïve route: guess and check!

- Guess a polynomial equation Pol satisfied by $A(x, y; t)$
(degrees $[18, 18, 17, 12]$ in x, y, t, A)
- Let $F(x, y; t)$ be the solution of Pol that coincides with $A(x, y; t)$ up to high order (in t)
- Prove that $F(x, y; t)$ is a formal power series in t with **polynomial coefficients in x and y** $\Rightarrow F(x, 0; t)$ and $F(0, y; t)$ are well-defined
- By taking resultants, prove that $F(x, y; t)$ satisfies the above functional equation.
- Then

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyF(y, 0; t) - txF(x, 0; t).$$



Example. When $\mathcal{S} = \{\bar{1}0, 0\bar{1}, 11\}$, the equation reads

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyA(0, y; t) - txA(x, 0; t).$$

A less naïve route

- Guess a polynomial equation Pol satisfied by $A(x, 0; t) = A(0, x; t)$ (degrees [6, 10, 6] in x, t, A)
- Let $F(x, 0; t)$ be the solution of Pol that coincides with $A(x, 0; t)$ up to high order (in t)
- Prove that $F(x, 0; t)$ is a formal power series in t with **polynomial coefficients in x**
- By taking resultants, prove that F satisfies the equation obtained by canceling the kernel:

$$txF(x, 0; t) + tY_0F(Y_0, 0; t) = xY_0.$$

- Then

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyF(y, 0; t) - txF(x, 0; t).$$



Example. When $\mathcal{S} = \{\bar{1}0, 0\bar{1}, 11\}$, the equation reads

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyA(0, y; t) - txA(x, 0; t).$$

A less naïve route

- Guess a polynomial equation Pol satisfied by $A(x, 0; t) = A(0, x; t)$ (degrees $[6, 10, 6]$ in x, t, A)
- Let $F(x, 0; t)$ be the solution of Pol that coincides with $A(x, 0; t)$ up to high order (in t)
- Prove that $F(x, 0; t)$ is a formal power series in t with **polynomial coefficients in x**
- By taking resultants, prove that F satisfies the equation obtained by canceling the kernel:

$$txF(x, 0; t) + tY_0F(Y_0, 0; t) = xY_0.$$

- Then

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyF(y, 0; t) - txF(x, 0; t).$$



Example. When $\mathcal{S} = \{\bar{1}0, 0\bar{1}, 11\}$, the equation reads

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyA(0, y; t) - txA(x, 0; t).$$

A less naïve route

- Guess a polynomial equation Pol satisfied by $A(x, 0; t) = A(0, x; t)$ (degrees $[6, 10, 6]$ in x, t, A)
- Let $F(x, 0; t)$ be the solution of Pol that coincides with $A(x, 0; t)$ up to high order (in t)
- Prove that $F(x, 0; t)$ is a formal power series in t with **polynomial coefficients in x**
- By taking resultants, prove that F satisfies the equation obtained by canceling the kernel:

$$txF(x, 0; t) + tY_0F(Y_0, 0; t) = xY_0.$$

- Then

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyF(y, 0; t) - txF(x, 0; t).$$



Example. When $\mathcal{S} = \{\bar{1}0, 0\bar{1}, 11\}$, the equation reads

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyA(0, y; t) - txA(x, 0; t).$$

A less naïve route

- Guess a polynomial equation Pol satisfied by $A(x, 0; t) = A(0, x; t)$ (degrees $[6, 10, 6]$ in x, t, A)
- Let $F(x, 0; t)$ be the solution of Pol that coincides with $A(x, 0; t)$ up to high order (in t)
- Prove that $F(x, 0; t)$ is a formal power series in t with **polynomial coefficients in x**
- By taking resultants, prove that F satisfies the equation obtained by canceling the kernel:

$$txF(x, 0; t) + tY_0F(Y_0, 0; t) = xY_0.$$

- Then

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyF(y, 0; t) - txF(x, 0; t).$$



Example. When $\mathcal{S} = \{\bar{1}0, 0\bar{1}, 11\}$, the equation reads

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyA(0, y; t) - txA(x, 0; t).$$

A less naïve route

- Guess a polynomial equation Pol satisfied by $A(x, 0; t) = A(0, x; t)$ (degrees $[6, 10, 6]$ in x, t, A)
- Let $F(x, 0; t)$ be the solution of Pol that coincides with $A(x, 0; t)$ up to high order (in t)
- Prove that $F(x, 0; t)$ is a formal power series in t with **polynomial coefficients in x**
- By taking resultants, prove that F satisfies the equation obtained by canceling the kernel:

$$txF(x, 0; t) + tY_0F(Y_0, 0; t) = xY_0.$$

- Then

$$(1 - t(\bar{x} + \bar{y} + xy))xyA(x, y; t) = xy - tyF(y, 0; t) - txF(x, 0; t).$$



Algebraicity of Gessel's model [Bostan-Kauers 10]

- When $\mathcal{S} = \{10, \bar{1}0, 11, \bar{1}\bar{1}\}$, the series $A(x, y; t)$ is algebraic (degree 72).
- In particular, the series $A(0, 0; t)$, which counts loops, has degree 8, and the following expansion:

$$A(0, 0; t) = \sum_{n \geq 0} 16^n \frac{(5/6)_n (1/2)_n}{(5/3)_n (2)_n} t^{2n}, \quad (1)$$

with $(i)_n = i(i+1) \cdots (i+n-1)$.


(1): Conjectured around 2001, first proof by [Kauers, Koutschan & Zeilberger 09] using computer algebra.

The algebraicity of $A(x, y; t)$ has now been re-proved using a complex analysis approach [Bostan, Kurkova & Raschel 13(a)]

A complex analysis approach

Markov chains with small steps in the quadrant: **stationary distribution(s)**
[Malyshev 71+]

Le petit livre jaune
[Fayolle, Iasnogorodski & Malyshev 99]



⇒ Reduction to a boundary value problem of the Riemann-Carleman type

A complex analysis approach

An expression of Q for any non-singular model \mathcal{S}

$$\tilde{K}(x, 0; t)A(x, 0; t) - \tilde{K}(0, 0; t)A(0, 0; t) = xY_0(x; t) + \frac{1}{2i\pi} \int_{x_1(t)}^{x_2(t)} u [Y_0(u; t) - Y_1(u; t)] \left[\frac{\partial_u w(u; t)}{w(u; t) - w(x; t)} - \frac{\partial_u w(u; t)}{w(u; t) - w(0; t)} \right] du$$

where Y_0 , Y_1 , x_1 and x_2 are explicit algebraic series and w is explicit/
very well understood.

[Raschel 12] + Fayolle, Kurkova

+ other formulas that complete the characterization of $A(x, y; t)$

A complex analysis approach

An expression of Q for any non-singular model \mathcal{S}

$$\tilde{K}(x, 0; t)A(x, 0; t) - \tilde{K}(0, 0; t)A(0, 0; t) = xY_0(x; t) + \frac{1}{2i\pi} \int_{x_1(t)}^{x_2(t)} u [Y_0(u; t) - Y_1(u; t)] \left[\frac{\partial_u w(u; t)}{w(u; t) - w(x; t)} - \frac{\partial_u w(u; t)}{w(u; t) - w(0; t)} \right] du$$

where Y_0 , Y_1 , x_1 and x_2 are explicit algebraic series and w is explicit/very well understood.

In particular, w is D-finite (in fact, algebraic!) iff the group is finite.

[Raschel 12] + Fayolle, Kurkova

+ other formulas that complete the characterization of $A(x, y; t)$

A complex analysis approach: climax(es)

Theorems

- If \mathcal{S} has an infinite group and is not singular, then $A(x, y; t)$ is not D-finite in x (\equiv no differential equation with respect to x)

[Kurkova & Raschel 12]

- A new proof of the algebraicity of Gessel's model



[Bostan, Kurkova & Raschel 13(a)]