


# Eulerian orientations of planar maps


Mireille Bousquet-Mélou, CNRS, Bordeaux, France

Andrew Elvey Price, Bordeaux





ArXiv: 1803.08265



# Planar maps



# Planar maps




# Planar maps




There are finitely many maps with  $n$  edges


# Planar maps




# Planar maps




# Planar maps: rooted version



# Planar maps: rooted version




# Degree constraints: 4-valent maps and quadrangulations




A 4-valent map

# Degree constraints: 4-valent maps and quadrangulations




A 4-valent map




Its dual

# Degree constraints: 4-valent maps and quadrangulations




A 4-valent map




A quadrangulation

# Degree constraints: 4-valent maps and quadrangulations




A near-quadrangulation  
with outer degree 6




A quadrangulation

# A chronology of planar maps




- **Recursive approach:** Tutte, Brown, Bender, Canfield, Richmond, Goulden, Jackson, Wormald, Walsh, Lehman, Gao, Wanless...
- **Matrix integrals:** Brézin, Itzykson, Parisi, Zuber, Bessis, Ginsparg, Zinn-Justin, Boulatov, Kazakov, Mehta, Bouttier, Di Francesco, Guitter, Eynard...
- **Bijections:** Cori & Vauquelin, Schaeffer, Bouttier, Di Francesco & Guitter (BDG), Bernardi, Fusy, Poulalhon, mbm, Chapuy...
- **Geometric properties of random maps:** Chassaing & Schaeffer, BDG, Marckert & Mokkadem, **Jean-François Le Gall**, Miermont, Curien, Albenque, Bettinelli, Ménard, Angel, Sheffield, Miller, Gwynne...


# A chronology of planar maps




# A chronology of planar maps



# A chronology of planar maps



... but for maps equipped with an Eulerian orientation




# I. Counting planar maps: the early days


William Tutte and the recursive method



## Counting quadrangulations by edge deletion




## Counting quadrangulations by edge deletion



This increases the outer degree! (by 2)



## Counting quadrangulations by edge deletion



This increases the outer degree! (by 2)

OK... then count **near**-quadrangulations, and record the outer degree:  
 $q(n, d)$  near-quadrangulations with  $n$  inner faces and outer degree  $d$

# Counting quadrangulations by edge deletion



$q(n, d)$  near-quadrangulations with  $n$  inner faces and outer degree  $d$

# Counting quadrangulations by edge deletion



$q(n, d)$  near-quadrangulations with  $n$  inner faces and outer degree  $d$

# Counting quadrangulations by edge deletion



$q(n, d)$  near-quadrangulations with  $n$  inner faces and outer degree  $d$

**Recurrence relation:**  $q(0, 0) = 1$  and

$$q(n, d) = \sum_{n_1+n_2=n} \sum_{d_1+d_2=d-2} q(n_1, d_1)q(n_2, d_2) + q(n-1, d+2)$$

# Counting quadrangulations by edge deletion



$q(n, d)$  near-quadrangulations with  $n$  inner faces and outer degree  $d$

**Recurrence relation:**  $q(0, 0) = 1$  and

$$q(n, d) = \sum_{n_1+n_2=n} \sum_{d_1+d_2=d-2} q(n_1, d_1)q(n_2, d_2) + q(n-1, d+2)$$

**Generating function:**  $Q(t, x) = \sum_{n, d \geq 0} q(n, d)t^n x^d$

$$Q(t, x) = 1 + x^2 Q(t, x)^2 + \frac{t}{x^2} (Q(t, x) - 1 - x^2 Q_2(t)),$$

with

$$Q_2(t) = [x^2]Q(t, x) = \sum_{n \geq 0} q(n, 2)t^n.$$

# Counting quadrangulations by edge deletion

Functional equation:

$$Q(t, x) = 1 + x^2 Q(t, x)^2 + \frac{t}{x^2} (Q(t, x) - 1 - x^2 Q_2(t)),$$

with

$$Q_2(t) = \sum_{n \geq 0} q(n, 2) t^n.$$

# Counting quadrangulations by edge deletion

Functional equation:

$$Q(t, x) = 1 + x^2 Q(t, x)^2 + \frac{t}{x^2} (Q(t, x) - 1 - x^2 Q_2(t)),$$

with

$$Q_2(t) = \sum_{n \geq 0} q(n, 2) t^n.$$

The original solution: guess and check

$$Q_2(t) = \frac{(1 - 12t)^{3/2} - 1 + 18t}{54t^2} \quad \text{and} \quad q(n, 2) = 2 \cdot 3^n \frac{(2n)!}{n!(n+2)!}.$$

# Counting quadrangulations by edge deletion

Functional equation:

$$Q(t, x) = 1 + x^2 Q(t, x)^2 + \frac{t}{x^2} (Q(t, x) - 1 - x^2 Q_2(t)),$$


with

$$Q_2(t) = \sum_{n \geq 0} q(n, 2) t^n.$$

The original solution: guess and check

$$Q_2(t) = \frac{(1 - 12t)^{3/2} - 1 + 18t}{54t^2} \quad \text{and} \quad q(n, 2) = 2 \cdot 3^n \frac{(2n)!}{n!(n+2)!}.$$

This is also the number of quadrangulations with  $n$  faces.



# Counting quadrangulations by edge deletion

Functional equation:

$$Q(t, x) = 1 + x^2 Q(t, x)^2 + \frac{t}{x^2} (Q(t, x) - 1 - x^2 Q_2(t)),$$


with

$$Q_2(t) = \sum_{n \geq 0} q(n, 2) t^n.$$

The original solution: guess and check

$$Q_2(t) = \frac{(1 - 12t)^{3/2} - 1 + 18t}{54t^2} \quad \text{and} \quad q(n, 2) = 2 \cdot 3^n \frac{(2n)!}{n!(n+2)!}.$$

This is also the number of quadrangulations with  $n$  faces.



# The recursive method

- One may have to
  - ▶ introduce more general objects (near-quadrangulations)
  - ▶ introduce new enumeration parameters (the outer degree)
- Solving can be non-trivial!

$$Q(t, x) = 1 + x^2 Q(t, x)^2 + \frac{t}{x^2} (Q(t, x) - 1 - x^2 Q_2(t))$$

## II. Maps with an additional structure

Eulerian orientations  
and the 6-vertex model



# Maps equipped with an additional structure

In combinatorics, probability and statistical physics


- **How many maps equipped with...**

- a spanning tree [Mullin 67, Bernardi]
- a spanning forest? [Bouttier et al., Sportiello et al., mbm & Courtiel]
- a self-avoiding walk? [Duplantier & Kostov; Gwynne & Miller]
- a proper  $q$ -colouring? [Tutte 74-83, Bouttier et al.]
- a bipolar orientation? [Kenyon, Miller, Sheffield, Wilson, Fusy, mbm...]


- **What is the expected partition function of...**

- the Ising model? [Boulatov, Kazakov, mbm, Schaeffer, Chen, Turunen, Bouttier et al., Albenque, Ménard...]
- the hard-particle model? [mbm, Schaeffer, Jehanne, Bouttier et al.]
- the Potts model? [Eynard-Bonnet, Baxter, mbm-Bernardi, Guionnet et al., Borot et al., ...]
- ...

# Maps equipped with an additional structure




# Eulerian orientations




Each vertex has equally many incoming as outgoing edges.


# Eulerian orientations




The 4-valent case: the ice model




# Eulerian orientations




The 4-valent case: the ice model




The 6-vertex model



# Eulerian orientations




The 4-valent case: the ice model




$$\omega = 1$$


The 6-vertex model




# Eulerian orientations



The 4-valent case: the ice model



The 6-vertex model



# The number of planar Eulerian orientations


## Theorem [mbm & Elvey Price 18]

Let  $S \equiv S(t) = t - 2t^2 - 4t^3 - 20t^4 - 132t^5 - \dots$  be the unique power series with constant term 0 satisfying

$$t = \sum_{n=0}^{\infty} \frac{1}{n+1} \binom{2n}{n}^2 S^{n+1}.$$

The generating function of planar Eulerian orientations counted by edges is

$$\begin{aligned} G(t) &= \frac{1}{4t^2} (t - 2t^2 - S(t)) \\ &= t + 5t^2 + 33t^3 + \dots \end{aligned}$$



# The number of planar Eulerian orientations

## Theorem [mbm & Elvey Price 18]

Let  $S \equiv S(t) = t - 2t^2 - 4t^3 - 20t^4 - 132t^5 - \dots$  be the unique power series with constant term 0 satisfying

$$t = \sum_{n=0}^{\infty} \frac{1}{n+1} \binom{2n}{n}^2 S^{n+1}.$$

The generating function of planar Eulerian orientations counted by edges is

$$\begin{aligned} G(t) &= \frac{1}{4t^2} (t - 2t^2 - S(t)) \\ &= t + 5t^2 + 33t^3 + \dots \end{aligned}$$


Asymptotics:  $G(t) = \sum_n g_n t^n$  where

$$g_n \sim \frac{1}{16} \frac{(4\pi)^{n+2}}{n^2 (\log n)^2}$$


### III. Two bijections

1. The dual of an Eulerian orientation: a **labelled map**
2. A bijection between 4-valent Eulerian orientations **with no alternating vertex** ( $n$  vertices) and general Eulerian orientations with  $n$  edges

# 1. From Eulerian orientations to labelled maps




# 1. From Eulerian orientations to labelled maps




The orientation defines a height function on the faces


# 1. From Eulerian orientations to labelled maps





The orientation defines a height function on the faces



# 1. From Eulerian orientations to labelled maps




The orientation defines a height function on the faces




[Elvey Price & Guttmann 18]

# 1. From Eulerian orientations to labelled maps





The orientation defines a height function on the faces




*Labelled maps* are rooted planar maps with labelled vertices such that:

- the root edge is labelled from 0 to 1,
- adjacent labels differ by  $\pm 1$ .

# 1. From Eulerian orientations to labelled maps





The 4-valent case: labelled quadrangulations




*Labelled maps* are rooted planar maps with labelled vertices such that:

- the root edge is labelled from 0 to 1,
- adjacent labels differ by  $\pm 1$ .

# 1. From Eulerian orientations to labelled maps




The 4-valent case: labelled quadrangulations




*Labelled maps* are rooted planar maps with labelled vertices such that:

- the root edge is labelled from 0 to 1,
- adjacent labels differ by  $\pm 1$ .

## For the 2-vertex model




Alternating (weight  $t\omega$ )  
bicoloured face




Non-alternating (weight  $t$ )  
colourful face (3 labels)

## For the 2-vertex model




Alternating (weight  $t\omega$ )  
bicoloured face




Non-alternating (weight  $t$ )  
colourful face (3 labels)

### Colourful quadrangulation:





## 2. From colourful labelled quadrangulations to labelled maps





A bijection à la Cori-Vauquelin-Schaeffer [Miermont 09, Ambjörn & Budd 13]

## 2. From colourful labelled quadrangulations to labelled maps





A bijection à la Cori-Vauquelin-Schaeffer [Miermont 09, Ambjörn & Budd 13]

## 2. From colourful labelled quadrangulations to labelled maps





A bijection à la Cori-Vauquelin-Schaeffer [Miermont 09, Ambjörn & Budd 13]

## 2. From colourful labelled quadrangulations to labelled maps



A bijection à la Cori-Vauquelin-Schaeffer [Miermont 09, Ambjörn & Budd 13]

## 2. From colourful labelled quadrangulations to labelled maps




A bijection à la Cori-Vauquelin-Schaeffer [Miermont 09, Ambjörn & Budd 13]

### Corollary

The number of labelled maps with  $n$  edges is half the number of colourful quadrangulations with  $n$  faces.

## IV. Functional equations



# The recursive method

- One may have to
  - ▶ introduce more general objects (3 different classes of labelled maps)
  - ▶ introduce new enumeration parameters (2 of them)
- Solving can be non-trivial!

## A system for colourful quadrangulations [mbm, Elvey Price]

- There exists a unique 3-tuple of series, denoted  $P(t, y)$ ,  $C(t, x, y)$  and  $D(t, x, y)$ , belonging to  $\mathbb{Q}[[y, t]]$ ,  $\mathbb{Q}[x][[y, t]]$  and  $\mathbb{Q}[[x, y, t]]$ , such that:

$$P(t, y) = \frac{1}{y}[x^1]C(t, x, y),$$

$$D(t, x, y) = \frac{1}{1 - C\left(t, \frac{1}{1-x}, y\right)},$$

$$C(t, x, y) = xy[x^{\geq 0}] \left( P(t, tx)D\left(t, \frac{1}{x}, y\right) \right),$$

and  $P(t, 0) = 1$ .

## A system for colourful quadrangulations [mbm, Elvey Price]

- There exists a unique 3-tuple of series, denoted  $P(t, y)$ ,  $C(t, x, y)$  and  $D(t, x, y)$ , belonging to  $\mathbb{Q}[[y, t]]$ ,  $\mathbb{Q}[x][[y, t]]$  and  $\mathbb{Q}[[x, y, t]]$ , such that:

$$P(t, y) = \frac{1}{y}[x^1]C(t, x, y),$$

$$D(t, x, y) = \frac{1}{1 - C\left(t, \frac{1}{1-x}, y\right)},$$

$$C(t, x, y) = xy[x^{\geq 0}] \left( P(t, tx)D\left(t, \frac{1}{x}, y\right) \right),$$

and  $P(t, 0) = 1$ .

**Notation:** for a Laurent series  $F(x) = \sum_{i \in \mathbb{Z}} F_i x^i$ ,

$$[x^1]F(x) := F_1 \quad \text{and} \quad [x^{\geq 0}]F(x) := \sum_{i \geq 0} F_i x^i.$$

## A system for colourful quadrangulations [mbm, Elvey Price]

- There exists a unique 3-tuple of series, denoted  $P(t, y)$ ,  $C(t, x, y)$  and  $D(t, x, y)$ , belonging to  $\mathbb{Q}[[y, t]]$ ,  $\mathbb{Q}[x][[y, t]]$  and  $\mathbb{Q}[[x, y, t]]$ , such that:

$$P(t, y) = \frac{1}{y}[x^1]C(t, x, y),$$

$$D(t, x, y) = \frac{1}{1 - C\left(t, \frac{1}{1-x}, y\right)},$$

$$C(t, x, y) = xy[x^{\geq 0}] \left( P(t, tx)D\left(t, \frac{1}{x}, y\right) \right),$$

and  $P(t, 0) = 1$ .

- The generating function that counts colourful quadrangulations by faces is

$$Q^c(t) = [y^1]P(t, y) - 1.$$

## A system for colourful quadrangulations [mbm, Elvey Price]

- There exists a unique 3-tuple of series, denoted  $P(t, y)$ ,  $C(t, x, y)$  and  $D(t, x, y)$ , belonging to  $\mathbb{Q}[[y, t]]$ ,  $\mathbb{Q}[x][[y, t]]$  and  $\mathbb{Q}[[x, y, t]]$ , such that:

$$P(t, y) = \frac{1}{y}[x^1]C(t, x, y),$$

$$D(t, x, y) = \frac{1}{1 - C\left(t, \frac{1}{1-x}, y\right)},$$

$$C(t, x, y) = xy[x^{\geq 0}] \left( P(t, tx)D\left(t, \frac{1}{x}, y\right) \right),$$

and  $P(t, 0) = 1$ .

- The generating function that counts colourful quadrangulations by faces is


$$Q^c(t) = [y^1]P(t, y) - 1.$$

- Recall that  $Q^c(t) = 2G(t)$ , where  $G(t)$  counts Eulerian orientations by edges.


# The most interesting equation

$$C(t, x, y) = xy[x \geq 0] \left( P(t, tx) D \left( t, \frac{1}{x}, y \right) \right)$$

A contraction operation of a whole submap



## V. Solution: guess and check!



## Solution for colourful quadrangulations

- Let  $S \equiv S(t) = t - 2t^2 - 4t^3 - 20t^4 - 132t^5 - \dots$  be the unique power series with constant term 0 satisfying

$$t = \sum_{n=0}^{\infty} \frac{1}{n+1} \binom{2n}{n}^2 S^{n+1}.$$

- The generating function of planar Eulerian orientations counted by edges is

$$G(t) = Q^c(t)/2 = \frac{1}{4t^2}(t - 2t^2 - S)$$

## Solution for colourful quadrangulations

- Let  $S \equiv S(t) = t - 2t^2 - 4t^3 - 20t^4 - 132t^5 - \dots$  be the unique power series with constant term 0 satisfying

$$t = \sum_{n=0}^{\infty} \frac{1}{n+1} \binom{2n}{n}^2 S^{n+1}.$$

- There exists a unique 3-tuple of series, denoted  $P(t, y)$ ,  $C(t, x, y)$  and  $D(t, x, y)$ , belonging to  $\mathbb{Q}[[y, t]]$ ,  $\mathbb{Q}[x][[y, t]]$  and  $\mathbb{Q}[[x, y, t]]$ , such that:

$$P(t, y) = \frac{1}{y} [x^1] C(t, x, y),$$

$$D(t, x, y) = \frac{1}{1 - C\left(t, \frac{1}{1-x}, y\right)},$$

$$C(t, x, y) = xy [x^{\geq 0}] \left( P(t, tx) D\left(t, \frac{1}{x}, y\right) \right),$$

and  $P(t, 0) = 1$ .

## Solution for colourful quadrangulations

- Let  $S \equiv S(t) = t - 2t^2 - 4t^3 - 20t^4 - 132t^5 - \dots$  be the unique power series with constant term 0 satisfying

$$t = \sum_{n=0}^{\infty} \frac{1}{n+1} \binom{2n}{n}^2 S^{n+1}.$$

- The whole solution:

$$tP(t, ty) = \sum_{n \geq 0} \sum_{j=0}^n \frac{1}{n+1} \binom{2n}{n} \binom{2n-j}{n} y^j S^{n+1},$$

$$C(t, x, ty) = 1 - \exp \left( - \sum_{n \geq 0} \sum_{j=0}^n \sum_{i=0}^n \frac{1}{n+1} \binom{2n-i}{n} \binom{2n-j}{n} x^{i+1} y^{j+1} S^{n+1} \right)$$

$$D(t, x, ty) = \exp \left( \sum_{n \geq 0} \sum_{j=0}^n \sum_{i \geq 0} \frac{1}{n+1} \binom{2n-j}{n} \binom{2n+i+1}{n} x^i y^{j+1} S^{n+1} \right)$$

## A similar solution for all labelled quadrangulations

(with a more complicated system of functional equations)

- Let  $R(t) \equiv R = t - 3t^2 - 12t^3 - 105t^4 - 1206t^5 - \dots$ , be the unique formal power series with constant term 0 satisfying

$$t = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} \binom{3n}{n} R^{n+1}.$$

- Then the generating function of 4-valent Eulerian orientations, counted by vertices, is

$$\begin{aligned} Q(t) &= \frac{1}{3t^2} (t - 3t^2 - R(t)) \\ &= 4t + 35t^2 + 402t^3 + \dots \end{aligned}$$

- Asymptotics:

$$q_n \sim \frac{1}{18} \frac{(4\sqrt{3}\pi)^{n+2}}{n^2(\log n)^2}$$

## A similar solution for all labelled quadrangulations

(with a more complicated system of functional equations)

- Let  $R(t) \equiv R = t - 3t^2 - 12t^3 - 105t^4 - 1206t^5 - \dots$ , be the unique formal power series with constant term 0 satisfying

$$t = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} \binom{3n}{n} R^{n+1}.$$

- Then the generating function of 4-valent Eulerian orientations, counted by vertices, is

$$\begin{aligned} Q(t) &= \frac{1}{3t^2} (t - 3t^2 - R(t)) \\ &= 4t + 35t^2 + 402t^3 + \dots \end{aligned}$$

- Asymptotics:

$$q_n \sim \frac{1}{18} \frac{(4\sqrt{3}\pi)^{n+2}}{n^2(\log n)^2}$$

**How can one guess that?**

## Guess and check: the story

Let  $R = t - 3t^2 + \dots$  satisfy:

$$t = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} \binom{3n}{n} R^{n+1}.$$

Then the GF of labelled quadrangulations is:

$$Q(t) = \frac{1}{3t^2} (t - 3t^2 - R(t)).$$

## Guess and check: the story

Let  $R = t - 3t^2 + \dots$  satisfy:

$$t = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} \binom{3n}{n} R^{n+1}.$$

Then the GF of labelled quadrangulations is:

$$Q(t) = \frac{1}{3t^2} (t - 3t^2 - R(t)).$$

- [Elvey Price, Guttmann 18] a formidable system of functional equations
- about 100 coefficients  $q_n$  of  $Q(t)$
- guessed the growth constant:  $\lim q_n^{1/n} = 4\sqrt{3} \pi$

## Guess and check: the story

Let  $R = t - 3t^2 + \dots$  satisfy:

$$t = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} \binom{3n}{n} R^{n+1}.$$

Then the GF of labelled quadrangulations is:

$$Q(t) = \frac{1}{3t^2} (t - 3t^2 - R(t)).$$

- [Elvey Price, Guttmann 18] a formidable system of functional equations
- about 100 coefficients  $q_n$  of  $Q(t)$
- guessed the growth constant:  $\lim q_n^{1/n} = 4\sqrt{3}\pi$
- “oh, I’ve seen this constant in another map problem” [mbm-Courtiel 15]

## Guess and check: the story

Let  $R = t - 3t^2 + \dots$  satisfy:

$$t = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} \binom{3n}{n} R^{n+1}.$$



Then the GF of labelled quadrangulations is:

$$Q(t) = \frac{1}{3t^2} (t - 3t^2 - R(t)).$$

- [Elvey Price, Guttmann 18] a formidable system of functional equations
- about 100 coefficients  $q_n$  of  $Q(t)$
- guessed the growth constant:  $\lim q_n^{1/n} = 4\sqrt{3}\pi$
- “oh, I’ve seen this constant in another map problem” [mbm-Courtiel 15]

## Guess and check: the story

Let  $R = t - 3t^2 + \dots$  satisfy:

$$t = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} \binom{3n}{n} R^{n+1}.$$



Then the GF of labelled quadrangulations is:

$$Q(t) = \frac{1}{3t^2} (t - 3t^2 - R(t)).$$

- [Elvey Price, Guttmann 18] a formidable system of functional equations
- about 100 coefficients  $q_n$  of  $Q(t)$
- guessed the growth constant:  $\lim q_n^{1/n} = 4\sqrt{3}\pi$
- “oh, I’ve seen this constant in another map problem” [mbm-Courtiel 15]
- a (guessed) relation between the solutions of the two problems [Elvey Price]

# Guess and check: the story

Luck and Andrew EP

Let  $R = t - 3t^2 + \dots$  satisfy:

$$t = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} \binom{3n}{n} R^{n+1}.$$




Then the GF of labelled quadrangulations is:

$$Q(t) = \frac{1}{3t^2} (t - 3t^2 - R(t)).$$

- [Elvey Price, Guttmann 18] a formidable system of functional equations
- about 100 coefficients  $q_n$  of  $Q(t)$
- guessed the growth constant:  $\lim q_n^{1/n} = 4\sqrt{3}\pi$
- “oh, I’ve seen this constant in another map problem” [mbm-Courtiel 15]
- a (guessed) relation between the solutions of the two problems [Elvey Price]

## VI. The 6-vertex model

An additional weight  $\omega$  on alternating vertices of 4-valent maps



This was already solved... [Kostov, Nucl. Physics B 00]

This was already solved... [Kostov, Nucl. Physics B 00]

Recall: solution when  $\omega = 1$

Let  $R(t) \equiv R = t - 3t^2 - 12t^3 - 105t^4 - 1206t^5 - \dots$ , be the unique formal power series with constant term 0 satisfying

$$t = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} \binom{3n}{n} R^{n+1}.$$

Then the generating function of 4-valent Eulerian orientations, counted by vertices, is

$$Q(t) = \frac{1}{3t^2} (t - 3t^2 - R(t)).$$

This was already solved... [Kostov, Nucl. Physics B 00]

(Once digested/corrected by Paul Zinn-Justin)

Write  $\omega = -2 \cos(2\alpha)$ .

This was already solved... [Kostov, Nucl. Physics B 00]

(Once digested/corrected by Paul Zinn-Justin)

Write  $\omega = -2 \cos(2\alpha)$ . Let  $q(t, \omega) \equiv q = t + (6\omega + 6)t^2 + \dots$  satisfy:

$$t = \frac{\cos \alpha}{64 \sin^3 \alpha} \left( -\frac{\vartheta(\alpha, q)\vartheta'''(\alpha, q)}{\vartheta'(\alpha, q)^2} + \frac{\vartheta''(\alpha, q)}{\vartheta'(\alpha, q)} \right)$$

This was already solved... [Kostov, Nucl. Physics B 00]

(Once digested/corrected by Paul Zinn-Justin)

Write  $\omega = -2 \cos(2\alpha)$ . Let  $q(t, \omega) \equiv q = t + (6\omega + 6)t^2 + \dots$  satisfy:

$$t = \frac{\cos \alpha}{64 \sin^3 \alpha} \left( -\frac{\vartheta(\alpha, q)\vartheta'''(\alpha, q)}{\vartheta'(\alpha, q)^2} + \frac{\vartheta''(\alpha, q)}{\vartheta'(\alpha, q)} \right)$$

where

$$\vartheta(z, q) = 2 \sin(z) q^{1/8} \prod_{n=1}^{\infty} (1 - 2 \cos(2z) q^n + q^{2n})(1 - q^n).$$

# This was already solved... [Kostov, Nucl. Physics B 00]

(Once digested/corrected by Paul Zinn-Justin)

Write  $\omega = -2 \cos(2\alpha)$ . Let  $q(t, \omega) \equiv q = t + (6\omega + 6)t^2 + \dots$  satisfy:

$$t = \frac{\cos \alpha}{64 \sin^3 \alpha} \left( -\frac{\vartheta(\alpha, q)\vartheta'''(\alpha, q)}{\vartheta'(\alpha, q)^2} + \frac{\vartheta''(\alpha, q)}{\vartheta'(\alpha, q)} \right)$$

where

$$\vartheta(z, q) = 2 \sin(z) q^{1/8} \prod_{n=1}^{\infty} (1 - 2 \cos(2z) q^n + q^{2n})(1 - q^n).$$

• Define the series  $R(t, \omega)$  by

$$R(t, \omega) = \frac{\cos^2 \alpha}{96 \sin^4 \alpha} \frac{\vartheta(\alpha, q)^2}{\vartheta'(\alpha, q)^2} \left( -\frac{\vartheta'''(\alpha, q)}{\vartheta'(\alpha, q)} + \frac{\vartheta'''(0, q)}{\vartheta'(0, q)} \right).$$

# This was already solved... [Kostov, Nucl. Physics B 00]

(Once digested/corrected by Paul Zinn-Justin)

Write  $\omega = -2 \cos(2\alpha)$ . Let  $q(t, \omega) \equiv q = t + (6\omega + 6)t^2 + \dots$  satisfy:

$$t = \frac{\cos \alpha}{64 \sin^3 \alpha} \left( -\frac{\vartheta(\alpha, q)\vartheta'''(\alpha, q)}{\vartheta'(\alpha, q)^2} + \frac{\vartheta''(\alpha, q)}{\vartheta'(\alpha, q)} \right)$$

where

$$\vartheta(z, q) = 2 \sin(z) q^{1/8} \prod_{n=1}^{\infty} (1 - 2 \cos(2z) q^n + q^{2n})(1 - q^n).$$

- Define the series  $R(t, \omega)$  by

$$R(t, \omega) = \frac{\cos^2 \alpha}{96 \sin^4 \alpha} \frac{\vartheta(\alpha, q)^2}{\vartheta'(\alpha, q)^2} \left( -\frac{\vartheta'''(\alpha, q)}{\vartheta'(\alpha, q)} + \frac{\vartheta'''(0, q)}{\vartheta'(0, q)} \right).$$

- Then the generating function of quartic rooted planar Eulerian orientations, counted by vertices, with a weight  $\omega$  per alternating vertex is

$$Q(t, \omega) = \frac{1}{(\omega + 2)t^2} (t - (\omega + 2)t^2 - R(t, \omega)).$$

## Two questions

- Relate this to the MBM-EP solution when  $\omega = 0$  or  $\omega = 1$   
[Elvey Price & Zinn Justin 19+]
- Find a new proof of Kostov's result via our functional equations  
[mbm & Elvey Price 19+]

## Complements and perspectives

- record the **number of faces** in Eulerian orientations (equivalently, the number of oriented faces in 4-valent Eulerian orientations) — **done**
- record the label of a vertex in a pointed labelled map — **in progress**.  
Scaling and limit law?
- asymptotics and phase transitions when  $\omega$  varies

## Something to take home

Let  $R = t - 3t^2 + \dots$  satisfy:


$$t = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} \binom{3n}{n} R^{n+1}.$$

The GF of 4-valent Eulerian orientations is

$$Q(t) = \frac{1}{3t^2} (t - 3t^2 - R(t)).$$

### Another interpretation [mbm-Courtiel 15]

Consider ternary trees with black and white leaves. Define the **charge** at a node to be the number of white leaves minus the number of black leaves in the associated subtree. Then  $(t - R)/t = 3t(1 + Q)$  counts (by the number of nodes) trees in which the root is the **only** vertex of charge 1.




Bijection?

## Something to take home



### Another interpretation [mbm-Courtiel 15]

Consider ternary trees with black and white leaves. Define the **charge** at a node to be the number of white leaves minus the number of black leaves in the associated subtree. Then  $(t - R)/t = 3t(1 + Q)$  counts (by the number of nodes) trees in which the root is the **only** vertex of charge 1.



Bijection?