


# On the shape of binary trees

Mireille Bousquet-Mélou, CNRS, LaBRI, Bordeaux




ArXiv math.CO/0501266

ArXiv math.PR/0500322 (with Svante Janson)

<http://www.labri.fr/~bousquet>

## A complete binary tree




$n$  internal vertices, called nodes (size  $n$ )

$n + 1$  external vertices, called leaves



## A plane tree




Each node has a (possibly empty) **ordered sequence** of children

$n$  edges,  $n + 1$  nodes (size  $n$ )


What is the shape of a (large) *random* binary tree?

# The (horizontal) profile of a binary tree




A binary tree of height 4 (or 5...)

## The horizontal profile of a plane tree



A plane tree of height 3.

## NEW! The *vertical* profile of a binary tree



A binary tree of **right width** 3, and vertical profile  $[2, 2; 4, 2, 1, 1]$ .

**Why study the vertical profile of trees?**

**Why study the vertical profile of trees?**

**Wait and see!**

## Limit results on the shape of trees: general approach

- **Enumerative combinatorics** (decompositions of trees, recurrence relations, generating functions...)
- **Singularity analysis** [Flajolet-Odlyzko 90]: extract from the generating functions the asymptotic behaviour of their coefficients


## Decomposition and enumeration of binary trees

Let  $a_n$  be the number of binary trees with  $n$  nodes (internal vertices):

$$\begin{cases} a_0 = 1 \\ a_n = \sum_{m=0}^{n-1} a_m a_{n-m-1} \end{cases}$$

Let  $A(t) := \sum_{n \geq 0} a_n t^n$  be the associated **generating function**:

$$A(t) = 1 + tA(t)^2.$$




## Decomposition and enumeration of plane trees

Let  $a_n^*$  be the number of plane trees with  $n$  edges ( $n + 1$  vertices):

$$\begin{cases} a_0^* &= 1 \\ a_n^* &= \sum_{m=0}^{n-1} a_m^* a_{n-m-1}^* \end{cases}$$

Let  $A^*(t) := \sum_{n \geq 0} a_n^* t^n$  be the associated **generating function**:

$$A^*(t) = 1 + tA^*(t)^2.$$



## The Catalan numbers

There are as many **binary trees** with  $n$  nodes as **plane trees** with  $n$  edges:

$$a_n = a_n^*$$

The associated generating function is

$$A(t) = A^*(t) = \sum_n a_n t^n$$

It satisfies


$$A(t) = 1 + tA(t)^2 \Rightarrow A(t) = \frac{1 - \sqrt{1 - 4t}}{2t} = \sum_{n \geq 0} \frac{1}{n+1} \binom{2n}{n} t^n.$$

Hence

$$a_n = a_n^* = \frac{1}{n+1} \binom{2n}{n}$$

is the  $n$ th **Catalan number**.

# Part 1. Height and width of binary trees




## Counting trees of bounded height

Let  $H_{\leq j}(t) \equiv H_{\leq j}$  be the generating function of binary trees of height at most  $j$ :

$$H_{\leq j} = \sum_{n \geq 0} h_{n, \leq j} t^n$$

$$H_{\leq 0} = 1$$


$$H_{\leq j} = 1 + tH_{\leq j-1}^2$$



Let  $H_{\leq j}^*(t) \equiv H_{\leq j}^*$  be the generating function of plane trees of height at most  $j$ , where  $t$  counts edges.


$$H_{\leq 0}^* = 1$$

$$H_{\leq j}^* = 1 + tH_{\leq j-1}^* H_{\leq j}^*$$





## Counting trees of bounded (right) width

Let  $W_{\leq j}(t) \equiv W_{\leq j}$  be the generating function of binary trees of right width at most  $j$ .



	$W_{\leq -1} = 1$
For $j \geq 0$ ,	$W_{\leq j} = 1 + tW_{\leq j+1}W_{\leq j-1}$

## Trees of bounded height or bounded width: functional equations

Family of trees	Bounded height	Bounded (right) width
	$H_{\leq 0} = 1$ $H_{\leq j} = 1 + tH_{\leq j-1}^2$	$W_{\leq -1} = 1$ $W_{\leq j} = 1 + tW_{\leq j+1}W_{\leq j-1}$ $(j \geq 0)$
	$H_{\leq 0}^* = 1$ $H_{\leq j}^* = 1 + tH_{\leq j-1}^*H_{\leq j}^*$	

## Generating functions for trees of bounded height/width

**Proposition:** The generating function of plane trees of height  $\leq j$  is:

$$H_{\leq j}^* = A \frac{1 - Q^{j+1}}{1 - Q^{j+2}}$$

where  $A$  counts plane trees and  $Q = A - 1$  [de Bruijn, Knuth, Rice 72].

## Generating functions for trees of bounded height/width

**Proposition:** The generating function of plane trees of height  $\leq j$  is:

$$H_{\leq j}^* = A \frac{1 - Q^{j+1}}{1 - Q^{j+2}}$$

where  $A$  counts plane trees and  $Q = A - 1$  [de Bruijn, Knuth, Rice 72].



**Proposition** [mbm 06]: The generating function of binary trees of right width  $\leq j$  is:

$$W_{\leq j} = A \frac{(1 - Z^{j+2})(1 - Z^{j+7})}{(1 - Z^{j+4})(1 - Z^{j+5})},$$

where  $A$  counts binary trees and  $Z \equiv Z(t)$  is the unique series in  $t$  such that

$$Z = t \frac{(1 + Z^2)^2}{1 - Z + Z^2} \quad \text{and} \quad Z(0) = 0.$$

**But. . . w h y ?**

## Limit results on the shape of trees: general approach

- **Enumerative combinatorics**: Using a recursive decomposition of trees, write equations for the relevant generating functions and **solve** them...
- Use the results (or the technique) of **singularity analysis** [Flajolet-Odlyzko 90] to extract from these series the asymptotic behaviour of their coefficients
  - Results: “If  $B(t) = \sum_n b_n t^n$  behaves like this in the neighborhood of its dominant singularity  $t_s$ , then its coefficients  $b_n$  behave asymptotically like that.”
  - Technique: Cauchy’s formula


$$b_n = \frac{1}{2i\pi} \int_{\mathcal{C}_n} B(t) \frac{dt}{t^{n+1}}$$

for a carefully chosen contour  $\mathcal{C}_n$ .

## The average height of binary trees: experimental approach

Consider the **average height** of plane trees of size  $n$ :

$$\mathbb{E}(H_n^*) := \frac{1}{a_n} \sum_{|\tau|=n} h(\tau) = \frac{1}{a_n} \sum_{j \geq 0} j (h_{n, \leq j}^* - h_{n, \leq j-1}^*).$$




## Convergence in law of the height: experimental approach

The number of trees of size  $n$  and height at most  $j$  is  $h_{n, \leq j}^*$ . For  $x \geq 0$ , consider

$$F_n(x) := \mathbb{P} \left( \frac{H_n^*}{\sqrt{n}} \leq x \right) = \frac{1}{a_n} h_{n, \leq x\sqrt{n}}^*,$$

the **proportion** of trees of size  $n$  having height at most  $x\sqrt{n}$ . It is the **probability** that a random tree of size  $n$  has height at most  $x\sqrt{n}$ .

Graph of  $F_n(x)$  :




## Limit results for the height of trees

**Convergence in law:** For all  $x \geq 0$ ,

$$\mathbb{P} \left( \frac{H_n^*}{\sqrt{n}} \leq x \right) \rightarrow F(x),$$

where

$$\begin{aligned} F(x) &= -\frac{1}{i\sqrt{\pi}} \int_{\mathcal{H}} \coth(x\sqrt{-z}) \sqrt{-z} e^{-z} dz \\ &= \sum_{k=-\infty}^{+\infty} e^{-k^2 x^2} (1 - 2k^2 x^2) \end{aligned}$$



**Convergence of the moments:** Let  $H_n^*$  be the (random) height of a plane tree with  $n$  edges. As  $n \rightarrow \infty$ ,

$$\frac{\mathbb{E}(H_n^*)}{\sqrt{n}} \rightarrow \sqrt{\pi}, \quad \mathbb{E} \left( \left( \frac{H_n^*}{\sqrt{n}} \right)^k \right) \rightarrow k(k-1)\Gamma(k/2)\zeta(k) \quad \text{for } k \geq 2.$$

[Flajolet-Odlyzko 82], [Brown-Schubert 84]

+ similar results for binary trees

## Limit results for the right width of binary trees [mbm 06]

**Convergence of the moments:** Let  $W_n$  be the (random) right width of a binary tree with  $n$  nodes. Then, as  $n \rightarrow \infty$ ,


$$\frac{\mathbb{E}(W_n)}{n^{1/4}} \rightarrow \frac{3\sqrt{\pi}}{\sqrt{2}\Gamma(3/4)}, \quad \mathbb{E}\left(\left(\frac{W_n}{n^{1/4}}\right)^2\right) \rightarrow 6\sqrt{\pi},$$

$$\mathbb{E}\left(\left(\frac{W_n}{n^{1/4}}\right)^k\right) = \frac{24\sqrt{\pi}k!\zeta(k-1)}{\sqrt{2}^k\Gamma((k-2)/4)} \quad \text{for } k \geq 3.$$


**Convergence in law:**

$$\mathbb{P}\left(\frac{W_n}{n^{1/4}} \geq x\right) \rightarrow G(x)$$


$$G(x) = \frac{3}{i\sqrt{\pi}} \int_{\mathcal{H}} \frac{\sqrt{-z}e^{-z}}{\sinh^2(x(-z^{1/4}/\sqrt{2}))} dz$$




## Height and width of binary trees




## Part 2. What about the profiles?



## Gallery of (horizontal and vertical) profiles




Horizontal profiles of random binary trees with 1000 nodes.




Vertical profiles of random binary trees with 1000 nodes.

## The average (horizontal and vertical) profiles

Average horizontal profile of plane trees of size 10, 20, 30, 40, 50:



Average vertical profile of binary trees of size 5, 10, 20:



## One new ingredient: bivariate generating functions

**Before:** Given  $j$ , how many plane trees of size  $n$  have height at most  $j$ ?

$$\Rightarrow \text{series } H_{\leq j}^*(t) = \sum_n h_{n, \leq j} t^n$$

**Now:** Given  $j$ , how many plane trees of size  $n$  have exactly  $k$  nodes at height  $j$ ?


$$\Rightarrow \text{series } P_j^*(t, u) = \sum_{n, k} p_{n, k, j} t^n u^k$$

## The number of nodes at height $j$ (plane trees)

Let  $P_j^* \equiv P_j^*(t, u)$  be the generating function of plane trees, counted by edges (variable  $t$ ) and by the number of nodes at height  $j$  (variable  $u$ ):

$$P_0^* = uA(t)$$

$$P_j^* = 1 + tP_{j-1}^*P_j^*$$



**Proposition** [???:

$$P_j^* = A \frac{1 - MQ^j}{1 - MQ^{j+1}}$$

where  $A$  counts plane trees,  $Q = A - 1$  and

$$M = \frac{A - u - tuA^2}{u + A(1 - u) + tuA^2(1 - A)}.$$


## The number of nodes at abscissa $j$ (binary trees)

Let  $V_j \equiv V_j(t, u)$  be the generating function of binary trees, counted by nodes ( $t$ ) and by the number of nodes at abscissa  $j$  (variable  $u$ ).

$$V_0 = 1 + utV_1^2$$

For  $j \geq 1$ ,

$$V_j = 1 + tV_{j+1}V_{j-1}$$



**Proposition** [mbm 06]: the series  $V_j$  are algebraic. Moreover,

$$V_j = A \frac{(1 + MZ^j)(1 + MZ^{j+5})}{(1 + MZ^{j+2})(1 + MZ^{j+3})}$$

where  $A$  counts binary trees,

$$Z = t \frac{(1 + Z^2)^2}{1 - Z + Z^2},$$

and  $M \equiv M(t, u)$  is the unique power series in  $t$  such that

$$M = (u - 1) \frac{Z(1 + MZ)^2(1 + MZ^2)(1 + MZ^6)}{(1 + Z)^2(1 + Z + Z^2)(1 - Z)^3(1 - M^2Z^5)}.$$

## The horizontal profile

A tree of size  $n$  has, on average, height  $O(\sqrt{n})$ . How many nodes are there at a given (horizontal) level?

???

## The horizontal profile

A tree of size  $n$  has, on average, height  $O(\sqrt{n})$ . How many nodes are there at a given (horizontal) level?

About  $\sqrt{n}$ .

## The horizontal profile

A tree of size  $n$  has, on average, height  $O(\sqrt{n})$ . How many nodes are there at a given (horizontal) level?

About  $\sqrt{n}$ .

Let  $X_n^*(j)$  be the number of nodes located at height  $j$  in a random plane tree with  $n$  edges.

We study the quantity

$$\frac{X_n^*([\lambda\sqrt{n}])}{\sqrt{n}}.$$

## The vertical profile

A binary tree of size  $n$  has, on average, width  $O(n^{1/4})$ . How many nodes are there on a given (vertical) layer?

About  $n^{3/4}$ .

Let  $Y_n(j)$  be the number of nodes located at abscissa  $j$  in a random binary tree with  $n$  nodes.

We study the quantity

$$\frac{Y_n(\lfloor \lambda n^{1/4} \rfloor)}{n^{3/4}}.$$


## The horizontal profile of plane trees

- The sequence  $\frac{X_n^*([\lambda\sqrt{n}])}{\sqrt{n}}$  converges in law for each  $\lambda$  (and as a process [Drmot-Gittenberger 97] )

- The first moment:

$$\mathbb{E} \left( \frac{X_n^*([\lambda\sqrt{n}])}{\sqrt{n}} \right) \rightarrow 2\lambda e^{-\lambda^2}.$$

On average, there are about  $2\lambda e^{-\lambda^2} \sqrt{n}$  nodes at height  $[\lambda\sqrt{n}]$  in a plane tree having  $n$  edges.



In other words, the height of a random node in a random tree, once normalized by  $\sqrt{n}$ , follows a law of density  $2\lambda e^{-\lambda^2}$ .

## The vertical profile of binary trees


Let  $Y_n(j)$  be the (random) number of nodes at abscissa  $j$  in a binary tree having  $n$  nodes.

- The sequence  $\frac{Y_n(\lfloor \lambda n^{1/4} \rfloor)}{n^{3/4}}$  **converges in law** to a random variable  $Y(\lambda)$  described explicitly by its Laplace transform for every  $\lambda$  (and as a process [mbm-Janson 06])

- Moreover,

$$\mathbb{E} \left( \frac{Y_n(\lfloor \lambda n^{1/4} \rfloor)}{n^{3/4}} \right) \rightarrow \frac{1}{\sqrt{2\pi}} \sum_{m \geq 0} \frac{(-\sqrt{2}|\lambda|)^m}{m!} \cos \frac{(m+1)\pi}{4} \Gamma \left( \frac{m+3}{4} \right).$$

This gives the average number of nodes at abscissa  $\lfloor \lambda n^{1/4} \rfloor$  in a random binary tree having  $n$  nodes.



[mbm 06]

## An example: nodes at abscissa 0

- The random variable

$$\frac{Y_n(0)}{n^{3/4}},$$

which gives the (normalized) number of nodes at abscissa 0, converges in law to a variable  $Y(0)$  such that

$$E(Y(0)^k) = \left(\frac{\sqrt{2}}{3}\right)^k \frac{\Gamma(1 + 3k/4)}{\Gamma(1 + k/2)}.$$

Hence (...)  $Y(0) = \frac{\sqrt{2}}{3\sqrt[3]{T_{2/3}}}$ , where  $T_{2/3}$  follows a unilateral stable law of parameter  $2/3$ .

$$E(e^{-aT_{2/3}}) = e^{-a^{2/3}} \text{ for } a \geq 0.$$

Merci Alain Rouault !

## An example: nodes at abscissa 0

- The random variable

$$\frac{Y_n(0)}{n^{3/4}},$$

which gives the (normalized) number of nodes at abscissa 0, converges in law to a variable  $Y(0)$  such that

$$E(Y(0)^k) = \left(\frac{\sqrt{2}}{3}\right)^k \frac{\Gamma(1 + 3k/4)}{\Gamma(1 + k/2)}.$$

Hence (...)  $Y(0) = \frac{\sqrt{2}}{3\sqrt[3]{T_{2/3}}}$ , where  $T_{2/3}$  follows a unilateral stable law of parameter  $2/3$ .

$$E(e^{-aT_{2/3}}) = e^{-a^{2/3}} \text{ for } a \geq 0.$$

Merci **Alain Rouault** !

- The number of nodes lying at a positive abscissa, normalized by  $n$ , converges to  $U(0, 1)$  [**Aldous**].



## Why study embedded trees?

- They have interesting combinatorial properties (mysterious algebraic series)

## Why study embedded trees?

- They have interesting combinatorial properties (mysterious algebraic series)

The GF of embedded plane trees with increments  $\pm 1$  having largest label at most  $j$  is


$$T_j = T \frac{(1 - Z^{j+1})(1 - Z^{j+5})}{(1 - Z^{j+2})(1 - Z^{j+4})},$$

where  $T = 1 + 2tT^2$  and

$$Z = t \frac{(1 + Z)^4}{1 + Z^2}.$$


## Why study embedded trees?

- They have interesting combinatorial properties (mysterious algebraic series)
- Embedded trees with **nonnegative** labels are related **bijectionally** to planar maps [Cori-Vauquelin 81], [Chassaing-Schaeffer 04], [Del Lungo, Del Ristoro, Penaud 00], [Bouttier-Di Francesco-Guitter 03]



## Why study embedded trees?


- They have interesting combinatorial properties (mysterious algebraic series)
- Embedded trees with **nonnegative** labels are related **bijectionally** to planar maps [Cori-Vauquelin 81], [Chassaing-Schaeffer 04], [Del Lungo, Del Ristoro, Penaud 00], [Bouttier-Di Francesco-Guitter 03]



- They are related to ISE (the Integrated SuperBrownian Excursion) [Aldous 93], [Borgs et al. 99]

## The Integrated SuperBrownian Excursion

The ISE is a (random) probability distribution on  $\mathbb{R}^d$  that occurs (almost) everywhere. At least, as soon as a branching structure (tree) is combined with an embedding of the nodes in the space. The ISE describes how the space is occupied by the nodes [Aldous 93], [Marckert-Mokkadem 03].





## Embedded trees and ISE

- Let  $T_n$  denote a random embedded tree with  $n$  nodes. Let  $\mu_n$  be the occupation measure of  $T_n$ :

$$\mu_n = \sum_{v \in T_n} \delta(\ell(v)),$$

where  $\ell(v)$  denotes the label (position) of the node  $v$ , and  $\delta(x)$  is the Dirac measure at  $x$ .





## Embedded trees and ISE

- Let  $T_n$  denote a random embedded tree with  $n$  nodes. Let  $\mu_n$  be the **occupation measure** of  $T_n$ :

$$\mu_n = \frac{1}{n} \sum_{v \in T_n} \delta(\ell(v)),$$

where  $\ell(v)$  denotes the label (position) of the node  $v$ , and  $\delta(x)$  is the Dirac measure at  $x$ . Then  $\mu_n$  is a **probability distribution** (total weight 1).





## Embedded trees and ISE

- Let  $T_n$  denote a random embedded tree with  $n$  nodes. Let  $\mu_n$  be the **occupation measure** of  $T_n$ :

$$\mu_n = \frac{1}{n} \sum_{v \in T_n} \delta(\ell(v)n^{-1/4}),$$

where  $\ell(v)$  denotes the label (position) of the node  $v$ , and  $\delta(x)$  is the Dirac measure at  $x$ . Then  $\mu_n$  is a **probability distribution** (total weight 1).

It is **random** since it depends on the **random tree**  $T_n$ .



## Embedded trees and ISE

- Let  $T_n$  denote a random embedded tree with  $n$  nodes. Let  $\mu_n$  be the occupation measure of  $T_n$ :



$$\mu_n = \frac{1}{n} \sum_{v \in T_n} \delta(\ell(v)n^{-1/4}),$$

where  $\ell(v)$  denotes the label (position) of the node  $v$ , and  $\delta(x)$  is the Dirac measure at  $x$ .

- Thm.** As  $n$  grows,  $\mu_n \rightarrow \mu_{\text{ise}}$ , where  $\mu_{\text{ise}}$  is the ISE.


[Aldous 93, Borgs et al. 99, Janson-Marckert 04]

This holds for “many” families of random embedded trees!





# Main objective: study ISE (in 1D) via embedded trees



- Largest label  $\rightarrow$  law of the **maximum of the support** of the ISE
- Number of nodes at abscissa  $j = \lambda n^{1/4} \rightarrow$  **law of the density** of the ISE at a fixed point  $\lambda$
- Number of nodes at abscissa  $\leq j = \lambda n^{1/4} \rightarrow$  **law of the distribution function** of the ISE at a fixed point  $\lambda$

## A beautiful slide for Wjch.

- Let  $M_i$  be the  $i$ th moment of the ISE ( $M_i$  is a random variable!)

$$M_i = \lim_n \frac{1}{n} \sum_{v \in T_n} (\ell(v) n^{-1/4})^i$$

## A beautiful slide for Wjcch.

- Let  $M_i$  be the  $i$ th moment of the ISE ( $M_i$  is a random variable!)

$$M_i = \lim_n \frac{1}{n} \sum_{v \in T_n} (\ell(v) n^{-1/4})^i$$

- Then  $\mathbb{E}(M_1^{2k}) = 0$  and  $\mathbb{E}(M_1^{2k}) = \frac{a_k \Gamma(1/2)}{2^{k/2} \Gamma((5k-1)/2)}$ , where  $a_0 = -2$  and for  $k > 0$ ,

$$4 a_k = \sum_{i=1}^{k-1} \binom{2k}{2i} a_i a_{k-i} + k(2k-1)(5k-4)(5k-6) a_{k-1}$$

## A beautiful slide for Wjcch.

- Let  $M_i$  be the  $i$ th moment of the ISE ( $M_i$  is a random variable!)

$$M_i = \lim_n \frac{1}{n} \sum_{v \in T_n} (\ell(v) n^{-1/4})^i$$

- Then  $\mathbb{E}(M_1^{2k}) = 0$  and  $\mathbb{E}(M_1^{2k}) = \frac{a_k \Gamma(1/2)}{2^{k/2} \Gamma((5k-1)/2)}$ , where  $a_0 = -2$  and for  $k > 0$ ,

$$4 a_k = \sum_{i=1}^{k-1} \binom{2k}{2i} a_i a_{k-i} + k(2k-1)(5k-4)(5k-6) a_{k-1}$$

- Also,


$$\mathbb{E}(M_1^{2k} M_2^\ell) = \frac{a_{k,\ell} \Gamma(1/2)}{2^{(k+\ell)/2} \Gamma((5k+3\ell-1)/2)},$$

where  $a_{0,0} = -2$  and the  $a_{k,\ell}$  are determined by induction on  $k + \ell$ :

$$a_{k,\ell} = \frac{1}{4} \sum_{(0,0) < (i,j) < (k,\ell)} \binom{2k}{2i} \binom{\ell}{j} a_{i,j} a_{k-i,\ell-j} + 2\ell(\ell-1) a_{k+1,\ell-2}$$

$$+ \frac{1}{4} k(2k-1)(5k+3\ell-4)(5k+3\ell-6) a_{k-1,\ell} + \frac{1}{2} (4k+1)\ell(5k+3\ell-4) a_{k,\ell-1}.$$

**That's it!**



## The Laplace transform of $Y(\lambda)$

Let  $\lambda \geq 0$ . The sequence  $Y_n(\lfloor \lambda n^{1/4} \rfloor)$  converges in distribution to a non-negative random variable  $Y(\lambda)$  whose Laplace transform is given, for  $|a| < 4/\sqrt{3}$ , by

$$\mathbb{E} \left( e^{aY(\lambda)} \right) = L(\lambda, a)$$

where

$$L(\lambda, a) = 1 + \frac{48}{i\sqrt{\pi}} \int_{\Gamma} \frac{A(a/v^3)e^{-2\lambda v}}{(1 + A(a/v^3)e^{-2\lambda v})^2} v^5 e^{v^4} dv,$$

$A(x) \equiv A$  is the unique solution of

$$A = \frac{x}{24} \frac{(1 + A)^3}{1 - A}$$

satisfying  $A(0) = 0$ , and the integral is taken over

$$\Gamma = \{1 - te^{-i\pi/4}, t \in (\infty, 0]\} \cup \{1 + te^{-i\pi/4}, t \in [0, \infty)\}.$$

## Did you say "ISE"?

