# Power domination in triangular grids


# Prosenjit Bose<sup>1</sup>, Claire Pennarun<sup>2</sup>, Sander Verdonschot<sup>1</sup>


School of Computer Science, Carleton University


LaBRI, University of Bordeaux


CCCG, July 26, 2017


Claire Pennarun








































Sets of vertices: S = selected, M = monitored





The **power domination number**  $\gamma_P(G)$  of *G* is the minimum size of *S* s.t. M = V(G) at the end.

### Power domination: some results

Background story: Placing a minimal number of "sensors" to monitor an electrical system


Power dominating set

**Input**:  $G, k \ge 0$ . **Question**: Is there a power dominating set  $S \subseteq V(G)$  with  $|S| \le k$ ? is NP-complete even for planar graphs, bipartite graphs, chordal graphs... [Guo et al. '05]

Well-studied classes: regular graphs and grids Hexagonal grids [Ferrero et al. '11], some products of paths [Dorfling and Henning '06, Dorbec et al. '08]...

## Power domination of triangular grids

**Triangular grid**  $T_k$ : regular triangular grid with border of hexagonal shape (*k* vertices on each side)




Theorem [Bose, P., Verdonschot 17+]


$$\gamma_P(T_k) = \left\lceil \frac{k}{3} \right\rceil$$

Upper bound: simple construction depending on the value of *k* modulo 3.


Lower bound: bound an invariant (the number of vertices potentially able to propagate at each step)

#### Monitoring set $\ensuremath{\mathbb{T}}$ is sufficient








#### Monitoring set $\ensuremath{\mathbb{T}}$ is sufficient





#### Monitoring set $\ensuremath{\mathbb{T}}$ is sufficient
















































### Monitoring set $\ensuremath{\mathbb{T}}$ is sufficient






Claire Pennarun








Claire Pennarun


General idea: study the size of the **border** of the set of monitored vertices (= vertices that could potentially propagate in the future)

Border  $B_A$  of a set A: vertices of A such that  $N(v) \setminus A \neq \emptyset$ 




General idea: study the size of the **border** of the set of monitored vertices (= vertices that could potentially propagate in the future)


Border  $B_A$  of a set A: vertices of A such that  $N(v) \setminus A \neq \emptyset$ 




**Shifting operation** in one direction *A* to *A*' (here direction  $\leftarrow$ )



**Shifting operation** in one direction *A* to *A*' (here direction  $\leftarrow$ )




**Shifting operation** in one direction *A* to *A*' (here direction  $\leftarrow$ )



Shifting does not increase the size of the border:  $|B_{A^*}| \le |B_A|$ .


**Shifting operation** in one direction *A* to *A*' (here direction  $\leftarrow$ )



Shifting does not increase the size of the border:  $|B_{A^*}| \le |B_A|$ .


Claire Pennarun

**Shifting operation** in one direction *A* to *A*' (here direction  $\leftarrow$ )



Shifting does not increase the size of the border:  $|B_{A^*}| \le |B_A|$ .


After shifting in directions  $\{\leftarrow, \checkmark, \land\}$  until stabilization,  $A^*$  has a staircase shape





# At any time, $|B_M| \le 6|S|$ (by induction on the propagation steps)


At any time,  $|B_M| \le 6|S|$  (by induction on the propagation steps) Consider *M* once it has  $\frac{n}{2}$  vertices. Shift  $M \to M^*$ 


At any time,  $|B_M| \le 6|S|$  (by induction on the propagation steps) Consider *M* once it has  $\frac{n}{2}$  vertices. Shift  $M \to M^*$ Each horizontal line has at least one vertex in  $B_{M^*}$ 














At any time,  $|B_M| \le 6|S|$  (by induction on the propagation steps) Consider *M* once it has  $\frac{n}{2}$  vertices. Shift  $M \to M^*$ Each horizontal line has at least one vertex in  $B_{M^*}$ 




At any time,  $|B_M| \le 6|S|$  (by induction on the propagation steps) Consider *M* once it has  $\frac{n}{2}$  vertices. Shift  $M \to M^*$ Each horizontal line has at least one vertex in  $B_{M^*}$ 




At any time,  $|B_M| \le 6|S|$  (by induction on the propagation steps) Consider *M* once it has  $\frac{n}{2}$  vertices. Shift  $M \to M^*$ Each horizontal line has at least one vertex in  $B_{M^*}$ 




At any time,  $|B_M| \le 6|S|$  (by induction on the propagation steps) Consider *M* once it has  $\frac{n}{2}$  vertices. Shift  $M \to M^*$ Each horizontal line has at least one vertex in  $B_{M^*}$ 




At any time,  $|B_M| \le 6|S|$  (by induction on the propagation steps) Consider *M* once it has  $\frac{n}{2}$  vertices. Shift  $M \to M^*$ Each horizontal line has at least one vertex in  $B_{M^*}$ 



At any time,  $|B_M| \le 6|S|$  (by induction on the propagation steps) Consider *M* once it has  $\frac{n}{2}$  vertices. Shift  $M \to M^*$ Each horizontal line has at least one vertex in  $B_{M^*}$ 




 $\rightarrow |B_{M^*}| \ge 2k - 1$  and  $|B_M| \ge |B_{M^*}|$ 



$$\rightarrow |B_{M^*}| \ge 2k - 1 \text{ and } |B_M| \ge |B_{M^*}|$$
$$\rightarrow |B_{M^*}| \ge 2k - 1$$

At any time,  $|B_M| \le 6|S|$  (by induction on the propagation steps) Consider *M* once it has  $\frac{n}{2}$  vertices. Shift  $M \to M^*$ Each horizontal line has at least one vertex in  $B_{M^*}$ 



$$2k-1 \le |B_M| \le 6|S| \to |S| \ge \frac{2k-1}{6}.$$

# Result


$$\frac{2k-1}{6} \le \gamma_P(T_k) \le \lceil \frac{k}{3} \rceil.$$
  

$$\gamma_P(T_k) \text{ is an integer + there are no integers between } \frac{2k-1}{6} \text{ and } \lceil \frac{k}{3} \rceil.$$
  

$$\rightarrow \boxed{\gamma_P(T_k) = \lceil \frac{k}{3} \rceil}.$$


#### Relation with the hexagonal grid

**Hexagonal grid**  $H_{\ell}$  with dimension  $\ell$  (number of hexagons per side) is an induced subgraph of  $T_{2\ell}$ ...



### Relation with the hexagonal grid

**Hexagonal grid**  $H_{\ell}$  with dimension  $\ell$  (number of hexagons per side) is an induced subgraph of  $T_{2\ell}$ ...



... and  $\gamma_P(H_\ell) = \gamma_P(T_{2\ell})$ 


Claire Pennarun

Power domination in triangular grids

CCCG, July 26, 2017 11 / 12

## Relation with the hexagonal grid

**Hexagonal grid**  $H_{\ell}$  with dimension  $\ell$  (number of hexagons per side) is an induced subgraph of  $T_{2\ell}$ ...



... and  $\gamma_P(H_\ell) = \gamma_P(T_{2\ell})$ When/why does this happen? Look at induced regular grids?

Claire Pennarun

# Thank you!

Claire Pennarun

Power domination in triangular grids

CCCG, July 26, 2017 12 / 12