

On the construction of an UBA

Jérôme Leroux

LSV - CNRS & ENS de Cachan


General case

Let $b \in \{0, \dots, r - 1\}$ be a **digit** where $r \geq 2$ is the basis of decomposition.

Let $\gamma_b : \mathbb{N}^m \rightarrow \mathbb{N}^m$ be the affine function defined by:

$$\gamma_b(x_1, \dots, x_m) = (x_2, \dots, x_m, r \cdot x_1 + b)$$

An infinite graph is associated to any subset $X \subseteq \mathbb{N}^m$:



General case

Let $b \in \{0, \dots, r - 1\}$ be a **digit** where $r \geq 2$ is the basis of decomposition.

Let $\gamma_b : \mathbb{N}^m \rightarrow \mathbb{N}^m$ be the affine function defined by:

$$\gamma_b(x_1, \dots, x_m) = (x_2, \dots, x_m, r \cdot x_1 + b)$$

Case $m = 2, r = 2$.

Let $X = \{(x_1, x_2) \in \mathbb{N}^2; \alpha x_1 + \beta x_2 = \gamma\}$, we have:

$$\begin{aligned} (x'_1, x'_2) \in \gamma_b^{-1}(X) &\iff \gamma_b(x'_1, x'_2) \in X \\ &\iff (x'_2, 2x'_1 + b) \in X \\ &\iff \alpha x'_2 + \beta(2x'_1 + b) = \gamma \\ &\iff 2\beta x'_1 + \alpha x'_2 = \gamma - b\beta \end{aligned}$$

General case

Let $b \in \{0, \dots, r - 1\}$ be a **digit** where $r \geq 2$ is the basis of decomposition.

Let $\gamma_b : \mathbb{N}^m \rightarrow \mathbb{N}^m$ be the affine function defined by:

$$\gamma_b(x_1, \dots, x_m) = (x_2, \dots, x_m, r \cdot x_1 + b)$$

Case $m = 2$, $r = 2$.

Let $X = \{(x_1, x_2) \in \mathbb{N}^2; \alpha x_1 + \beta x_2 = \gamma\}$, we have:

$$\begin{aligned} (x'_1, x'_2) \in \gamma_b^{-1}(X) &\iff \gamma_b(x'_1, x'_2) \in X \\ &\iff (x'_2, 2x'_1 + b) \in X \\ &\iff \alpha x'_2 + \beta(2x'_1 + b) = \gamma \\ &\iff 2\beta x'_1 + \alpha x'_2 = \gamma - b\beta \end{aligned}$$

$$\boxed{\alpha x_1 + \beta x_2 = \gamma} \xrightarrow{b} \boxed{2\beta x_1 + \alpha x_2 = \gamma - b\beta}$$

Case $2x_1 - x_2 = 0$

$$\boxed{\alpha x_1 + \beta x_2 = \gamma} \xrightarrow{b} \boxed{2\beta x_1 + \alpha x_2 = \gamma - b\beta}$$

$$\boxed{2x_1 - x_2 = 0}$$

Case $2x_1 - x_2 = 0$

$$\boxed{\alpha x_1 + \beta x_2 = \gamma} \xrightarrow{b} \boxed{2\beta x_1 + \alpha x_2 = \gamma - b\beta}$$

$$\begin{array}{c} \boxed{-2x_1 + 2x_2 = 1} \\ \uparrow \\ \boxed{2x_1 - x_2 = 0} \end{array}$$

Case $2x_1 - x_2 = 0$


$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$

$$\begin{aligned} -2x_1 + 2x_2 &= 1 \\ \iff (x_1, x_2) &\in \emptyset \\ 2x_1 - x_2 &= 0 \end{aligned}$$

↑
1


Case $2x_1 - x_2 = 0$

$$(\alpha x_1 + \beta x_2 = \gamma) \xrightarrow{b} (2\beta x_1 + \alpha x_2 = \gamma - b\beta)$$




Case $2x_1 - x_2 = 0$

$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$




Case $2x_1 - x_2 = 0$

$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$




Case $2x_1 - x_2 = 0$

$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$




Case $2x_1 - x_2 = 0$

$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$




Case $2x_1 - x_2 = 0$

$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$




Case $2x_1 - x_2 = 0$

$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$




Case $2x_1 - x_2 = 0$

$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$




Case $2x_1 - x_2 = 0$

$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$




Case $2x_1 - x_2 = 0$

$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$




Case $2x_1 - x_2 = 0$

$$\alpha x_1 + \beta x_2 = \gamma \xrightarrow{b} 2\beta x_1 + \alpha x_2 = \gamma - b\beta$$




Case $2x_1 - x_2 = 0$




$$x_1 \quad = \quad (\text{binary}) \quad = 0 \quad (\text{decimal})$$

$$x_2 \quad = \quad (\text{binary}) \quad = 0 \quad (\text{decimal})$$

Case $2x_1 - x_2 = 0$




Case $2x_1 - x_2 = 0$




$$\begin{array}{ll} x_1 &= 1 \\ x_2 &= 0 \end{array} \quad \begin{array}{l} (\text{binary}) \\ (\text{binary}) \end{array} \quad \begin{array}{ll} = 1 \\ = 0 \end{array} \quad \begin{array}{l} (\text{decimal}) \\ (\text{decimal}) \end{array}$$

Case $2x_1 - x_2 = 0$




$$\begin{array}{lll} x_1 & = 1 & \text{(binary)} \\ x_2 & = 0 1 & \leftarrow \text{(binary)} \end{array} \quad \begin{array}{lll} & = 1 & \text{(decimal)} \\ & = 2 & \text{(decimal)} \end{array}$$

Case $2x_1 - x_2 = 0$



Case $2x_1 - x_2 = 0$



$$x_1 = 1 \ 0 \quad (\text{binary}) \quad = 1 \quad (\text{decimal})$$

$$x_2 = 0 \ 1 \quad (\text{binary}) \quad = 2 \quad (\text{decimal})$$