

Automates et grammaires algébriques :  
universalité de l'asymptotique  
et  
biodiversité des lois limites

Hanane Tafat Bouzid

University of Paris North,

03 février 2012  
JCB 2012, Bordeaux

# Plan

## 1 Introduction

## 2 Regular grammar

- Asymptotics
- Limits laws of motif
  - In strongly connected graph
  - In non-strongly connected graph
- Applications

## 3 Algebraic grammar

- asymptotics
  - In strongly connected graph
  - In non-strongly connected graph

## 4 Conclusion

A **strongly connected graph** is a directed graph that has a path from each vertex to every other vertex.


Language

$$\mathcal{L} = \{(ab^*a \cup ba^*b)^*\}$$

Grammar

$$\begin{aligned} A &\rightarrow aB \mid bC \mid \epsilon \\ B &\rightarrow aA \mid bB \\ C &\rightarrow aC \mid bA \end{aligned}$$


Dependency Graph



strongly connected graph.

$$\mathcal{L} = \{a^n b^m a^m b^n, n \geq 0, m \geq 0\}$$

$$\begin{aligned} A &\rightarrow aAb \mid B \\ B &\rightarrow bBa \mid \epsilon \end{aligned}$$



non-strongly connected graph.

# Plan

## 1 Introduction

## 2 Regular grammar

- Asymptotics
- Limits laws of motif
  - In strongly connected graph
  - In non-strongly connected graph
- Applications

## 3 Algebraic grammar

- asymptotics
  - In strongly connected graph
  - In non-strongly connected graph

## 4 Conclusion

# Rational generating function

- $\mathcal{L} = \{(a \cdot b)^n, n \geq 0\}$


Right regular grammar

$$\begin{aligned} L &\rightarrow a \cdot A \mid \epsilon \\ A &\rightarrow b \cdot L \end{aligned}$$

Left regular grammar

$$\begin{aligned} L &\rightarrow A \cdot b \mid \epsilon \\ A &\rightarrow L \cdot a \end{aligned}$$

Automate



- A language is regular iff it is recognized by a finite deterministic automaton.

## Theorem [Kleene, 1956]

The generating function of a regular language belongs to the set  $\mathbb{Q}(z)$  of rational functions:

$$\mathbb{N}^{rat}[[z]] \subsetneq \mathbb{Q}(z).$$

- $\mathbb{N}^{rat}[[z]]$  is also characterized as the smallest set of generating functions closed by addition, product, quasi-inverse.

## Closure properties

$$\mathcal{A} = \{a^n, n \geq 0\}, \mathcal{B} = \{b^n, n \geq 0\}$$

- Hadamard Product  $\mathcal{A} \odot \mathcal{B} := \{u.v, |u| = |v|, u \in \mathcal{A} \text{ and } v \in \mathcal{B}\}$

$$\mathcal{A} \odot \mathcal{B} = \{a^n \cdot b^n, n \geq 0\} \quad L \rightarrow a \cdot L \cdot b \mid \epsilon$$

- Shuffle  $\mathcal{A} \sqcup \mathcal{B} := \bigcup_{w_1 \in \mathcal{A}, w_2 \in \mathcal{B}} (w_1 \sqcup w_2)$

$$\mathcal{A} \sqcup \mathcal{B} = \{(a \cup b)^n, n \geq 0\} \quad L \rightarrow a \cdot L \mid b \cdot L \mid \epsilon$$

- Derivation  $\mathcal{A}' := \frac{\partial}{\partial \Sigma} \mathcal{A}$

$$\mathcal{A} = \{a^n, n \geq 0\} \quad \mathcal{A}' = \{a^n \cdot a^m, n \geq 0, m \geq 0\}$$

$$A \rightarrow a \cdot A \mid \epsilon \quad \begin{aligned} A' &\rightarrow a \cdot A' \mid A \\ A &\rightarrow a \cdot A \mid \epsilon \end{aligned}$$

- Complement, intersection, square root...

## Asymptotics of rational series (folklore)

- Let  $F(z)$  be a rational function analytic at 0 then

$$[z^n]F(z) = [z^n] \frac{P(z)}{Q(z)} = [z^n] \sum_{\rho \in \text{pole}(F)} \sum_{r=1}^{\text{order}(\rho)} \frac{C_{\rho,r}}{(z - \rho)^r}$$

$$= \sum_{\rho \in \text{pole}(F)} \sum_{r=1}^{\text{order}(\rho)} C_{\rho,r} (-1)^r \binom{n+r-1}{r-1} \rho^{-n-r}$$

- Let  $\rho$  be **the unique dominant singularity** and let  $r$  be its order, then

$$[z^n]F(z) \sim C_{\rho,r} (-1)^r \binom{n+r-1}{r-1} \rho^{-n-r}$$

- However, some difficult problems:  $\exists f_n = 0$ ? (Pisot's problem), fluctuations of the positivity of the  $(f_n)$ 's (conjugate roots of same modulus, e.g.  $F(z) = \frac{1}{1-6z+25z^2}$ )

- Let  $F(z, u)$  be a bivariate rational generating function

$$F(z, u) = \sum_{n \geq 0} f_n(u) z^n = \sum_{n \geq 0} \sum_{k=0}^n f_{n,k} u^k z^n,$$

where the number of words of length  $n$  with  $k$  “ $b$ ” is  $f_{n,k}$ .

- $X_n$ : a random variable counting the number of letters “ $b$ ” in words of length  $n$

$$\Pr(X_n = k) = \frac{f_{n,k}}{f_n(1)}.$$

What is the limit distribution of  $X_n$ ?

## Limit law in strongly connected graph

- [Edward E.Bender, 1973]: Central and Local Limit Theorems Applied to Asymptotic Enumeration.
- [Guibas–Odlyzko, 1981]: String overlaps, pattern matching, and nontransitive games.
- [Régnier–Szpankowski, 1998]: On pattern frequency occurrences in a Markovian sequence.
- [Bertoni–Choffrut–Goldwurm–Lonati, 2006]: Asymptotics in a regular language of the number of words with a fixed number of occurrences.

### Theorem [Nicodème–Salvy–Flajolet, 2002]

If the language  $\mathcal{L}$  is rational with strongly connected dependency graph, then  $X_n$  is asymptotically Gaussian with mean  $\mu_n$  and variance  $\sigma_n^2$ :


$$\Pr \left[ \frac{X_n - \mu_n}{\sigma_n} \leq x \right] \rightarrow \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x \exp\left(-\frac{t^2}{2}\right) dt.$$

Sketch of proof: Perron–Frobenius and Hwang quasi-power theorem.


# Non-Strongly Connected Graph

Theorem [BaBoPoTa, 2011]

For any continuous curve  $C$ , there exists a regular language in which  $X_n$  (number of "b" in words of length  $n$ ) has a limit distribution which  $\epsilon$ -near from  $C$ .



$$(x, y) \rightarrow \left( \frac{x}{n}, \frac{y}{f_n(1)} \right)$$



## Theorem [BaBoPoTa, 2011]

For any polynomial  $P(n, k)$  in  $\mathbb{R}^{\geq 0}[n, k]$ , there exists a power series  $F(z, u) \in \mathbb{N}^{\text{rat}}[[z]]$  such that

$$\Pr(X_n = k) = \frac{[z^n][u^k]F(z, u)}{[z^n]F(z, 1)} = \frac{P(n, k)}{\sum_{i=0}^n P(n, i)}.$$

### Sketch of proof

- $P(n, k)$  is a linear combination of  $n^\alpha k^\beta$
- $n^\alpha k^\beta = [z^n][u^k] \sum_{n \geq 0} n^\alpha z^n \sum_{k \geq 0} k^\beta u^k$ .

□

## Lemma

The power series  $\sum_{n \geq 0} n^i z^n$  belongs to  $\mathbb{N}^{rat}[[z]]$  and more precisely


$$\sum_{n \geq 0} n^i z^n = \sum_{j=0}^i j! \begin{Bmatrix} i \\ j \end{Bmatrix} \frac{z^j}{(1-z)^{j+1}} = \frac{\sum_{j=0}^i \begin{Bmatrix} i \\ j \end{Bmatrix} z^{j+1}}{(1-z)^{i+1}}.$$

Proof (by induction):


- apply  $z\partial_z$  to the equality (equivalent to pointing  $\Theta$  in the symbolic method),
- use the recurrence defining:

- ▶ the Stirling numbers of the second kind  $\begin{Bmatrix} i+1 \\ j \end{Bmatrix} = \begin{Bmatrix} i \\ j \end{Bmatrix} + j \begin{Bmatrix} i \\ j \end{Bmatrix}$
- ▶ the Eulerian numbers  $\begin{Bmatrix} i+1 \\ j \end{Bmatrix} = (j+1) \begin{Bmatrix} i \\ j \end{Bmatrix} + (i+1-j) \begin{Bmatrix} i \\ j-1 \end{Bmatrix}$ . □


# Examples




$$\frac{1}{(1-z)} \frac{1}{(1-uz)}$$




$$\frac{1}{(1-z)} \frac{1}{(1-u^3 z)}$$



$$\frac{(u+u^2+u^3)}{(1-z)(1-u^3 z)}$$



$$\frac{1}{(1-z)^2} \frac{1}{(1-uz)}$$



$$\frac{1}{(1-z)} \frac{1}{(1-uz)^2}$$



- $\mathcal{C} = \mathcal{A} \cup \mathcal{B}$
- $\Pr(X_n^{\mathcal{C}} = k) = \frac{[u^k][z^n]C(z,u)}{[z^n]C(z,1)} = \frac{[u^k][z^n](A(z,u)+B(z,u))}{[z^n](A(z,1)+B(z,1))}$

- ▶ If  $(\rho_{\mathcal{A}} < \rho_{\mathcal{B}})$  or  $(\rho_{\mathcal{A}} = \rho_{\mathcal{B}}$  and  $r_{\mathcal{A}} < r_{\mathcal{B}}$ ) then

$$\Pr(X_n^{\mathcal{C}} = k) \sim \Pr(X_n^{\mathcal{A}} = k)$$

- ▶ If  $\rho_{\mathcal{A}} = \rho_{\mathcal{B}}$  and  $r_{\mathcal{A}} = r_{\mathcal{B}}$  then







$$\Pr(X_n^{\mathcal{C}} = k) = c_1 \Pr(X_n^{\mathcal{A}} = k) + c_2 \Pr(X_n^{\mathcal{B}} = k)$$



$$\frac{1}{(1-z)^2(1-u^2z)} + \frac{1}{(1-z)(1-u^2z)^2}$$

$$\frac{1}{(1-z)^2(1-u^2z)} + \frac{u}{(1-z)(1-u^2z)^2} + \frac{1}{(1-z)^2(1-u^2z)}$$

# Examples



# Boltzmann Sampling

- Boltzmann samplers allows the generation of uniform random decomposable combinatorial objects.
- [P. Duchon–P. Flajolet–G. Louchard–G. Schaeffer, 2004]: Boltzmann Samplers for the random generation of combinatorial structures.
- [O. Bodini–Y. Ponty, 2010]: Multidimensional Boltzmann Sampling of languages.
- The Boltzmann sampler returns an object  $\gamma$  of size  $n$  and parameter  $k$  with

$$\Pr_{x,u}(\gamma) = \frac{x^n u^k}{F(x,u)}.$$

# Boltzmann Sampling

## Theorem [BaBoPoTa, 2011] intrinsic efficiency for the size

For any fixed tolerance  $\varepsilon > 0$ , the Boltzmann sampler size generates, in average, in  $O(n)$  independently in  $u_0$ , an object of size  $N$  in the interval  $[(1 - \varepsilon), (1 + \varepsilon)n]$ .

## Theorem [BaBoPoTa, 2011] intrinsic impossibility for the frequency

For any fixed tolerance  $0 \leq \delta_u < 1$ , the Boltzmann sampler size-frequency could be unable to generate an object with a frequency of  $u$  in an interval  $[(\mu_u - \delta_u), (\mu_u + \delta_u)]$  where  $\mu_u$  is the mean of  $u$ .

# Plan

## 1 Introduction

## 2 Regular grammar

- Asymptotics
- Limits laws of motif
  - In strongly connected graph
  - In non-strongly connected graph
- Applications

## 3 Algebraic grammar

- asymptotics
  - In strongly connected graph
  - In non-strongly connected graph

## 4 Conclusion

# Algebraic Branches

Combinatorial object	Algebraic grammar	Functional Equation	Branches
Binary trees	$T \rightarrow a \cdot T \cdot T   a$	$z = T(z) + zT^2(z)$	$\left\{ \begin{array}{l} \frac{1+\sqrt{1-4z^2}}{2z} \\ \frac{1-\sqrt{1-4z^2}}{2z} \end{array} \right.$
Dyck words	$D \rightarrow a \cdot D \cdot b \cdot D   \epsilon$	$1 - D(z) + z^2 D^2(z)$	$\left\{ \begin{array}{l} \frac{1+\sqrt{1-4z^2}}{2z^2} \\ \frac{1-\sqrt{1-4z^2}}{2z^2} \end{array} \right.$

Algebraic grammar with  $m$  non terminals:

- ① Polynomial system of equations  $Y_j = P_i(z, Y_1, \dots, Y_m)$ ,
- ②  $P_j$  have non-negative coefficients,
- ③  $\vec{Y}^{(0)} = (0, \dots, 0)$ ,  $\vec{Y}^{(h+1)} = P_i(\vec{Y}^{(h)})$ ,  $\vec{Y} = \lim_{h \rightarrow \infty} \vec{Y}^{(h)}$

A unique branch analytic with non-negative Taylor coefficients around 0.

# Asymptotics of algebraic grammars

## Theorem of Newton-Puiseux

Let  $f(z)$  the branch of an algebraic equation  $P(z, f(z)) = 0$ , near its singularities  $\rho$ ,  $f(z)$  admits a convergent fractional series expansion of the form:  $f(z) = \sum_{k \geq k_0} c_k \cdot (z - \rho)^{\frac{k}{K}}$  with  $k_0 \in \mathbb{Z}$  and  $K \geq 1$ .

asymptotics:

$$f_n \sim c_{k_0} \frac{1}{\Gamma(-\alpha)} \rho^{-n} n^{-\alpha-1} \text{ with } \alpha = \frac{k_0}{K}$$

What are the possible values of  $\alpha$ ?

# Strongly connected graph

## Theorem of [Drmota–Lalley–Woods, 97–93–97]

Let  $\vec{Y} = P_i(\vec{Y})$  a system satisfying:

- ① All polynomials  $P_i$  have positive coefficients .
- ② The system admits a fixed point  $\vec{Y}$ .
- ③ The dependency graph of the system is strongly connected.

Then:

- ① all solutions behave like  $Y_j(z) = C_{0j} - C_j \cdot \sqrt{1 - \frac{z}{\rho}} + O(1 - \frac{z}{\rho})$  in neighborhood  $z = \rho$ ,
- ②  $[z^n] Y_j(z) \sim C_j \cdot \frac{1}{2\sqrt{\pi}} \cdot \rho^{-n} \cdot n^{-\frac{3}{2}}$  with  $C_{0j}, C_j, \rho$  algebraic numbers.

## Example

$$\begin{cases} \mathcal{A} \rightarrow a \cdot \mathcal{A}|b \cdot \mathcal{B} \cdot \mathcal{B}|c \\ \mathcal{B} \rightarrow b \cdot \mathcal{A}|a \cdot \mathcal{B} \cdot \mathcal{B}|c \end{cases}$$


- $\begin{cases} A(z) = zA(z) + zB(z)^2 + z \\ B(z) = zA(z) + zB(z)^2 + z \end{cases}$
- Algebraic equation:  $zA(z)^2 + (z - 1)A(z) + z = 0$
- Branches of  $A(z)$ :  
$$-\frac{z - 1 + \sqrt{-3z^2 - 2z + 1}}{2z}, \quad \frac{-z + 1 + \sqrt{-3z^2 - 2z + 1}}{2z}$$
- The singularities:  $-1, 0, 1/3$
- Puiseux expansion:  $A(z) = \frac{5}{2} - \sqrt{3(1 - 3z)} + O(1 - 3z)$

## non-strongly connected graph

general trees

$$\mathcal{T} = z \text{Seq}(\mathcal{T})$$

$$\left\{ \begin{array}{l} \mathcal{T} \rightarrow a \cdot \mathcal{T} \cdot \mathcal{A}|a \\ \mathcal{A} \rightarrow \mathcal{T} \cdot \mathcal{A}|\epsilon \end{array} \right.$$




strongly connected.

super bicolore trees:

$$\mathcal{B} = \mathcal{T}[z.\mathcal{T}]$$

$$\mathcal{B} = 2z\mathcal{T}\text{Seq}(\mathcal{B})$$

$$\left\{ \begin{array}{l} \mathcal{B} \rightarrow a \cdot \mathcal{T} \cdot \mathcal{U}|b \cdot \mathcal{T} \cdot \mathcal{U} \\ \mathcal{U} \rightarrow \mathcal{B} \cdot \mathcal{U}|\epsilon \\ \mathcal{T} \rightarrow a \cdot \mathcal{T} \cdot \mathcal{A}|a \\ \mathcal{A} \rightarrow \mathcal{T} \cdot \mathcal{A}|\epsilon \end{array} \right.$$



non-strongly connected.

Branch:  $B(z) = \frac{1}{2} - \frac{1}{2}\sqrt{1 - 4z + 4z\sqrt{1 - 4z}}$

Puiseux expansion:  $B(z) = \frac{1}{2} - \frac{1}{2}(1 - 4z)^{1/4} + O((1 - 4z)^{3/4})$

- The behavior asymptotics of bicolored trees  $\mathcal{T}$  is  $1/2$
- Super-bicolored trees:  $\mathcal{B} = \mathcal{T}[(z + z).\mathcal{T}]$ 
  - ▶ schema of composition critique,
  - ▶  $\frac{1}{2^2}$  is the behavior asymptotics.
- $k$ -Super-bicolored trees:  $\mathcal{B}_k = \mathcal{T}[(z + z).\mathcal{B}_{k-1}]$ 
  - ▶ schema of  $k$ - composition critique,
  - ▶  $\frac{1}{2^{k+1}}$  is the behavior asymptotics.

Using Maple (Algolib:Combstruct, gdev) we obtain: for  $k$  compositions de  $\mathcal{B}$   $\frac{1}{2^{k+1}}$  behavior.

## Theorem

Suppose that  $\vec{Y} = P(z, \vec{Y})$  is a positive polynomial system such that there is a unique solution  $(Y_1(z), \dots, Y_m(z))$  that is analytic at  $z = 0$ .


What is more, if  $\frac{\partial^2 P_j}{\partial^2 Y_j} \neq 0$  then for every  $j$  there exists an integer  $k_j \geq 1$  such that locally

$$Y_j(z) = \sum_{n \geq 0} c_n \cdot (z - \rho)^{n/2^{k_j}}$$

super bicolore trees:

$$\mathcal{B} = 2z\mathcal{T} \text{Seq}(\mathcal{B})$$

$$\left\{ \begin{array}{l} \mathcal{B} \rightarrow a \cdot \mathcal{T} \cdot \mathcal{U} | b \cdot \mathcal{T} \cdot \mathcal{U} \\ \mathcal{U} \rightarrow \mathcal{B} \cdot \mathcal{U} | \epsilon \\ \mathcal{T} \rightarrow a \cdot \mathcal{T} \cdot \mathcal{A} | a \\ \mathcal{A} \rightarrow \mathcal{T} \cdot \mathcal{A} | \epsilon \end{array} \right.$$



non-strongly connected.


## the derivative

- The generating function of Super-bicolored trees:

$$B(z) = \frac{1}{2} - \frac{1}{2}(1-4z)^{1/4} + O((1-4z)^{3/4})$$

$$\frac{\partial}{\partial z} B(z) = \frac{1}{2}(1-4z)^{-3/4} + O((1-4z)^{-1/4})$$

$$\left\{ \begin{array}{l} \mathcal{B}' \rightarrow \mathcal{T} \cdot \mathcal{U}|a \cdot \mathcal{T}' \cdot \mathcal{U}|a \cdot \mathcal{T} \cdot \mathcal{U}'|b \cdot \mathcal{T}' \cdot \mathcal{U}|b \cdot \mathcal{T} \cdot \mathcal{U}' \\ \mathcal{B} \rightarrow a \cdot \mathcal{T} \cdot \mathcal{U}|b \cdot \mathcal{T} \cdot \mathcal{U} \\ \mathcal{T}' \rightarrow \mathcal{T}\mathcal{A}|a \cdot \mathcal{T}' \cdot \mathcal{A}|a \cdot \mathcal{T} \cdot \mathcal{A}'|\epsilon \\ \mathcal{T} \rightarrow a \cdot \mathcal{T} \cdot \mathcal{A}|a \\ \mathcal{U}' \rightarrow \mathcal{B}' \cdot \mathcal{U}|\mathcal{B} \cdot \mathcal{U}'|\epsilon \\ \mathcal{U} \rightarrow \mathcal{B} \cdot \mathcal{U}|\epsilon \\ \mathcal{A}' \rightarrow \mathcal{T}' \cdot \mathcal{A}|\mathcal{T} \cdot \mathcal{A}'|\epsilon \\ \mathcal{A} \rightarrow \mathcal{T} \cdot \mathcal{A}|\epsilon \end{array} \right.$$



non-strongly connected.

The dependency graph still non-strongly connected

## Theorem

Suppose that  $\vec{Y} = P(z, \vec{Y})$  is a positive polynomial system such that there is a unique solution  $(Y_1(z), \dots, Y_m(z))$  that is analytic at  $z = 0$ .

What is more, if for all  $(i, j)$   $\frac{\partial P_j}{\partial Y_i} \neq 0$  and  $\frac{\partial^2 P_j}{\partial Y_i \partial Y_h} \neq 0$  then for every  $j$  there exists an integer  $k_j \geq 1$  such that locally

$$Y_j(z) = \sum_{n \geq 0} c_n \cdot (z - \rho)^{n/2^{k_j}}$$

or

$$Y_j(z) = \sum_{n \geq 0} c_n \cdot (z - \rho)^{n \frac{1}{2^{k_j}} - 1}$$

$$\left\{ \begin{array}{l} \mathcal{B}' \rightarrow \mathcal{T} \cdot \mathcal{U} | a \cdot \mathcal{T}' \cdot \mathcal{U} | a \cdot \mathcal{T} \cdot \mathcal{U}' | b \cdot \mathcal{T}' \cdot \mathcal{U} | b \cdot \mathcal{T} \cdot \mathcal{U}' \\ \mathcal{B} \rightarrow a \cdot \mathcal{T} \cdot \mathcal{U} | b \cdot \mathcal{T} \cdot \mathcal{U} \\ \vdots \end{array} \right.$$

## Theorem

Suppose that  $\vec{Y} = P(z, \vec{Y})$  is a positive polynomial system of entire functions such that there is a unique solution  $(Y_1(z), \dots, Y_m(z))$  that is analytic at  $z = 0$ .


$$Y_j(z) = \sum_{n \geq 0} c_n \cdot (z - \rho)^{n/2^{k_j}}$$

or

$$Y_j(z) = \sum_{n \geq 0} c_n \cdot (z - \rho)^{n \frac{1}{2^{k_j}} - m}$$

## Example

$$\left\{ \begin{array}{l} \mathcal{A} \rightarrow a \cdot \mathcal{A} \cdot \mathcal{A} \cdot \mathcal{A} | b \cdot \mathcal{A} \\ \mathcal{B} \rightarrow a | a \cdot \mathcal{BC} | b \cdot \mathcal{BC} \\ \mathcal{C} \rightarrow a | b \cdot \mathcal{CC} \end{array} \right.$$



$$\left\{ \begin{array}{l} A(z) = \frac{z^4(1-4z^2)^{-3/2}}{1-z} \\ B(z) = z(1-4z^2)^{-1/2} \\ C(z) = \frac{1-(1-4z^2)^{1/2}}{2z} \end{array} \right.$$

$$\left\{ \begin{array}{l} A(z) = \frac{\sqrt{2}}{32}(1-2z)^{-3/2} + \dots \\ B(z) = \frac{\sqrt{2}}{4}(1-2z)^{-1/2} + \dots \\ C(z) = 1 - \sqrt{2}(1-2z)^{1/2} + \dots \end{array} \right.$$

$$[z^n]A(z) \sim \frac{\sqrt{2}}{56\sqrt{\pi}} 2^n n^{1/2}$$

# Plan

## 1 Introduction

## 2 Regular grammar

- Asymptotics
- Limits laws of motif
  - In strongly connected graph
  - In non-strongly connected graph
- Applications

## 3 Algebraic grammar

- asymptotics
  - In strongly connected graph
  - In non-strongly connected graph

## 4 Conclusion

# Conclusion

- We gave distributions which can be reached as a pattern occurrence distribution (+ application: random generation).
- Work in progress: the asymptotic behavior and the possible limit laws in non strongly connected context free-grammars?
- Implementation (in SageMath):
  - ▶ Input: (`language`, `pattern`),
  - ▶ Output: (`limit law`, `multivariate Boltzmann sampler`).