

Central limit theorem for random Young diagrams with respect to Jack measure (joint work with Valentin Féray)

Maciej Dołęga


LIAFA, Université Paris Diderot,
Instytut Matematyczny, Uniwersytet Wrocławski

12 II 2014

Young diagrams

Definition

A **partition** λ is a finite non-increasing sequence of positive integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$. It can be represented by a **Young diagram** λ . The **size** of the Young diagram λ is defined by $|\lambda| := \sum_i \lambda_i$.



Problem

We want to investigate some asymptotic properties of Young diagrams as their size is tending to infinity. How to do it?


Solution

Look on 'large Young diagrams' from a 'large perspective' and treat these discrete objects as continuous ones!

Young diagrams

Definition

A **partition** λ is a finite non-increasing sequence of positive integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$. It can be represented by a **Young diagram** λ . The **size** of the Young diagram λ is defined by $|\lambda| := \sum_i \lambda_i$.



Problem

We want to investigate some asymptotic properties of Young diagrams as their size is tending to infinity. How to do it?


Solution

Look on 'large Young diagrams' from a 'large perspective' and treat these discrete objects as continuous ones!

Young diagrams

Definition

A **partition** λ is a finite non-increasing sequence of positive integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$. It can be represented by a **Young diagram** λ . The **size** of the Young diagram λ is defined by $|\lambda| := \sum_i \lambda_i$.



Problem


We want to investigate some asymptotic properties of Young diagrams as their size is tending to infinity. How to do it?

Solution

Look on 'large Young diagrams' from a 'large perspective' and treat these discrete objects as continuous ones!


Young diagrams as continuous objects

French convention:




Young diagrams as continuous objects

French convention:




Young diagrams as continuous objects

Russian convention:




Young diagrams as continuous objects

Russian convention:



Young diagrams as continuous objects

Russian convention:



Definition

A **profile** of a Young diagram λ is a function $\omega(\lambda) : \mathbb{R} \rightarrow \mathbb{R}_+$ such that its graph is a profile of λ drawn in Russian convention.

Continuous Young diagrams

Definition

A **continuous Young diagram** is a function $\omega : \mathbb{R} \rightarrow \mathbb{R}_+$ such that

- $\omega(x) - |x|$ has compact support;
- $|\omega(x_1) - \omega(x_2)| \leq |x_1 - x_2|$ for any $x_1, x_2 \in \mathbb{R}$.

An **area** of a continuous Young diagram ω is given by:

$$\text{Area}(\omega) := \frac{1}{2} \int_{\mathbb{R}} |\omega(x) - |x|| \, dx.$$

Remark

Let λ - Young diagram with $|\lambda| = n$. Then

$$\text{Area}(\omega(\lambda)) = n.$$

Normalized Young diagrams

Problem

How to look on the 'large Young diagrams' from 'large perspective'?

Solution

Normalize them in a way that their areas are constant.

Definition

Let λ - Young diagram with $|\lambda| = n$. We define **scaled** (continuous) Young diagram

$$\omega(D_{\sqrt{n}^{-1}}(\lambda))(x) := \sqrt{n}^{-1} \omega(\lambda)(\sqrt{nx}).$$

Remark

$$\text{Area} \left(\omega(D_{\sqrt{n}^{-1}}(\lambda)) \right) = 1.$$

Normalized Young diagrams

Problem

How to look on the 'large Young diagrams' from 'large perspective'?

Solution

Normalize them in a way that their areas are constant.

Definition

Let λ - Young diagram with $|\lambda| = n$. We define **scaled** (continuous) Young diagram

$$\omega(D_{\sqrt{n}^{-1}}(\lambda))(x) := \sqrt{n}^{-1} \omega(\lambda)(\sqrt{n}x).$$

Remark

$$\text{Area} \left(\omega(D_{\sqrt{n}^{-1}}(\lambda)) \right) = 1.$$

Asymptotic shape of large Young diagrams

Let $(\lambda_{(n)})_{n \in \mathbb{N}_+}$ - sequence of Young diagrams with $|\lambda_{(n)}| = n$.

Definition

We say that $(\lambda_{(n)})_{n \in \mathbb{N}_+}$ has a **limit shape** ω if

$$\left\| \omega(D_{\sqrt{n}^{-1}}(\lambda_{(n)})) - \omega \right\| \rightarrow 0,$$

as $n \rightarrow \infty$, where $\|f\| = \sup_{x \in \mathbb{R}} |f(x)|$.

Problem

Let us choose $(\lambda_{(n)})_{n \in \mathbb{N}_+}$ randomly according with some 'nice' distribution. Does it have a limit shape with a high probability? Is it unique? Can we say more about it?

Asymptotic shape of large Young diagrams

Let $(\lambda_{(n)})_{n \in \mathbb{N}_+}$ - sequence of Young diagrams with $|\lambda_{(n)}| = n$.

Definition

We say that $(\lambda_{(n)})_{n \in \mathbb{N}_+}$ has a **limit shape** ω if


$$\left\| \omega(D_{\sqrt{n}^{-1}}(\lambda_{(n)})) - \omega \right\| \rightarrow 0,$$

as $n \rightarrow \infty$, where $\|f\| = \sup_{x \in \mathbb{R}} |f(x)|$.

Problem


Let us choose $(\lambda_{(n)})_{n \in \mathbb{N}_+}$ randomly according with some 'nice' distribution. Does it have a limit shape with a high probability? Is it unique? Can we say more about it?

'Nice' distribution = Plancherel distribution




$a(\bullet)$ = number of boxes to the right of the given box

'Nice' distribution = Plancherel distribution




$\ell(\bullet)$ = number of boxes above the given box

'Nice' distribution = Plancherel distribution



$$\text{hook}^{(1)}(\bullet) := a(\bullet) + \ell(\bullet) + 1.$$

'Nice' distribution = Plancherel distribution




$$\mathbb{P}_n^{(1)}(\lambda) = \frac{\dim(\lambda)^2}{n!},$$

where (hook formula:)

$$\dim(\lambda) = \frac{n!}{\prod_{\square \in \lambda} \text{hook}(\square)}.$$

'Nice' distribution = Plancherel distribution



$$\mathbb{P}_n^{(1)}(\lambda) = \frac{\dim(\lambda)^2}{n!},$$

where (**hook formula:**)

$$\dim(\lambda) = \frac{n!}{\prod_{\square \in \lambda} \text{hook}(\square)}.$$

$$\mathbb{P}_n^{(1)}(\lambda) = \frac{n!}{\prod_{\square \in \lambda} (\text{hook}(\square))^2}.$$

Plancherel measure $\mathbb{P}_n^{(1)}$ is a **probability measure** on the set \mathbb{Y}_n of Young diagrams of size n .

Vershik-Kerov, Logan-Shepp limit shape




Figure : Scaled random Young diagram of size 100 distributed according with Plancherel measure

Vershik-Kerov, Logan-Shepp limit shape




Figure : Scaled random Young diagram of size 1000 distributed according with Plancherel measure

Vershik-Kerov, Logan-Shepp limit shape




Figure : Scaled random Young diagram of size 5000 distributed according with Plancherel measure

Vershik-Kerov, Logan-Shepp limit shape





Figure :
$$\Omega(x) = \begin{cases} |x| & \text{if } |x| \geq 2; \\ \frac{2}{\pi} \left(x \cdot \arcsin \frac{x}{2} + \sqrt{4 - x^2} \right) & \text{otherwise.} \end{cases}$$

First order asymptotic = 'law of large numbers'



$$\Omega(x) = \begin{cases} |x| & \text{if } |x| \geq 2; \\ \frac{2}{\pi} \left(x \cdot \arcsin \frac{x}{2} + \sqrt{4 - x^2} \right) & \text{otherwise.} \end{cases}$$

Theorem (Vershik-Kerov, Logan-Shepp '77)

Let $\lambda_{(n)}$ be a random Young diagram of size n distributed with *Plancherel measure* $\mathbb{P}_n^{(1)}$. Then, in probability, as $n \rightarrow \infty$

$$\left\| \omega(D_{1/\sqrt{n}}(\lambda_{(n)})) - \Omega \right\| \rightarrow 0.$$

Second order asymptotic

Problem

Can we describe the second order asymptotic? What does it really mean?

Solution

We should look on the fluctuations around the limit shape.

Let $\lambda_{(n)}$ - random Young diagram distributed according with $\mathbb{P}_n^{(1)}$.

- We know that $\|\omega(D_{1/\sqrt{n}}(\lambda_{(n)})) - \Omega\| \rightarrow 0$ in probability.
- We would like to investigate behaviour of random variables:

$$m_k(\lambda_{(n)}) := \int_{\mathbb{R}} x^k \Delta(\lambda_{(n)})(x) dx,$$

where

$$\Delta(\lambda)(x) := \sqrt{n} \frac{\omega(D_{1/\sqrt{n}}(\lambda))(x) - \Omega(x)}{2}.$$

Second order asymptotic

Problem

Can we describe the second order asymptotic? What does it really mean?

Solution

We should look on the fluctuations around the limit shape.

Let $\lambda_{(n)}$ - random Young diagram distributed according with $\mathbb{P}_n^{(1)}$.

- We know that $\|\omega(D_{1/\sqrt{n}}(\lambda_{(n)})) - \Omega\| \rightarrow 0$ in probability.
- We would like to investigate behaviour of random variables:

$$m_k(\lambda_{(n)}) := \int_{\mathbb{R}} x^k \Delta(\lambda_{(n)})(x) dx,$$

where

$$\Delta(\lambda)(x) := \sqrt{n} \frac{\omega(D_{1/\sqrt{n}}(\lambda))(x) - \Omega(x)}{2}.$$

Second order asymptotic = 'central limit theorem'

- $u_k(x) = U_k(x/2) = \sum_{0 \leq j \leq \lfloor k/2 \rfloor} (-1)^j \binom{k-j}{j} x^{k-2j};$
- $u_k(2 \cos(\theta)) = \frac{\sin((k+1)\theta)}{\sin(\theta)};$
- $u_k(\lambda) = \int_{\mathbb{R}} u_k(x) \Delta(\lambda)(x) dx.$

Theorem (Kerov, 1993)

Choose a sequence $(\Xi_k)_{k=2,3,\dots}$ of independent standard Gaussian random variables and let $\lambda_{(n)}$ be a random Young diagram of size n distributed with Plancherel measure. As $n \rightarrow \infty$, we have:

$$(u_k(\lambda_{(n)}))_{k=1,2,\dots} \xrightarrow{d} \left(\frac{\Xi_{k+1}}{\sqrt{k+1}} \right)_{k=1,2,\dots}.$$

Second order asymptotic = 'central limit theorem'

- $u_k(x) = U_k(x/2) = \sum_{0 \leq j \leq \lfloor k/2 \rfloor} (-1)^j \binom{k-j}{j} x^{k-2j}$;
- $u_k(2 \cos(\theta)) = \frac{\sin((k+1)\theta)}{\sin(\theta)}$;
- $u_k(\lambda) = \int_{\mathbb{R}} u_k(x) \Delta(\lambda)(x) dx$.

Theorem (Kerov, 1993)

Choose a sequence $(\Xi_k)_{k=2,3,\dots}$ of independent standard Gaussian random variables and let $\lambda_{(n)}$ be a random Young diagram of size n distributed with Plancherel measure. As $n \rightarrow \infty$, we have:

$$(u_k(\lambda_{(n)}))_{k=1,2,\dots} \xrightarrow{d} \left(\frac{\Xi_{k+1}}{\sqrt{k+1}} \right)_{k=1,2,\dots}.$$

Jack measure

Let us recall that

$$\text{hook}(\square) = a(\square) + \ell(\square) + 1.$$

Let $\alpha \in \mathbb{R}_+$. α -deformations of hook length:

- $\text{hook}^{(\alpha)}(\square) := \sqrt{\alpha}(a(\square) + 1) + \sqrt{\alpha^{-1}}\ell(\square),$
- $(\text{hook}^{(\alpha)})'(\square) := \sqrt{\alpha}a(\square) + \sqrt{\alpha^{-1}}(\ell(\square) + 1).$

Definition

is a probability measure \mathbb{P}_n on the set \mathbb{Y}_n defined by

$$\mathbb{P}_n(\lambda) := \frac{n!}{\prod_{\square \in \lambda} (\text{hook}(\square) (\text{hook}')(\square))}$$

Jack measure

Let us recall that

$$\text{hook}(\square) = a(\square) + \ell(\square) + 1.$$

Let $\alpha \in \mathbb{R}_+$. **α -deformations of hook length:**

- $\text{hook}^{(\alpha)}(\square) := \sqrt{\alpha}(a(\square) + 1) + \sqrt{\alpha}^{-1}\ell(\square),$
- $(\text{hook}^{(\alpha)})'(\square) := \sqrt{\alpha}a(\square) + \sqrt{\alpha}^{-1}(\ell(\square) + 1).$

Definition

is a probability measure \mathbb{P}_n on the set \mathbb{Y}_n defined by

$$\mathbb{P}_n(\lambda) := \frac{n!}{\prod_{\square \in \lambda} (\text{hook}(\square) (\text{hook}')(\square))}$$

Jack measure

Let us recall that

$$\text{hook}(\square) = a(\square) + \ell(\square) + 1.$$

Let $\alpha \in \mathbb{R}_+$. **α -deformations of hook length:**

- $\text{hook}^{(\alpha)}(\square) := \sqrt{\alpha}(a(\square) + 1) + \sqrt{\alpha}^{-1}\ell(\square),$
- $\left(\text{hook}^{(\alpha)}\right)'(\square) := \sqrt{\alpha}a(\square) + \sqrt{\alpha}^{-1}(\ell(\square) + 1).$

Definition

Jack measure is a probability measure $\mathbb{P}_n^{(\alpha)}$ on the set \mathbb{Y}_n defined by

$$\mathbb{P}_n^{(\alpha)}(\lambda) := \frac{n!}{\prod_{\square \in \lambda} \left(\text{hook}^{(\alpha)}(\square) \left(\text{hook}^{(\alpha)}\right)'(\square)\right)},$$

where $\alpha \in \mathbb{R}_+$.

Jack measure

Let us recall that

$$\text{hook}(\square) = a(\square) + \ell(\square) + 1.$$

Let $\alpha \in \mathbb{R}_+$. α -deformations of hook length:

- $\text{hook}^{(\alpha)}(\square) := \sqrt{\alpha}(a(\square) + 1) + \sqrt{\alpha}^{-1}\ell(\square),$
- $\left(\text{hook}^{(\alpha)}\right)'(\square) := \sqrt{\alpha}a(\square) + \sqrt{\alpha}^{-1}(\ell(\square) + 1).$

Definition

Plancherel measure is a probability measure $\mathbb{P}_n^{(1)}$ on the set \mathbb{Y}_n defined by

$$\mathbb{P}_n^{(1)}(\lambda) := \frac{n!}{\prod_{\square \in \lambda} (\text{hook}^{(1)}(\square) \left(\text{hook}^{(1)}\right)'(\square))} = \frac{n!}{\prod_{\square \in \lambda} (\text{hook}(\square))^2}$$


- for $\alpha = 1$ Jack measure \equiv Plancherel measure.

α -anisotropic Young diagrams

Let λ be a Young diagram.

Definition

α -anisotropic Young diagram $A_\alpha(\lambda)$ (for $\alpha \in \mathbb{R}_+$) - continuous Young diagram obtained from λ (considered in French convention) by a horizontal stretching of ratio $\sqrt{\alpha}$ and a vertical stretching of ratio $\sqrt{\alpha}^{-1}$.




$$\lambda \mapsto A_4(\lambda)$$


α -anisotropic Young diagrams


Let λ be a Young diagram.

Definition

α -anisotropic Young diagram $A_\alpha(\lambda)$ (for $\alpha \in \mathbb{R}_+$) - continuous Young diagram obtained from λ (considered in French convention) by a horizontal stretching of ratio $\sqrt{\alpha}$ and a vertical stretching of ratio $\sqrt{\alpha}^{-1}$.



$$\lambda \mapsto A_4(\lambda)$$






α -anisotropic Young diagrams

Let λ be a Young diagram.

Definition

α -anisotropic Young diagram $A_\alpha(\lambda)$ (for $\alpha \in \mathbb{R}_+$) - continuous Young diagram obtained from λ (considered in French convention) by a horizontal stretching of ratio $\sqrt{\alpha}$ and a vertical stretching of ratio $\sqrt{\alpha}^{-1}$.






α -anisotropic Young diagrams

Let λ be a Young diagram.

Definition

α -anisotropic Young diagram $A_\alpha(\lambda)$ (for $\alpha \in \mathbb{R}_+$) - continuous Young diagram obtained from λ (considered in French convention) by a horizontal stretching of ratio $\sqrt{\alpha}$ and a vertical stretching of ratio $\sqrt{\alpha}^{-1}$.




α -anisotropic Young diagrams


Let λ be a Young diagram.

Definition

α -anisotropic Young diagram $A_\alpha(\lambda)$ (for $\alpha \in \mathbb{R}_+$) - continuous Young diagram obtained from λ (considered in French convention) by a horizontal stretching of ratio $\sqrt{\alpha}$ and a vertical stretching of ratio $\sqrt{\alpha}^{-1}$.



$$\lambda \mapsto A_4(\lambda)$$






α -anisotropic Young diagrams

Let λ be a Young diagram.

Definition

α -anisotropic Young diagram $A_\alpha(\lambda)$ (for $\alpha \in \mathbb{R}_+$) - continuous Young diagram obtained from λ (considered in French convention) by a horizontal stretching of ratio $\sqrt{\alpha}$ and a vertical stretching of ratio $\sqrt{\alpha}^{-1}$.



First order asymptotic = 'law of large numbers'

Theorem (D., Féray)

Let $\lambda_{(n)}$ be a random Young diagram of size n distributed with **Jack measure** $\mathbb{P}_n^{(\alpha)}$. Then, in probability, as $n \rightarrow \infty$

$$\|\omega(D_{1/\sqrt{n}}(A_\alpha(\lambda_{(n)}))) - \Omega\| \rightarrow 0.$$

Remark

Plugging $\alpha = 1$ we recover Vershik-Kerov, Logan-Shepp limit shape for Plancherel measure.

Second order asymptotic = 'central limit theorem'

Theorem (D. Féray)

Choose a sequence $(\Xi_k)_{k=2,3,\dots}$ of independent standard Gaussian random variables and let $\lambda_{(n)}$ be a random Young diagram of size n distributed with Jack measure. As $n \rightarrow \infty$, we have:

$$\left(u_k^{(\alpha)}(\lambda_{(n)}) \right)_{k=1,2,\dots} \xrightarrow{d} \left(\frac{\Xi_{k+1}}{\sqrt{k+1}} - \frac{\gamma}{k+1} [k \text{ is odd}] \right)_{k=1,2,\dots},$$

where $u_k^{(\alpha)}(\lambda) = \int_{\mathbb{R}} u_k(x) \Delta(A_\alpha(\lambda))(x) dx$, $\gamma := \sqrt{\alpha} - \sqrt{\alpha}^{-1}$, and we use the usual notation $[condition]$ for the indicator function of the corresponding condition.

Remark

Plugging $\alpha = 1$ we recover central limit theorem of Kerov for Plancherel measure.

Jack polynomials and Jack characters

Jack polynomials $J_\lambda^{(\alpha)}$:

- symmetric functions introduced by Jack;
- generalization of Schur symmetric function (for $\alpha = 1$);
- special case of Macdonald polynomials

Expand Jack polynomial in power-sum symmetric basis:

$$J_\lambda^{(\alpha)} = \sum_{\substack{\rho: \\ |\rho|=|\lambda|}} \theta_\rho^{(\alpha)}(\lambda) p_\rho.$$

We call quantities $\theta_\rho^{(\alpha)}(\lambda)$ **Jack characters** (for $\alpha = 1$ they coincide with the irreducible characters of the symmetric groups up to some normalization constant).

Jack polynomials and Jack characters

Jack polynomials $J_\lambda^{(\alpha)}$:

- symmetric functions introduced by Jack;
- generalization of Schur symmetric function (for $\alpha = 1$);
- special case of Macdonald polynomials

Expand Jack polynomial in power-sum symmetric basis:

$$J_\lambda^{(\alpha)} = \sum_{\substack{\rho: \\ |\rho|=|\lambda|}} \theta_\rho^{(\alpha)}(\lambda) p_\rho.$$

We call quantities $\theta_\rho^{(\alpha)}(\lambda)$ **Jack characters** (for $\alpha = 1$ they coincide with the irreducible characters of the symmetric groups up to some normalization constant).

Characterization of Jack measure

$$\mathbb{E}_{\mathbb{P}_n^{(\alpha)}}(\theta_{\mu}^{(\alpha)}) = \begin{cases} 1 & \text{if } \mu = \mathbf{1}^n, \\ 0 & \text{otherwise.} \end{cases}$$

Let $\lambda \in \mathbb{Y}_n$.

Proposition

$\int_{\mathbb{R}} x^k \Delta(A_{\alpha}(\lambda))(x) dx$ can be expressed as a function of

$$\theta_{(\mathbf{1}^n)}^{(\alpha)}, \theta_{(2, \mathbf{1}^{n-2})}^{(\alpha)}, \dots, \theta_{(k-1, \mathbf{1}^{n-k+1})}^{(\alpha)}.$$

Corollary

Our central limit theorem has equivalent, algebraic version!

Algebraic central limit theorem

Theorem (D., Féray)

Choose a sequence $(\Xi_k)_{k=2,3,\dots}$ of independent standard Gaussian random variables. As $n \rightarrow \infty$, we have:

$$\left(\frac{\sqrt{k} \theta_{(k, 1^{n-k})}^{(\alpha)}(\lambda(n))}{n^{k/2}} \right)_{k=2,3,\dots} \xrightarrow{d} (\Xi_k)_{k=2,3,\dots},$$

where the distribution of $\lambda(n)$ is Jack measure of size n and where \xrightarrow{d} means convergence in distribution of the finite-dimensional law.

We can prove this theorem using algebraic methods (Jack characters after normalization span a very nice algebra)!

Polynomials functions

We define

$$\text{Ch}_\mu^{(\alpha)}(\lambda) = \begin{cases} \alpha^{-\frac{|\mu|-\ell(\mu)}{2}} \binom{|\lambda|-|\mu|+m_1(\mu)}{m_1(\mu)} z_\mu \theta_{\mu,1^{|\lambda|-|\mu|}}^{(\alpha)}(\lambda) & \text{if } |\lambda| \geq |\mu|; \\ 0 & \text{if } |\lambda| < |\mu|, \end{cases}$$

where

- $z_\mu = \mu_1 \mu_2 \cdots m_1(\mu)! m_2(\mu)! \cdots$,
- $m_i(\mu)$ - number of parts of μ equal to i .

Theorem (Lassalle, 2009)

The family $(\text{Ch}_\mu^{(\alpha)})_\mu$ span linearly an algebra $\Lambda_\star^{(\alpha)}$ of α -shifted symmetric functions.

What do we have and what do we miss?

In order to prove our main theorem:

- We want to estimate mixed moments of Jack characters;
- Expectation of the Jack characters is easy to compute;
- Suitably normalized Jack characters span linearly some nice algebra $\Lambda_{\star}^{(\alpha)}$;
- We want to expand a product:

$$\text{Ch}_{\mu}^{(\alpha)} \text{Ch}_{\nu}^{(\alpha)} = \sum_{\rho} g_{\mu, \nu; \pi}^{(\alpha)} \text{Ch}_{\pi}^{(\alpha)}$$

as a linear combination of suitably normalized Jack characters.

Problem

What can we say about $g_{\mu, \nu; \pi}^{(\alpha)}$?

Main result for structure constants

Theorem (D., Féray)

Let

$$\text{Ch}_\mu^{(\alpha)} \text{Ch}_\nu^{(\alpha)} = \sum_{\rho} g_{\mu,\nu;\pi}^{(\alpha)} \text{Ch}_\pi^{(\alpha)}.$$

Then, *structure constants* $g_{\mu,\nu;\pi}^{(\alpha)}$ are polynomials in $\gamma := \alpha^{1/2} - \alpha^{-1/2}$ of degree less than

$$\min_{i=1,2,3} (n_i(\mu) + n_i(\nu) - n_i(\pi)),$$

with rational coefficients, where $n_i(\lambda)$ - natural valued function of λ .

- It is crucial for proving central limit theorem;
- It is applicable to different problems.

Projection on the set of Young diagrams of a fixed size

Let $\mu, \nu, \pi \in \mathbb{Y}_n$.

$$\theta_{\mu}^{(\alpha)}(\lambda) \theta_{\nu}^{(\alpha)}(\lambda) = \sum_{|\pi|=n} c_{\mu, \nu; \pi}^{(\alpha)} \theta_{\pi}^{(\alpha)}.$$

Hence

$$c_{\mu, \nu; \pi}^{(\alpha)} = \frac{\alpha^{d(\mu, \nu; \pi)/2}}{z_{\tilde{\mu}} z_{\tilde{\nu}}} \sum_{0 \leq i \leq m_1(\pi)} g_{\tilde{\mu}, \tilde{\nu}; \tilde{\pi} 1^i}^{(\alpha)} \cdot z_{\tilde{\pi}} \cdot i! \cdot \binom{n - |\tilde{\pi}|}{i},$$

where

- $\tilde{\mu}$ is created from μ by removing all parts equal to 1,
- $d(\mu, \nu; \pi) = |\mu| - \ell(\mu) + |\nu| - \ell(\nu) - (|\pi| - \ell(\pi))$.

$\alpha = 1$ - Structure constants of the $Z(\mathbb{C}[\mathfrak{S}_n])$

Let $\mathbb{C}[\mathfrak{S}_n] := \{f : \mathfrak{S}_n \rightarrow \mathbb{C}\}$ be a **group algebra of the symmetric group**. This is algebra with the multiplication defined by:

$$f \cdot g(\sigma) := \sum_{\sigma_1 \sigma_2 = \sigma} f(\sigma_1)g(\sigma_2).$$

Let

$$Z(\mathbb{C}[\mathfrak{S}_n]) := \{f \in \mathbb{C}[\mathfrak{S}_n] : \forall g \in \mathbb{C}[\mathfrak{S}_n], fg = gf\}$$

be the **center** of that algebra. It has a basis $(f_\mu)_{|\mu|=n}$, where

$$f_\mu(\sigma) = \begin{cases} 1 & \text{if } \sigma \text{ has cycle type } \mu, \\ 0 & \text{otherwise.} \end{cases}$$

$\alpha = 1$ - Structure constants of the $Z(\mathbb{C}[\mathfrak{S}_n])$

Let

$$f_\mu f_\nu = \sum_{|\rho|=n} c_{\mu,\nu;\rho} f_\rho.$$

Lemma

The structure constant $c_{\mu,\nu;\rho}$ is equal to the number of pairs of permutation (σ_1, σ_2) such that

- σ_1 has cycle type μ ,
- σ_2 has cycle type ν ,
- $\sigma_1 \sigma_2 = \sigma$, where σ is a fixed permutation of the cycle-type ρ .

$\alpha = 1$ - Structure constants of the $Z(\mathbb{C}[\mathfrak{S}_n])$

One has a following relation:

$$c_{\mu,\nu;\rho}^{(1)} = c_{\mu,\nu;\rho}.$$

Remark

From the previous theorem and a relation between $c^{(\alpha)}$ and $g^{(\alpha)}$ one can deduce a classical result of Farahat and Higman: $c_{\mu\mathbf{1}^{n-|\mu|},\nu\mathbf{1}^{n-|\nu|};\rho\mathbf{1}^{n-|\rho|}}$ is a polynomial in n .

$\alpha = 2$ - Structure constants of the Hecke algebra of (\mathfrak{S}_{2n}, H_n)

Let \mathfrak{S}_{2n} acts on the set $X_n := \{1, \bar{1}, \dots, n, \bar{n}\}$ by permutations and let

$$\mathfrak{S}_{2n} > H_n := \{\sigma \in \mathfrak{S}_{2n} : \forall i \in X_n \sigma(\bar{i}) = \sigma(i)\}$$

be a **hyperoctahedral subgroup**.

Hecke algebra $\mathbb{C}[H_n \backslash \mathfrak{S}_{2n} / H_n] < \mathbb{C}[\mathfrak{S}_{2n}]$ of the pair (\mathfrak{S}_{2n}, H_n) is defined by:

$$\mathbb{C}[H_n \backslash \mathfrak{S}_{2n} / H_n] := \{x \in \mathbb{C}[\mathfrak{S}_{2n}] : hxh' = x \forall h, h' \in H_n\}.$$

Double-cosets: equivalence classes for the relation $x \sim hxh'$ (for $x \in \mathfrak{S}_{2n}$ and $h, h' \in H_n$)

- naturally indexed by **partitions of size n** ;
- $F_\mu = \sum_{x \text{ of type } \mu} \delta_x$ - linear basis of $\mathbb{C}[H_n \backslash \mathfrak{S}_{2n} / H_n]$.

$\alpha = 2$ - Structure constants of the Hecke algebra of (\mathfrak{S}_{2n}, H_n)

Let

$$F_\mu F_\nu = \sum_{|\rho|=n} h_{\mu,\nu;\rho} F_\rho.$$

Then

$$c_{\mu,\nu;\rho}^{(2)} = \frac{h_{\mu,\nu;\rho}}{2^n n!}.$$

Remark

From the previous theorem and a relation between $c^{(\alpha)}$ and $g^{(\alpha)}$ one can deduce a result of Tout (2013):

$$\frac{h_{\mu 1^{n-|\mu|}, \nu 1^{n-|\nu|}; \pi 1^{n-|\pi|}}}{n! 2^n}$$

is a polynomial in n .

Fin

MERCI