

# Advanced Graph Theory (4TiN917U)

Marthe BONANZY

## Lectures 5 and 8: Discharging and Fixed Parameter Tractability (Or how to get rid of useless stuff)


Flashback to the previous lecture: complexity to recognize if a graph is planar? Poly-time since characterized by finitely many forbidden minors.

Example of infinitely many pairwise incomparable (for the induced subgraph relation) trees? 


Consider a problem (say  $k$ -coloring) and an instance of it.

Goal: Simplify the instance as much as possible.

Examples:



: Vertices of degree  $5k-1$  can be ignored (deleted).



: In any solution, u and v have the same color as merge u and v into a single vertex.

Things to worry about:

- Did I lose information? (ex: there is a vertex adjacent to all the others, and the graph is planar)
- Did I modify the problem? (ex:  $k$ -coloring  $\rightarrow$   $(k-1)$ -coloring)
- Did I leave the graph class? (ex: the graph is planar and I merged vertices)

New goal: Prove that all planar graphs are 5-colorable.


21

Problem: 5-coloring      Instance: A planar graph  $G$ .

① Vertices of degree  $\leq 4$  are irrelevant for the problem  
(can be deleted)

② Vertices of degree 5 are also irrelevant.

Proof 1:



If  $G$  contains  $u$  with  $d(u) = 5$

Consider assume  $G \setminus u$  admits a 5-coloring of  
Consider a planar embedding of  $G$  and let  $v_1, \dots, v_5$  be the neighbors of  $u$  in clockwise order.  
If  $\{v_1, v_2, v_3, v_4, v_5\}$  use  $\leq 4$  colors in  $\alpha$ ,

then  $\alpha$  extends to  $G$ . Otherwise, assume without loss of generality  
that  $d(v_i) = i \quad \forall i \in \{1, 2, 3, 4, 5\}$ . Consider a ~~Kempe~~ Kempe  
chain (1,3) on vertex  $v_1$  (Remember, this is a maximal component colored 1 and 3  
in  $\alpha$  that contains  $v_1$ ). If it does not contain  $v_3$ , we can swap the  
two colors and go back to the previous case. So there is a path  
colored 1-3 between  $v_1$  and  $v_3$  (possibly  $v_1v_3$  is an edge). By  
the same analysis, there is a path (possibly just an edge) colored 2-4  
between  $v_2$  and  $v_4$ . The two paths are vertex-disjoint since they  
use disjoint colors. Since this contradicts the planarity of the embedding  
 $G$ .

Proof 2: If  $v_i v_j$  is an edge  $\wedge i \neq j$ , then  $G$  contains a degenerate  
5 vertices  $\Rightarrow$  a contradiction with planarity. Therefore, if  $i \neq j$   
contracting the edges  $v_i u$  and  $v_j u$ .  $G'$  is planar since planarity  
is closed under taking minors. Assume  $G'$  admits a 5-coloring  $\alpha$ .  
We derive a 5-coloring  $\alpha'$  of  $G \setminus u$  that uses  $\leq 4$  colors on  $\{v_1, \dots, v_5\}$ , hence the conclusion.

# Advanced Graph Theory (4TIN917U)


Markus BONAMY


## Lectures 5 and 8: Discharging and Fixed Parameter Tractability (Or how to get rid of useless stuff)

Flashback to the previous lecture: complexity to recognize if a graph is planar? Poly-time since characterized by finitely many forbidden minors. Example of infinitely many pairwise incomparable (for the induced subgraph relation) trees?  $\mathcal{X}, \mathcal{Y}, \mathcal{Z}, \mathcal{W}, \mathcal{V}, \dots$

Consider a problem (say  $k$ -coloring) and an instance of it.

Goal: Simplify the instance as much as possible.

Example:  Vertices of degree  $\leq k-1$  can be ignored (deleted).

 digraph  $k-1$  vertices.

In any solution,  $u$  and  $v$  have the same color as merge  $u$  and  $v$  into a single vertex.

Things to worry about:

- Did I lose information? (ex: there is a vertex adjacent to all the others, and the graph is planar)
- Did I modify the problem? (ex:  $k$ -coloring  $\rightarrow (k-1)$ -coloring)
- Did I leave the graph class? (ex: the graph is planar and I merged vertices)

It suffices now to argue that all planar graphs whose vertices all have degree  $\geq 6$  are 5-colorable. 3/

We use

$\Delta$  Euler's formula: for every planar embedding  $(V, E, F)$  of a planar graph  $G$ , we have  $|V| - |E| + |F| \geq 0$

\* (in fact  $= 1 + \# \text{ connected components}$ )

The traditional proof is by easy induction, here we show an of and prof following a discharging argument.

$\Delta$  Discharging = Double counting

"Silly" example: Gauss' formula for  $\sum_{i=1}^m i$

Weight assignment  $w: i \mapsto i$

Discharging rule:  $\forall i < \frac{m+1}{2}$  gives all its weight to the element  $m+1-i$ .

Let  $w'$  be the final weight (after application of the discharging rule).

$$\forall i < \frac{m+1}{2}, w'(i) = 0$$

$$\forall i > \frac{m+1}{2}, w'(i) = m+1$$

$$\text{If } m \text{ is odd, } w'\left(\frac{m+1}{2}\right) = w\left(\frac{m+1}{2}\right) = \frac{m+1}{2}.$$

$$\sum_{i=1}^m i = \sum_{i=1}^m w(i) = \underbrace{\sum_{i=1}^{\frac{m+1}{2}} w(i)}_{0+0+0+4+8+8+8 \leftarrow w'} + \underbrace{\sum_{i=\frac{m+1}{2}+1}^m w(i)}_{m+1+m+1+\dots+m+1 \leftarrow w'} = \frac{m(m+1)}{2}$$

Easier to count

$$1 \quad 2 \quad 3 \quad 4 \quad 5 \quad 6 \quad 7 \leftarrow w \\ 0 \quad 0 \quad 0 \quad 4 \quad 8 \quad 8 \quad 8 \leftarrow w'$$

## Euler's Formula

4/


$$|V| - |E| + |F| = \sum_{v \in V} 1 + \sum_{e \in E} (-1) + \sum_{f \in F} 1$$

Weight assignment:  $\forall v \in V, w(v) = 1$

$\forall e \in E, w(e) = -1$

$\forall f \in F, w(f) = 1$

Consider ~~an orientation~~ for which no edge is perfectly horizontal.




Axios

Right

Discharging rule: Every vertex and every edge gives all its weight to the face on its right.

$w'$  the resulting weight assignment.



$\forall v \in V, w'(v) = 0$

$\forall e \in E, w'(e) = 0$

$\forall f \in F$  that is an internal face,  $w'(f) = 0$

$w'(\text{external face}) = 1 + \# \text{ connected components}$

We have

$$|V|-|E|+|F| = \sum_{x \in V \cup E \cup F} w(x) = \sum_{x \in V \cup E \cup F} w'(x) = w'(\text{external face}) = 1 + \# \text{ connected components} > 0$$

Note that

$$\sum_{v \in V} (d(v) - 6) + \sum_{f \in F} (2d(f) - 6)$$

$$= 2|E| - 6|V| + 4|E| - 6|F| < 0$$

Note that as soon as  $G$  contains at least two edges

we have  $d(f) \geq 3 \quad \forall f \in F$ .

Therefore  $\sum_{f \in F} (2d(f) - 6) \geq 0$ .

If every vertex has degree at least 6,

$$\text{then } \sum_{v \in V} (d(v) - 6) \geq 0$$

$$\text{A contradiction since } \sum_{v \in V} (d(v) - 6) + \sum_{f \in F} (2d(f) - 6) < 0.$$

Consequently, every planar graph  $G$  with  $V(G) \neq \emptyset$  contains a vertex of degree  $\leq 5$ .

This implies that all planar graphs are 5-colorable, as desired.

Can be seen as a proof by induction

Usually:  $H_0$  holds. If  $H_m$  holds then  $H_{m+1}$  holds.  
Here: the property holds for the empty graph. If  $G$  contains a vertex  $v$  of degree  $\leq 5$ , then the property for  $G \setminus v$  implies the property for  $G$

Work to guarantee all graphs are built

A finer discharging argument.

Goal: Every planar graph  $G$  with  $V(G) \neq \emptyset$

contains

(a) a vertex of degree  $\leq 4$ , or

(b) a vertex of degree 5 adjacent to a vertex of degree  $\leq 6$ .

$(V, E, F)$  a planar embedding of  $G$ .



Proof:  $w(v) := d(v) - 6 \quad \forall v \in V(G)$

$w(f) := 2d(f) - 6 \quad \forall f \in F$ .

Then  $G$  contains at least two edges, so  $\forall f \in F, 2d(f) - 6 > 0$ .

$\sum_{x \in V \cup F} w(x) < 0$  by Euler's formula.

Discharging rules:



$$d(f) \geq 4$$

Let  $w'$  be the resulting weight assignment.

Assume for a contradiction that  $V(G) \neq \emptyset$  and  $G$  contains neither (a) nor (b).

Let  $v \in V$ .  $d(v) \geq 5$ .

If  $d(v)=5$ ,  $w(v)=-1$ . If  $v$  is incident to a face of degree  $\geq 4$ , by R<sub>2</sub>  $v$  receives 1 at least, so  $w'(v) \geq 0$ . If  $v$  is only incident to triangles, then all its neighbors have degree 3, 7 so R<sub>1</sub> applies 5 times, here  $v$  receives  $5 \times \frac{1}{3} = 1$  so  $w'(v) \geq 0$ .

If  $d(v) = 6$ , no rule applying so  $w'(v) = w(v) = 0$ .

If  $d(v) \geq 3$  by R<sub>1</sub> and since there is no edge (5)-5 by assumption on G, R<sub>1</sub> is applied at most  $\left\lfloor \frac{d(v)}{2} \right\rfloor$  times on v.

Therefore  $w'(v) \geq w(v) - \frac{1}{5} \times \left\lfloor \frac{d(v)}{2} \right\rfloor \geq \frac{9}{10}d(v) - 6$

Note that  $w'(v) \geq 0$  when  $d(v) \geq 7$ .

Therefore,  $\forall v \in V, w'(v) \geq 0$

Let  $f \in F$ .

If  $d(f) = 3$  no rule applies so  $w'(f) = w(f) = 0$

If  $d(f) \geq 4$  since there is no edge 5-5, R<sub>2</sub> is applied at most  $\frac{d(f)}{2}$  times.

Hence  $w'(f) \geq w(f) - 1 \times \frac{d(f)}{2} \geq \frac{3d(f)}{2} - 6$

Note that  $w'(f) \geq 0$  when  $d(f) \geq 4$ .

Consequently,  $\sum_{x \in V \cup F} w'(x) \geq 0$

Since  $\sum_{x \in V \cup F} w(x) = \sum_{x \in V \cup F} w(x) < 0$ , we reach a contradiction. D.

## Fixed Parameter Tractability

Problem is FPT w.r.t parameter  $\lambda$   
 if there is an algorithm in time  $f(\lambda) \cdot \text{poly}(\text{instance})$   
 for some computable function  $f$ .


Equivalently, the problem admits a kernel of size  $g(\lambda)$   
 that can be computed in  $\text{poly}(\text{instance})$ .

Example: Vertex Cover parametrized by the size  
 of the solution?

Given  $(G, \lambda)$ , does there exist  $S \subseteq V(G)$  s.t.

$$\left\{ \begin{array}{l} |S| \leq \lambda \\ \forall u \in V(G), \{u, v\} \cap S \neq \emptyset \end{array} \right.$$

Reduction 1: If  $G$  contains a vertex  $u$  of degree 0,  
 return  $(G \setminus \{u\}, \lambda)$ .

Reduction 2:  If  $G$  contains a vertex  $v$  of degree  $k$ ,  
 return  $(G \setminus \{v\}, \lambda - 1)$ .  
 "v has to be in the solution if any  
 "v has to be in the solution if any