


Coloring the squares of graphs


Marthe Bonamy

Clermont-Ferrand
April 17, 2014






Square coloring



χ : Minimum number of colors to ensure that

$$\textcircled{a} \text{---} \textcircled{b} \Rightarrow a \neq b.$$

Square coloring




χ : Minimum number of colors to ensure that

$$\textcircled{a} \text{---} \textcircled{b} \Rightarrow a \neq b.$$

χ^2 : Minimum number of colors to ensure that

$$\textcircled{a} \text{---} \textcircled{b} \text{ or } \textcircled{a} \text{---} \bigcirc \text{---} \textcircled{b} \Rightarrow a \neq b.$$



χ : Minimum number of colors to ensure that

$$\textcircled{a} \text{---} \textcircled{b} \Rightarrow a \neq b.$$

χ^2 : Minimum number of colors to ensure that

$$\textcircled{a} \text{---} \textcircled{b} \text{ or } \textcircled{a} \text{---} \bigcirc \text{---} \textcircled{b} \Rightarrow a \neq b.$$

Δ : Maximum degree of the graph.

$$\Delta^2 + 1 \geq \chi^2 \geq \Delta + 1.$$

A graph is **planar** if it can be drawn in a plane without crossing edges.

Conjecture (Wegner 1977)

If G is a *planar* graph, then:

- $\chi^2 \leq 7$ if $\Delta = 3$
- $\chi^2 \leq \Delta + 5$ if $4 \leq \Delta \leq 7$
- $\chi^2 \leq \lfloor \frac{3\Delta}{2} \rfloor + 1$ if $\Delta \geq 8$

A graph is **planar** if it can be drawn in a plane without crossing edges.

Conjecture (Wegner 1977)

If G is a *planar* graph, then:

- $\chi^2 \leq 7$ if $\Delta = 3$
- $\chi^2 \leq \Delta + 5$ if $4 \leq \Delta \leq 7$
- $\chi^2 \leq \lfloor \frac{3\Delta}{2} \rfloor + 1$ if $\Delta \geq 8$

✓ C. Thomassen

A graph is **planar** if it can be drawn in a plane without crossing edges.

Conjecture (Wegner 1977)

If G is a *planar* graph, then:

- $\chi^2 \leq 7$ if $\Delta = 3$
- $\chi^2 \leq \Delta + 5$ if $4 \leq \Delta \leq 7$
- $\chi^2 \leq \lfloor \frac{3\Delta}{2} \rfloor + 1$ if $\Delta \geq 8$


✓ C. Thomassen

Sparse planar graphs

g : girth = minimum length of a cycle.


Sparse planar graphs

g : girth = minimum length of a cycle.



Sparse planar graphs

g : girth = minimum length of a cycle.



Theorem (Bu, Zhu '11)

Wegner's conjecture is *true* for planar graphs with $g \geq 6$.

Sparse planar graphs (2)

Question (Wang, Lih '01)

$$g \geq k, \Delta \geq d \Rightarrow \chi^2 = \Delta + 1.$$

Sparse planar graphs (2)

Question (Wang, Lih '01)

In the case of planar graphs,
 $\exists? k_0, \forall k \geq k_0, \exists d,$
 $g \geq k, \Delta \geq d \Rightarrow \chi^2 = \Delta + 1.$


Sparse planar graphs (2)

Question (Wang, Lih '01)

In the case of *planar* graphs,
 $\exists? k_0, \forall k \geq k_0, \exists d,$
 $g \geq k, \Delta \geq d \Rightarrow \chi^2 = \Delta + 1.$

Theorem (Borodin, Glebov, Ivanova, Neustroeva, Tashkinov '04)

False for $k \leq 6$. *True* for $k \geq 7$.



Theorem (Borodin, Ivanova, Neustroeva '08 & Ivanova '11)

For any *planar* graph, $\chi^2 = \Delta + 1$ in each of the following cases :

- $g \geq 24$ and $\Delta \geq 3$
- $g \geq 15$ and $\Delta \geq 4$
- $g \geq 12$ and $\Delta \geq 5$
- $g \geq 10$ and $\Delta \geq 6$
- $g \geq 8$ and $\Delta \geq 10$
- $g \geq 7$ and $\Delta \geq 16$

Theorem (Borodin, Ivanova, Neustroeva '08 & Ivanova '11)

For any *planar* graph, $\chi^2 = \Delta + 1$ in each of the following cases :

- $g \geq 24$ and $\Delta \geq 3$
- $g \geq 15$ and $\Delta \geq 4$
- $g \geq 12$ and $\Delta \geq 5$
- $g \geq 10$ and $\Delta \geq 6$
- $g \geq 8$ and $\Delta \geq 10$
- $g \geq 7$ and $\Delta \geq 16$

$$|E| - |V| - |F| = -2$$

(proof)

Theorem (Borodin, Ivanova, Neustroeva '08 & Ivanova '11)

For any *planar* graph, $\chi^2 = \Delta + 1$ in each of the following cases :

- $g \geq 24$ and $\Delta \geq 3$
- $g \geq 15$ and $\Delta \geq 4$
- $g \geq 12$ and $\Delta \geq 5$
- $g \geq 10$ and $\Delta \geq 6$
- $g \geq 8$ and $\Delta \geq 10$
- $g \geq 7$ and $\Delta \geq 16$

$$|E| - |V| - |F| = -2$$

(proof)

$$2|E| - g|F| + (g - 2)|E| - g|V| < 0$$

Theorem (Borodin, Ivanova, Neustroeva '08 & Ivanova '11)

For any *planar* graph, $\chi^2 = \Delta + 1$ in each of the following cases :

- $g \geq 24$ and $\Delta \geq 3$
- $g \geq 15$ and $\Delta \geq 4$
- $g \geq 12$ and $\Delta \geq 5$
- $g \geq 10$ and $\Delta \geq 6$
- $g \geq 8$ and $\Delta \geq 10$
- $g \geq 7$ and $\Delta \geq 16$

$$|E| - |V| - |F| = -2$$

(proof)

$$2|E| - g|F| + (g - 2)|E| - g|V| < 0$$

$$\sum_{f \in F} \underbrace{(d(f) - g)}_{\geq 0} + c \cdot \sum_{v \in V} \left(d(v) - \frac{2g}{g-2} \right) < 0$$

Another measure of sparseness

ad: average degree = $\frac{\sum d(v)}{|V|} = \frac{2|E|}{|V|}$.

Another measure of sparseness

ad: average degree = $\frac{\sum d(v)}{|V|} = \frac{2|E|}{|V|}$.

mad: maximum average degree = $\max \frac{2|E(H)|}{|V(H)|}$.

Another measure of sparseness

ad: average degree = $\frac{\sum d(v)}{|V|} = \frac{2|E|}{|V|}$.

mad: maximum average degree = $\max \frac{2|E(H)|}{|V(H)|}$.

Lemma (Derived from Euler's Formula)

For any *planar* graph, $(\text{mad} - 2)(g - 2) < 4$

Another measure of sparseness

ad: average degree = $\frac{\sum d(v)}{|V|} = \frac{2|E|}{|V|}$.

mad: maximum average degree = $\max \frac{2|E(H)|}{|V(H)|}$.

Lemma (Derived from Euler's Formula)

For any *planar* graph, $(\text{mad} - 2)(g - 2) < 4$

Question

$$\text{mad} < m, \Delta \geq d \Rightarrow \chi^2 = \Delta + 1.$$

Another measure of sparseness

ad: average degree = $\frac{\sum d(v)}{|V|} = \frac{2|E|}{|V|}$.

mad: maximum average degree = $\max \frac{2|E(H)|}{|V(H)|}$.

Lemma (Derived from Euler's Formula)

For any *planar* graph, $(\text{mad} - 2)(g - 2) < 4$

Question

M maximal | $\forall m < M, \exists d,$
 $\text{mad} < m, \Delta \geq d \Rightarrow \chi^2 = \Delta + 1.$

First results on non-planar sparse graphs

Theorem (B., Lévêque, Pinlou '11, Cranston, Škrekovski '13)

For *any* graph, $\chi^2 = \Delta + 1$ in each of the following cases :

• ~~$g \geq 24$ and $\Delta \geq 3$~~

• $\underbrace{mad < \frac{7}{3}}_{g \geq 15 \text{ } 14}$ and $\Delta \geq 4$

• $\underbrace{mad < \frac{18}{7}}_{g \geq 9}$ and $\Delta \geq 8$

• $\underbrace{mad < \frac{70}{29}}_{g \geq 12}$ and $\Delta \geq 5$

• $\underbrace{mad < \frac{8}{3}}_{g \geq 8}$ and $\Delta \geq 14 > 10$

• $\underbrace{mad < \frac{5}{2}}_{g \geq 10}$ and $\Delta \geq 6$

• $\underbrace{mad < \frac{14}{5}}_{g \geq 7}$ and $\Delta \geq 1200 \gg 16$

First results on non-planar sparse graphs

Theorem (B., Lévêque, Pinlou '11, Cranston, Škrekovski '13)

For *any* graph, $\chi^2 = \Delta + 1$ in each of the following cases :

• ~~$g \geq 24$ and $\Delta \geq 3$~~

• $mad < \frac{7}{3}$ and $\Delta \geq 4$
 $\underbrace{\hspace{10em}}_{g \geq 15 \text{ } 14}$

• $mad < \frac{18}{7}$ and $\Delta \geq 8$
 $\underbrace{\hspace{10em}}_{g \geq 9}$

• $mad < \frac{70}{29}$ and $\Delta \geq 5$
 $\underbrace{\hspace{10em}}_{g \geq 12}$

• $mad < \frac{8}{3}$ and $\Delta \geq 14 > 10$
 $\underbrace{\hspace{10em}}_{g \geq 8}$

• $mad < \frac{5}{2}$ and $\Delta \geq 6$
 $\underbrace{\hspace{10em}}_{g \geq 10}$

• $mad < \frac{14}{5}$ and $\Delta \geq 1200 \gg 16$
 $\underbrace{\hspace{10em}}_{g \geq 7}$

What about $mad < 3$?

First results on non-planar sparse graphs (2)

Theorem (B., Lévêque, Pinlou '13)

For a small enough ϵ , $\text{mad} < 3 - \epsilon$ and $\Delta \geq \frac{3}{\epsilon^2} \Rightarrow \chi^2 = \Delta + 1$.

First results on non-planar sparse graphs (2)

Theorem (B., Lévêque, Pinlou '13)

For a small enough ϵ , $\text{mad} < 3 - \epsilon$ and $\Delta \geq \frac{3}{\epsilon^2} \Rightarrow \chi^2 = \Delta + 1$.

Theorem (Borodin, Kostochka, Woodall '97)

For any *bipartite multigraph* G , if L is an edge list assignment such that $|L(u, v)| \geq d(u), d(v)$ for every (u, v) , then G is *L-edge-choosable*.

First results on non-planar sparse graphs (2)

Theorem (B., Lévêque, Pinlou '13)

For a small enough ϵ , $\text{mad} < 3 - \epsilon$ and $\Delta \geq \frac{3}{\epsilon^2} \Rightarrow \chi^2 = \Delta + 1$.


Theorem (Borodin, Kostochka, Woodall '97)

For any *bipartite multigraph* G , if L is an edge list assignment such that $|L(u, v)| \geq d(u), d(v)$ for every (u, v) , then G is *L-edge-choosable*.

\Rightarrow Global discharging method.


List coloring?

χ_ℓ^2 : Minimum size of every $L(v)$ such that



List coloring?


χ_ℓ^2 : Minimum size of every $L(v)$ such that



$$\Delta^2 + 1 \geq \chi_\ell^2 \geq \chi^2 \geq \Delta + 1.$$

List coloring?

χ_ℓ^2 : Minimum size of every $L(v)$ such that



$$\Delta^2 + 1 \geq \chi_\ell^2 \geq \chi^2 \geq \Delta + 1.$$

What changes in the previous theorems?


Conjecture (Kostochka, Woodall '01)

For any graph, $\chi^2 = \chi_l^2$.

List coloring.

Conjecture (Kostochka, Woodall '01)


For any graph, $\chi^2 = \chi_\ell^2$.



List coloring.

Conjecture (Kostochka, Woodall '01)

For any graph, $\chi^2 = \chi_\ell^2$.




Theorem (Kim, Park '13)

$\chi_\ell^2 - \chi^2$ can be *arbitrarily large*.

List coloring.

Conjecture (Kostochka, Woodall '01)

For any graph, $\chi^2 = \chi_\ell^2$.



Theorem (Kim, Park '13)

$\chi_\ell^2 - \chi^2$ can be *arbitrarily large*.

Theorem (Kim, Kwon, Park '13, Kosar, Petrickova, Reiniger, Yeager '13)

For any k , $\chi_\ell^k - \chi^k$ can be *arbitrarily large*.

$$\Delta^2 + 1 \geq \chi_\ell^2 \geq \chi^2 \geq \Delta + 1.$$

Conjecture (Cranston, Kim '08)

If $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^2 = \Delta^2$.

Cranston, Kim '08: **True** for $\Delta = 3$.

Upper bounds

Conjecture (Cranston, Kim '08)

If $\Delta \geq 3$, *no* graph satisfies $\chi_l^2 = \Delta^2$.

Cranston, Kim '08: **True** for $\Delta = 3$.

Conjecture (Borodin, Kostochka '77)

If $\Delta \geq 9$ and $\omega \leq \Delta - 1$, then $\chi_l \leq \Delta - 1$.

Upper bounds


Conjecture (Cranston, Kim '08)

If $\Delta \geq 3$, *no* graph satisfies $\chi_l^2 = \Delta^2$.

Cranston, Kim '08: **True** for $\Delta = 3$.

Conjecture (Borodin, Kostochka '77)

If $\Delta \geq 9$ and $\omega \leq \Delta - 1$, then $\chi_l \leq \Delta - 1$.



Upper bounds

Conjecture (Cranston, Kim '08)

If $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^2 = \Delta^2$.

Cranston, Kim '08: **True** for $\Delta = 3$.

Conjecture (Borodin, Kostochka '77)

If $\Delta \geq 9$ and $\omega \leq \Delta - 1$, then $\chi_\ell \leq \Delta - 1$.

Conjecture (Miao, Fan '13)

For any $k \geq 2$, and $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^k = D(k, \Delta)$.

Upper bounds

Conjecture (Cranston, Kim '08)

If $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^2 = \Delta^2$.

Cranston, Kim '08: **True** for $\Delta = 3$.

Conjecture (Borodin, Kostochka '77)

If $\Delta \geq 9$ and $\omega \leq \Delta - 1$, then $\chi_\ell \leq \Delta - 1$.

Conjecture (Miao, Fan '13)

For any $k \geq 2$, and $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^k = D(k, \Delta)$.

Theorem (B., Bousquet '13)

For any $k \geq 3$, and $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^k = D(k, \Delta)$.

Upper bounds

Conjecture (Cranston, Kim '08)

If $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^2 = \Delta^2$.

Cranston, Kim '08: **True** for $\Delta = 3$.

Conjecture (Borodin, Kostochka '77)

If $\Delta \geq 9$ and $\omega \leq \Delta - 1$, then $\chi_\ell \leq \Delta - 1$.

Conjecture (Miao, Fan '13)

For any $k \geq 2$, and $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^k = D(k, \Delta)$.

Theorem (B., Bousquet '13)

For any $k \geq 3$, and $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^k \geq D(k, \Delta)$.

Upper bounds

Conjecture (Cranston, Kim '08)

If $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^2 = \Delta^2$.

Cranston, Kim '08: **True** for $\Delta = 3$.

Conjecture (Borodin, Kostochka '77)

If $\Delta \geq 9$ and $\omega \leq \Delta - 1$, then $\chi_\ell \leq \Delta - 1$.

Conjecture (Miao, Fan '13)

For any $k \geq 2$, and $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^k = D(k, \Delta)$.

Theorem (B., Bousquet '13)

For any $k \geq 3$, and $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^k \geq D(k, \Delta)$.

Theorem (Cranston, Rabern '13)

If $\Delta \geq 3$, *no* graph satisfies $\chi_\ell^2 = \Delta^2$.

Conjecture

For any $k \geq 3$, and $\Delta \geq 3$, *no* graph satisfies

$$\chi_l^k \geq D(k, \Delta) - k + 2.$$

Conclusion

Conjecture

For any $k \geq 3$, and $\Delta \geq 3$, no graph satisfies

$$\chi_k \geq D(k, \Delta) - k + 2.$$

Conjecture

For any C and $\epsilon > 0$,

$$\text{mad} < \frac{4C+2}{C+1} - \epsilon \text{ and } \Delta \geq f(C, \epsilon) \Rightarrow \chi_k^2 \leq \Delta + C.$$

