



An overview of discharging methods

Marthe Bonamy

May 6th, 2014








χ : Minimum number of colors to ensure that

$$\textcircled{a} - \textcircled{b} \Rightarrow a \neq b.$$



χ : Minimum number of colors to ensure that

$$\textcircled{a} \text{---} \textcircled{b} \Rightarrow a \neq b.$$

A graph is **planar** if it can be drawn in a plane without crossing edges.

- Introduced by Wernicke in 1904.

- Introduced by Wernicke in 1904.
- Goal:

Conjecture (Guthrie 1852)

For every planar graph, $\chi \leq 4$.

- Introduced by Wernicke in 1904.
- Goal:

Conjecture (Guthrie 1852)

For every *planar* graph, $\chi \leq 4$.

- Successful?

- Introduced by Wernicke in 1904.
- Goal:

Conjecture (Guthrie 1852)

For every *planar* graph, $\chi \leq 4$.

- Successful? ✓

- 1 Every planar graph contains some $\mathcal{C} \in \{C_1, C_2, \dots, C_p\}$.

Discharging, the outline

- 1 Every planar graph contains some $C \in \{C_1, C_2, \dots, C_p\}$.
- 2 For the 4-coloring of planar graphs, a graph that contains a C_i can be reduced to a smaller graph.

Discharging, the outline

- 1 Every planar graph contains some $C \in \{C_1, C_2, \dots, C_p\}$.
- 2 For the 4-coloring of planar graphs, a graph that contains a C_i can be reduced to a smaller graph.
- 3 Thus every planar graph is 4-colorable.

Discharging, the outline

- 1 Every planar graph contains some $C \in \{C_1, C_2, \dots, C_p\}$.
Euler's formula: $|E| - |V| - |F| = -2$ (Proof)
- 2 For the 4-coloring of planar graphs, a graph that contains a C_i can be reduced to a smaller graph.
- 3 Thus every planar graph is 4-colorable.

Discharging, the outline


- 1 Every planar graph contains some $C \in \{C_1, C_2, \dots, C_p\}$.
Euler's formula: $2|E| - 6|V| + 4|E| - 6|F| < 0$
- 2 For the 4-coloring of planar graphs, a graph that contains a C_i can be reduced to a smaller graph.
- 3 Thus every planar graph is 4-colorable.

Discharging, the outline

- 1 Every planar graph contains some $C \in \{C_1, C_2, \dots, C_p\}$.
Euler's formula: $\sum_v (d(v) - 6) + \sum_f (2d(f) - 6) < 0$
- 2 For the 4-coloring of planar graphs, a graph that contains a C_i can be reduced to a smaller graph.
- 3 Thus every planar graph is 4-colorable.

Discharging, the outline


- 1 Every planar graph contains some $C \in \{C_1, C_2, \dots, C_p\}$.
Euler's formula: $\sum_v (d(v) - 6) + \sum_f (2d(f) - 6) < 0$



- 2 For the 4-coloring of planar graphs, a graph that contains a C_i can be reduced to a smaller graph.
- 3 Thus every planar graph is 4-colorable.

Discharging, the outline

- 1 Every planar graph contains some $C \in \{C_1, C_2, \dots, C_p\}$.
Euler's formula: $\sum_v (d(v) - 6) + \sum_f (2d(f) - 6) < 0$




$\Rightarrow \forall v, f, w'(v) \geq 0, w'(f) \geq 0$ **Contradiction!**

- 2 For the 4-coloring of planar graphs, a graph that contains a C_i can be reduced to a smaller graph.
- 3 Thus every planar graph is 4-colorable.

Discharging, the outline


- 1 Every planar graph contains some $C \in \{C_1, C_2, \dots, C_p\}$.
Euler's formula: $\sum_v (d(v) - 6) + \sum_f (2d(f) - 6) < 0$



$\Rightarrow \forall v, f, w'(v) \geq 0, w'(f) \geq 0$ **Contradiction!**


- 2 For the 4-coloring of planar graphs, a graph that contains a C_i can be reduced to a smaller graph.
- 3 Thus every planar graph is 4-colorable.

(Proof)




χ' : Minimum number of colors to ensure that


$$\text{○} \xrightarrow{a} \text{○} \xrightarrow{b} \text{○} \Rightarrow a \neq b.$$



χ' : Minimum number of colors to ensure that




χ'_e : Minimum size of every $L(e)$ such that




Edge Coloring (2)

Vertex coloring of **Line graphs**.




Edge Coloring (2)

Vertex coloring of **Line graphs**.




Edge Coloring (2)

Vertex coloring of **Line graphs**.




Edge Coloring (2)

Vertex coloring of **Line graphs**.



Edge Coloring (2)

Vertex coloring of **Line graphs**.




Conjecture (List Coloring Conjecture '85)

For any graph, $\chi'_e = \chi'$.

Edge Coloring (2)

Vertex coloring of **Line graphs**.



Conjecture (List Coloring Conjecture '85)


For any graph, $\chi'_e = \chi'$.

Theorem (Kahn '93)

$\chi'_e - \chi' = o(\chi')$ when $\chi' \rightarrow \infty$.

Edge Coloring (2)

Vertex coloring of **Line graphs**.



Conjecture (List Coloring Conjecture '85)

For any graph, $\chi'_e = \chi'$.

Theorem (Kahn '93)

$\chi'_e - \chi' = o(\chi')$ when $\chi' \rightarrow \infty$.

Theorem (Vizing '64)

For any graph, $\chi' \leq \Delta + 1$.

On planar graphs:

Theorem (Borodin '90)

For any *planar* graph with $\Delta \geq 9$, $x'_e \leq \Delta + 1$.

On planar graphs:

Theorem (Borodin '90)

For any *planar* graph with $\Delta \geq 9$, $\chi'_e \leq \Delta + 1$.

Simplified in 2008 by Cohen and Havet. (Proof)

On planar graphs:

Theorem (Borodin '90)

For any *planar* graph with $\Delta \geq 9$, $\chi'_e \leq \Delta + 1$.

Simplified in 2008 by Cohen and Havet. (Proof)

Theorem (B. '13+)

For any *planar* graph with $\Delta \geq 8$, $\chi'_e \leq \Delta + 1$.

On planar graphs:

Theorem (Borodin '90)

For any *planar* graph with $\Delta \geq 9$, $\chi'_e \leq \Delta + 1$.

Simplified in 2008 by Cohen and Havet. (Proof)

Theorem (B. '13+)

For any *planar* graph with $\Delta \geq 8$, $\chi'_e \leq \Delta + 1$.

(Should be true for every single graph)

Key lemma to the previous proof (and many other):

Lemma

Even cycles are 2-choosable.

Key lemma to the previous proof (and many other):

Lemma

Even cycles are 2-choosable.

Theorem (Borodin, Kostochka, Woodall '97)

For any bipartite multigraph G , if L is an edge list assignment such that $|L(u, v)| \geq d(u), d(v)$ for every (u, v) , then G is L -edge-choosable.

More Global discharging

Key lemma to the previous proof (and many other):

Lemma

Even cycles are 2-choosable.

Theorem (Borodin, Kostochka, Woodall '97)

For any bipartite multigraph G , if L is an edge list assignment such that $|L(u, v)| \geq d(u), d(v)$ for every (u, v) , then G is L -edge-choosable.

Theorem (Borodin, Kostochka, Woodall '97)

For any graph, $\Delta > mad^2 \Rightarrow \chi'_e = \Delta$.

Can also be used in many similar settings.

Who's afraid of discharging?

- It's fun to do (not so much to read...)

Who's afraid of discharging?

- It's fun to do (not so much to read...)
- and can be very elegant (I said "can").

Who's afraid of discharging?

- It's fun to do (not so much to read...)
- and can be very elegant (I said "can").
- It typically yields a polynomial algorithm (usually linear or quadratic).

Who's afraid of discharging?

- It's fun to do (not so much to read...)
- and can be very elegant (I said "can").
- It typically yields a polynomial algorithm (usually linear or quadratic).
- You can make a linear program out of it.

Who's afraid of discharging?

- It's fun to do (not so much to read...)
- and can be very elegant (I said "can").
- It typically yields a polynomial algorithm (usually linear or quadratic).
- You can make a linear program out of it.
- There are some hopes of partial automatization.

Who's afraid of discharging?

- It's fun to do (not so much to read...)
- and can be very elegant (I said "can").
- It typically yields a polynomial algorithm (usually linear or quadratic).
- You can make a linear program out of it.
- There are some hopes of partial automatization.
- It's extremely powerful in some settings (planar graphs, bounded density graphs).

Who's afraid of discharging?

- It's fun to do (not so much to read...)
- and can be very elegant (I said "can").
- It typically yields a polynomial algorithm (usually linear or quadratic).
- You can make a linear program out of it.
- There are some hopes of partial automatization.
- It's extremely powerful in some settings (planar graphs, bounded density graphs).
- ...Completely useless in others (?).

Who's afraid of discharging?

- It's fun to do (not so much to read...)
- and can be very elegant (I said "can").
- It typically yields a polynomial algorithm (usually linear or quadratic).
- You can make a linear program out of it.
- There are some hopes of partial automatization.
- It's extremely powerful in some settings (planar graphs, bounded density graphs).
- ...Completely useless in others (?).
- But we should keep in mind it's just a counting argument (the world is not divided into discharging and non-discharging proofs).

Some directions though

- FPT meets discharging (but not enough).

Some directions though

- FPT meets discharging (but not enough).
- Examples of global discharging on planar graphs that cannot be made local?

Some directions though

- FPT meets discharging (but not enough).
- Examples of global discharging on planar graphs that cannot be made local?
- How to automatize?

- FPT meets discharging (but not enough).
- Examples of global discharging on planar graphs that cannot be made local?
- How to automatize?

Thanks!