

From Antimagic to Equitable Labellings


Julien Bensmail^a, Clara Marcille^b

a: I3S/INRIA, Université Côte d'Azur, France

b: LaBRI, Université de Bordeaux, France

Journée du Département, April 17, 2024

Magic labellings




Magic labellings

2	7	6
9	5	1
4	3	8


Magic labellings

2	7	6
9	5	1
4	3	8




Magic labellings

2	7	6
9	5	1
4	3	8




Magic labellings

2	7	6
9	5	1
4	3	8




Magic labellings

2	7	6
9	5	1
4	3	8





Magic labellings

2	7	6
9	5	1
4	3	8




Antimagic labellings




Antimagic labellings

9	7	6
2	5	1
4	8	3




Antimagic labellings

9	7	6
2	5	1
4	8	3



Antimagic labellings

9	7	6
2	5	1
4	8	3




Definition

A *labelling* is said to be (locally) *distinguishing* if any pair of (adjacent) vertices have different *resulting sum*.

Antimagic labellings

9	7	6
2	5	1
4	8	3



Definition

A *labelling* is said to be (locally) *distinguishing* if any pair of (adjacent) vertices have different *resulting sum*.



We will get back to all this terminology later.

Regular Graphs

Definition

A graph is *regular* if all its vertices have the same number of incident edges, called *degree*.

There are a lot of different regular graphs, for example:




Question

What is an *irregular* graph?

Irregular graphs

Observation

There are no simple graphs where every two vertices have different degree.




Local definition

A graph is *locally irregular* if every two adjacent vertices have different degree.

Irregularity Strength

Edges with multiplicity


We now allow two vertices to be joined by multiple edges.



Irregularity Strength

Edges with multiplicity

We now allow two vertices to be joined by multiple edges.



Irregularity Strength

Edges with multiplicity

We now allow two vertices to be joined by multiple edges.



Irregularity Strength

Edges with multiplicity

We now allow two vertices to be joined by multiple edges.



Irregularity Strength

Edges with multiplicity

We now allow two vertices to be joined by multiple edges.



Irregularity Strength

Edges with multiplicity

We now allow two vertices to be joined by multiple edges.



Definition

We call *irregularity strength* of a graph G the smallest multiplicity needed to make G irregular (every two vertices have distinct degree).

Labellings

Definition

A k -labelling of a graph G is a function $\ell : E(G) \rightarrow \{1, \dots, k\}$.

Labellings

Definition

A k -labelling of a graph G is a function $\ell : E(G) \rightarrow \{1, \dots, k\}$.

Definition

We call *resulting sum* (relative to a labelling ℓ) of a vertex u the sum of labels on edges incident to u . We denote it $\sigma_\ell(u)$.

Labellings

Definition

A k -labelling of a graph G is a function $\ell : E(G) \rightarrow \{1, \dots, k\}$.

Definition

We call *resulting sum* (relative to a labelling ℓ) of a vertex u the sum of labels on edges incident to u . We denote it $\sigma_\ell(u)$.

Definition (Irregularity Strength)

We say a labelling ℓ is *distinguishing* if for every two vertices u and v of G , $\sigma_\ell(u) \neq \sigma_\ell(v)$.

Labellings

Definition

A k -labelling of a graph G is a function $\ell : E(G) \rightarrow \{1, \dots, k\}$.


Definition

We call *resulting sum* (relative to a labelling ℓ) of a vertex u the sum of labels on edges incident to u . We denote it $\sigma_\ell(u)$.


Definition (Local)

We say a labelling ℓ is *distinguishing* if for every two adjacent vertices u and v of G , $\sigma_\ell(u) \neq \sigma_\ell(v)$.


Small example




Small example




Small example




Small example




Small example




Small example




Small example




Small example




Small example




Small example




Small example




Small example



Small example



Small example



Conjectures and knowledge

These are the main conjectures and theorems in the field:

1-2-3 Conjecture (Karoński *et al.*, 2004)

All graphs admit a distinguishing 3-labelling.

Conjectures and knowledge

These are the main conjectures and theorems in the field:

1-2-3 Conjecture (Karoński *et al.*, 2004)

All graphs admit a distinguishing 3-labelling.

1-2-3-4-5 Theorem (Kalkowski *et al.*, 2010)

All graphs admit a (distinguishing) 5-labelling.

Conjectures and knowledge

These are the main conjectures and theorems in the field:

1-2-3 Conjecture (Karoński *et al.*, 2004)

All graphs admit a distinguishing 3-labelling.

1-2-3 Theorem (Keusch, 2024)

All graphs admit a (distinguishing) 3-labelling.

Conjectures and knowledge

These are the main conjectures and theorems in the field:

1-2-3 Conjecture (Karoński *et al.*, 2004)

All graphs admit a distinguishing 3-labelling.

1-2-3 Theorem (Keusch, 2024)

All graphs admit a (distinguishing) 3-labelling.

1-2 Problem (Ahadi *et al.*, 2013)

Given a graph G , deciding if G admits a 2-labellings is NP-complete.

Equitable Labellings

Definition

A k -labelling is said to be *equitable* if for all $1 \leq i, j \leq k$, the number of edges with label i and the number of edges with label j differs by at most 1.

Equitable Labellings

Definition

A k -labelling is said to be *equitable* if for all $1 \leq i, j \leq k$, the number of edges with label i and the number of edges with label j differs by at most 1.

In particular, a locally Antimagic labelling can be defined as a special case of equitable $|E(G)|$ -labelling!

Equitable Labellings

Definition

A k -labelling is said to be *equitable* if for all $1 \leq i, j \leq k$, the number of edges with label i and the number of edges with label j differs by at most 1.

In particular, a locally Antimagic labelling can be defined as a special case of equitable $|E(G)|$ -labelling!

Definition

We denote $\overline{\chi}_{\Sigma}(G)$ the smallest integer k such that G admits an equitable k -labelling.

Equitable Labellings

Definition

A k -labelling is said to be *equitable* if for all $1 \leq i, j \leq k$, the number of edges with label i and the number of edges with label j differs by at most 1.

In particular, a locally Antimagic labelling can be defined as a special case of equitable $|E(G)|$ -labelling!

Definition


We denote $\overline{\chi}_{\Sigma}(G)$ the smallest integer k such that G admits an equitable k -labelling.

Theorem (consequence of Lyngsie and Zhong, 2018)

If G is a nice graph, then $\overline{\chi}_{\Sigma}(G) \leq |E(G)|$.


Examples

An equitable labelling which happens to be antimagic:




Examples

An equitable labelling which happens to be antimagic:



An equitable 3-labelling:



Conjecture and contribution

Theorem (consequence of Lyngsie and Zhong, 2018)

If G is a nice graph, then $\overline{\chi}_{\Sigma}(G) \leq |E(G)|$.

Conjecture and contribution

Theorem (consequence of Lyngsie and Zhong, 2018)

If G is a nice graph, then $\overline{\chi}_{\Sigma}(G) \leq |E(G)|$.

Conjecture (Bensmail, Fioravantes, Mc Inerney, Nisse, 2021)

If G is a nice graph different from K_4 , then $\overline{\chi}_{\Sigma}(G) \leq 3$.

Conjecture and contribution

Theorem (consequence of Lyngsie and Zhong, 2018)

If G is a nice graph, then $\overline{\chi_{\Sigma}}(G) \leq |E(G)|$.

Conjecture (Bensmail, Fioravantes, Mc Inerney, Nisse, 2021)

If G is a nice graph different from K_4 , then $\overline{\chi_{\Sigma}}(G) \leq 3$.

Theorem (Bensmail, M., 2024+)


If G is a nice graph, then $\overline{\chi_{\Sigma}}(G) \leq \left\lfloor \frac{|E(G)|}{2} \right\rfloor + 2$.

Preliminary work

We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k + 1, k + 1, k + 2, k + 2)$ (where $k = \lfloor \frac{|E(G)|}{2} \rfloor$).


Preliminary work

We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \lfloor \frac{|E(G)|}{2} \rfloor$).



Preliminary work


We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \lfloor \frac{|E(G)|}{2} \rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 4, 5, 5, 6, 6)$.

Preliminary work


We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \lfloor \frac{|E(G)|}{2} \rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 4, 5, 5, 6, 6)$.

Preliminary work


We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k + 1, k + 1, k + 2, k + 2)$ (where $k = \lfloor \frac{|E(G)|}{2} \rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 4, 5, 5, 6, 6)$.

Preliminary work


We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \lfloor \frac{|E(G)|}{2} \rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 4, 5, 5, 6, 6)$.

Preliminary work


We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k + 1, k + 1, k + 2, k + 2)$ (where $k = \lfloor \frac{|E(G)|}{2} \rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 5, 5, 6, 6)$.

Preliminary work

We consider the *sequence of labels* $L = (1, 1, 2, 2, \dots, k+1, k+1, k+2, k+2)$ (where $k = \lfloor \frac{|E(G)|}{2} \rfloor$).



Here, $L = (1, 1, 2, 2, 3, 3, 4, 5, 5, 6, 6)$.

Question

How do we carry the fact that some edges received a weight?

Weight function


Solution

We will keep the information by updating a *weight function*.

Weight function

Solution


We will keep the information by updating a *weight function*.



Weight function

Solution


We will keep the information by updating a *weight function*.



Weight function

Solution

We will keep the information by updating a *weight function*.



We now consider the weighted graph (G, c) where c is the weight of a vertex of G .

Idea of the proof

- **Proceed by induction on the number of vertices.**

We consider a vertex u and assign the labels of L to the edges incident to u progressively to build a labelling ℓ , then remove u and the used labels.

Idea of the proof

- **Proceed by induction on the number of vertices.**

We consider a vertex u and assign the labels of L to the edges incident to u progressively to build a labelling ℓ , then remove u and the used labels.

- **Build a partial labelling of the graph, and extend it.**

We carry the contribution of the labelled edges through a weight function on the vertices adjacent to u .

Idea of the proof

- **Proceed by induction on the number of vertices.**

We consider a vertex u and assign the labels of L to the edges incident to u progressively to build a labelling ℓ , then remove u and the used labels.

- **Build a partial labelling of the graph, and extend it.**

We carry the contribution of the labelled edges through a weight function on the vertices adjacent to u .

Problem

Once all the edges incident to u have been labelled, there is no way to change $\sigma_\ell(u)$.

Idea of the proof

- **Proceed by induction on the number of vertices.**

We consider a vertex u and assign the labels of L to the edges incident to u progressively to build a labelling ℓ , then remove u and the used labels.

- **Build a partial labelling of the graph, and extend it.**

We carry the contribution of the labelled edges through a weight function on the vertices adjacent to u .

- **Ensure that a vertex will have a resulting sum smaller than the vertex treated later in the induction.**

We find the vertex of lowest *potential* resulting sum.

Idea of the proof

- **Proceed by induction on the number of vertices.**

We consider a vertex u and assign the labels of L to the edges incident to u progressively to build a labelling ℓ , then remove u and the used labels.

- **Build a partial labelling of the graph, and extend it.**

We carry the contribution of the labelled edges through a weight function on the vertices adjacent to u .

- **Ensure that a vertex will have a resulting sum smaller than the vertex treated later in the induction.**

We find the vertex of lowest *potential* resulting sum.

Problem

Maybe $G - u$ has a component isomorphic to K_2 .

Idea of the proof

- **Proceed by induction on the number of vertices.**

We consider a vertex u and assign the labels of L to the edges incident to u progressively to build a labelling ℓ , then remove u and the used labels.

- **Build a partial labelling of the graph, and extend it.**

We carry the contribution of the labelled edges through a weight function on the vertices adjacent to u .

- **Ensure that a vertex will have a resulting sum smaller than the vertex treated later in the induction.**


We find the vertex of lowest *potential* resulting sum.

- **Handle exceptions on the way.**

Identify and prevent problematic cases from arising while labelling.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



Here, the minimum possible resulting sum for this vertex is $1 + 6 + 7 = 14$. For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



Here, the minimum possible resulting sum for this vertex is $3 + 6 = 9$.
 For instance, consider $L = (6, 7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.

Vertex of lowest potential


We want to find the vertex with the lowest potential resulting sum:



For instance, consider $L = (7, 7, 8, 8, 9, 9)$. Each vertex is annotated with its current weight.


Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



Handling exceptions


Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



We choose a vertex of highest degree amongst vertices of lowest potential.

Handling exceptions


Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



We choose a vertex of highest degree amongst vertices of lowest potential.

Handling exceptions


Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



How to assign each label?

Handling exceptions


Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



How to assign each label?

Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Problem

A component isomorphic to K_2 with constant weight cannot be labelled.


Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Handling exceptions

Consider this weighted graph with $L = (2, 3, 3, 4, 4)$:



Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Problem

Two adjacent vertices of lowest potential can be in conflict.

Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:




Problem

Two adjacent vertices of lowest potential can be in conflict.


Handling exceptions

Consider this weighted graph with $L = (1, 1, 2, 2, 3, 3, 4, 4)$:



Handling exceptions

Consider this weighted graph with $L = (2, 3, 3, 4, 4)$:




The extra labels

They are some configuration where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$


The extra labels

They are some configuration where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$




The extra labels

They are some configuration where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$





The extra labels

They are some configuration where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$




The extra labels

They are some configuration where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$





The extra labels

They are some configuration where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$





The extra labels

They are some configuration where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$





The extra labels

They are some configuration where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$





The extra labels

They are some configuration where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$



The extra labels

They are some configuration where you can not use the smallest labels. Assume for instance the sequence of labels is $(1, 1, 2, 2, 3, 3)$



Conclusion

Conclusion

- We essentially improved the bound on equitable labellings by a factor of 2.

Conclusion

- We essentially improved the bound on equitable labellings by a factor of 2.
- The question of a constant bound remains completely open.

Conclusion

- We essentially improved the bound on equitable labellings by a factor of 2.
- The question of a constant bound remains completely open.
- The tools we used do not generalize nicely to multiplicity higher than 2.

Conclusion

- We essentially improved the bound on equitable labellings by a factor of 2.
- The question of a constant bound remains completely open.
- The tools we used do not generalize nicely to multiplicity higher than 2.
- More work can be done to characterize the graphs that cannot be done with a sequence of $|E(G) + 1|$ labels and multiplicity 2.

Conclusion

- We essentially improved the bound on equitable labellings by a factor of 2.
- The question of a constant bound remains completely open.
- The tools we used do not generalize nicely to multiplicity higher than 2.
- More work can be done to characterize the graphs that cannot be done with a sequence of $|E(G) + 1|$ labels and multiplicity 2.

Thank you for your attention!