

[GTN LXV:1]

## BOUNDS FOR THE $(m, n)$ -MIXED CHROMATIC NUMBER AND THE ORIENTED CHROMATIC NUMBER

William F. Klostermeyer<sup>1</sup>, Gary MacGillivray<sup>2</sup>,  
André Raspaud<sup>3</sup>, and Eric Sopena<sup>3</sup>

<sup>1</sup>School of Computing  
University of North Florida  
Jacksonville, Florida 32224-2669, U.S.A.  
<klostermeyer@hotmail.com>

<sup>2</sup>Department of Mathematics and Statistics  
University of Victoria  
Victoria BC, CANADA V8W 3R4

<sup>3</sup>LaBRI, UMR CNRS 5800  
Universite Bordeaux I,  
33405 Talence Cedex, FRANCE

### 1. Introduction

A bound for the  $(n, m)$ -mixed chromatic number in terms of the chromatic number of the square of the underlying undirected graph is obtained. A similar bound holds when the chromatic number of the square is replaced by the *injective chromatic number*. When restricted to  $n = 1$  and  $m = 0$  (that is, *oriented graphs*) this provides a new bound for the *oriented chromatic number*. In this case, a slightly improved bound is obtained if the chromatic number of the square is replaced the 2-dipath chromatic number (defined in Section 4). In all cases, the method of proof generalizes an argument that has been used to obtain Brooks-type theorems for injective oriented colorings [1]–[8]. Similar, although not identical, arguments have appeared in the work of Sopena [9], and Nešetřil and Raspaud [10][11].

Colorings of mixed graphs were first studied by Nešetřil and Raspaud [11]. They gave a bound on the  $(n, m)$ -mixed chromatic number in terms of the *acyclic chromatic number* of the underlying undirected graph. Bounds for the  $(0, 2)$ -mixed chromatic number of planar graphs and other graph families have been obtained [12]. Some of these have been extended to arbitrary  $n$  and  $m$  [13].

### 2. Definitions

For basic definitions in graph theory, see the text by Bondy and Murty [14].

**Definition 2.1:** A  $(n, m)$ -colored mixed graph is an  $(m + n + 1)$ -tuple

$$G = (V, A_1, A_2, \dots, A_n, E_1, E_2, \dots, E_m),$$

where:

- (1)  $V$  is a set of vertices;
- (2) for  $i = 1, 2, \dots, n$ ,  $A_i$  is a set of ordered pairs of distinct vertices of  $G$  called the arcs of color  $i$ ;
- (3) for  $j = 1, 2, \dots, m$ ,  $E_j$  is a set of unordered pairs of distinct vertices of  $G$  called the edges of color  $j$ ; and
- (4) the underlying undirected graph  $U[G]$ , with vertex set  $V(U[G]) = V(G)$  and an edge joining  $x$  to  $y$  for every  $i$  for which the ordered pair  $xy \in A_i$  and for every  $j$  for which the unordered pair  $xy \in E_j$ , is simple. ■

Observe that a  $(0, 1)$ -colored mixed graph is a simple graph, and a  $(1, 0)$ -colored mixed graph is an oriented graph.

Homomorphisms between  $(n, m)$ -colored mixed graphs are defined in the natural way. For  $(n, m)$ -colored mixed graphs  $G$  and  $H$ , a *homomorphism* of  $G$  to  $H$  is a function  $f: V(G) \rightarrow V(H)$  such that for  $1 \leq i \leq n$ ,

$f(x)f(y) \in A_i(H)$  whenever  $xy \in A_i(G)$ , and for  $1 \leq j \leq m$ ,  $f(x)f(y) \in E_j(H)$  whenever  $xy \in E_j(G)$ . For more on homomorphisms, see [15].

Observe that homomorphisms of  $(n, m)$ -colored mixed graphs compose. That is, if  $F$ ,  $G$ , and  $H$  are  $(n, m)$ -colored mixed graphs,  $f$  is a homomorphism of  $F$  to  $G$ , and  $g$  is a homomorphism of  $G$  to  $H$ , then  $g \circ f$  is a homomorphism of  $F$  to  $H$ .

By analogy with the corresponding definitions for other coloring concepts for oriented graphs, a  $k$ -coloring of a  $(n, m)$ -colored mixed graph is defined to be a homomorphism to a  $(n, m)$ -colored mixed graph on  $k$  vertices. The smallest  $k$  for which there is a  $k$ -coloring of a  $(n, m)$ -colored graph  $G$  is called the  $(n, m)$ -mixed chromatic number of  $G$  and is denoted by  $\chi_{(n, m)}(G)$ .

Informally, a  $k$ -coloring of an  $(n, m)$ -colored mixed graph is a partition of the vertices of  $G$  into  $k$  independent sets (sets containing no edges or arcs of any color),  $X_1, X_2, \dots, X_k$ , such that, for any two independent sets  $X_p$  and  $X_q$ , there is only one type of adjacency; that is, only arcs of the same color and orientation; or only edges of the same color, between vertices in  $X_p$  and  $X_q$ .

**Example 1:** Consider a path on four vertices  $v_1, v_2, v_3, v_4$  with arcs  $v_1v_2$  and  $v_2v_3$  of color 1 and an edge  $v_3v_4$  of color 1. Call this  $(1, 1)$ -colored mixed graph  $G$ . Then  $\chi_{(n, m)}(G) = 3$ ; color  $v_1$  and  $v_4$  with color 1,  $v_2$  with color 2, and  $v_3$  with color 3.

**Example 2:** Consider a path on five vertices  $v_1, v_2, v_3, v_4, v_5$  with arcs  $v_1v_2$  and  $v_2v_3$  of color 1, an edge  $v_3v_4$  of color 1, and an edge  $v_4v_5$  of color 2. Call this  $(1, 2)$ -colored mixed graph  $G$ . Then  $\chi_{(n, m)}(G) = 4$ ; color  $v_1$  and  $v_4$  with color 1,  $v_2$  with color 2,  $v_3$  with color 3, and  $v_5$  with color 4.

We next define a  $(n, m)$ -colored mixed graph, Let  $t \geq 3$ . For  $k = 1, 2, \dots, t$ , let  $I_k$  be the set of all sequences of length  $t$  in which the  $k^{\text{th}}$  element is “ $\cdot$ ” and every other entry is a “ $+_i$ ” or “ $-_i$ ” ( $1 \leq i \leq n$ ), or “ $\sim_j$ ” ( $1 \leq j \leq m$ ).

**Definition 2.2:** The graph  $H_{(n, m)}^t$  is the  $(n, m)$ -colored mixed graph with vertex set

$$V(H_{(n, m)}^t) = I_1 \cup I_2 \cup \dots \cup I_t \text{ and}$$

- (1) an arc of color  $i$  joining sequence  $\alpha \in I_k$  to sequence  $\beta \in I_l$  if and only if the  $l^{\text{th}}$  entry of  $\alpha$  is  $+_i$  and the  $k^{\text{th}}$  entry of  $\beta$  is  $-_i$ , and
- (2) an edge of color  $j$  joining sequence  $\alpha \in I_k$  to sequence  $\beta \in I_l$  if and only if the  $l^{\text{th}}$  entry of  $\alpha$  and the  $k^{\text{th}}$  entry of  $\beta$  are both  $\sim_j$ . ■

In the next section, we bound the  $(n, m)$ -mixed chromatic number of  $H_{(n, m)}^t$ .

### 3. Main Result

The graph  $H_{(1, 0)}^t$  is used to prove bounds for injective oriented colorings [1]–[5]. Young proved that an oriented graph can be properly  $t$ -colored so that any two vertices joined by a directed path of length two get different colors if and only if it has an homomorphism to  $H_{(1, 0)}^t$  [8] (Min and Wang [16] call this a  $2$ -dipath  $k$ -coloring.) Our bound on the  $(n, m)$ -mixed chromatic number of a  $(n, m)$ -mixed colored graph  $G$  will be obtained by coloring the square of the underlying graph of  $G$  with  $t$  colors, and then refining this coloring to obtain a homomorphism of  $G$  to  $H_{(n, m)}^t$ .

**Lemma 3.1 (The Refinement Lemma):**

$$\begin{aligned} \chi_{(n, m)}(H_{(n, m)}^t) &\leq 1 + (2n + m) + (2n + m)^2 + \dots + (2n + m)^{t-1} \\ &= \begin{cases} t & \text{if } n = 0 \text{ and } m = 1 \\ \frac{(2n + m)^t - 1}{2n + m - 1} & \text{otherwise.} \end{cases} \end{aligned}$$

**Proof:** Let  $B = 1 + (2n + m) + (2n + m)^2 + \dots + (2n + m)^{t-1}$ . We describe a  $(n, m)$ -mixed  $B$ -coloring of  $H_{(n, m)}^t$ . For  $k = 1, 2, \dots, t$ , partition the independent set  $I_k$  into  $(2n + m)^{k-1}$  subsets  $I_{k, l}$ ,  $1 \leq l \leq (2n + m)^{k-1}$ , such that all sequences in  $I_{k, l}$  agree in the first  $k$  places. The total number of sets  $I_{p, q}$  is  $B$ .

By definition of these sets, if  $p < r$  then:

- (1) all arcs joining vertices in  $I_{p,q}$  and  $I_{r,s}$  are of color  $i$  and
  - (a) are oriented toward  $I_{p,q}$  if the  $p^{\text{th}}$  symbol of all sequences in  $I_{r,s}$  is  $+_i$  and the  $r^{\text{th}}$  symbol in all sequences in  $I_{r,s}$  is  $-_i$ ;
  - (b) oriented towards  $I_{r,s}$  if the  $p^{\text{th}}$  symbol of all sequences in  $I_{r,s}$  is  $-_i$  and the  $r^{\text{th}}$  symbol in all sequences in  $I_{r,s}$  is  $+_i$ ;
- (2) all edges joining vertices in  $I_{p,q}$  and  $I_{r,s}$  are of color  $j$  if the  $p^{\text{th}}$  symbol of all sequences in  $I_{r,s}$  and the  $r^{\text{th}}$  symbol of all sequences in  $I_{p,q}$  are both  $\sim_j$ .

Hence, by identifying all vertices in each set  $I_{k,l}$ ,  $1 \leq k \leq t$ ,  $1 \leq l \leq (2n + m)^{k-1}$ , a homomorphism of  $H_{(n,m)}^t$  onto the  $(n,m)$ -colored mixed graph  $O_{(n,m)}$  on  $B$  vertices is obtained. Therefore,  $\chi_{(n,m)}(H_{(n,m)}^t) \leq B$ . ■

Lemma 3.1 is dubbed the *Refinement Lemma* because the independent sets  $I_k$  are refined to obtain the homomorphism to  $O_{(n,m)}$ .

Let  $G$  be an  $(n,m)$ -colored mixed graph. Define  $S(G)$  to be square of the underlying undirected graph of  $G$ . That is,  $S(G)$  has the same vertex set as  $G$  and  $xy \in E(S(G))$  if and only if  $1 \leq \text{dist}_G(x, y) \leq 2$ .

**Lemma 3.2:** Let  $G$  be an  $(n,m)$ -colored mixed graph. If  $S(G)$  is  $t$ -colorable, then there is a homomorphism from  $G$  to  $H_{(n,m)}^t$ .

**Proof:** Let  $C_1, C_2, \dots, C_t$  be a  $t$ -coloring of  $S(G)$ . Let  $x \in C_k$ . Define the sets

$$S_{x,i} = \{p : xy \in A_i(G) \text{ for some } y \in C_p\}, R_{x,i} = \{p : yx \in A_i(G) \text{ for some } y \in C_p\}, \text{ and}$$

$$T_{x,j} = \{p : xy \in E_j(G) \text{ for some } y \in C_p\}.$$

These are, respectively, the sets of (vertex) colors to which  $x$  sends an arc of color  $i$  in  $D$ , from which  $x$  receives an arc of color  $i$  in  $D$ , or to which  $x$  is joined by an edge of color  $j$  in  $D$ . Since  $C_k$  is an independent set,  $x \notin S_{x,i} \cup R_{x,i'} \cup T_{x,j}$  for any  $i, i'$ , and  $j$ . By construction of  $S(G)$ , for all  $i, i'$ , and  $j$  the intersection between any two of  $S_{x,i}, R_{x,i'}$ , and  $T_{x,j}$  is empty.

Thus, each vertex in  $C_k$  can be associated with a sequence with  $k^{\text{th}}$  entry “.” and in which the  $l^{\text{th}}$  entry is “ $+_i$ ” if  $k \in S_{x,i}$ , is “ $-_i$ ” if  $k \in R_{x,i}$ , is “ $\sim_j$ ” if  $k \in T_{x,j}$ , and is “ $\sim_i$ ” otherwise. This is a homomorphism of  $G$  to  $H_{(n,m)}^t$ . ■

**Corollary 3.3:** Let  $G$  be an  $(n,m)$ -colored mixed graph. If  $S(G)$  is  $t$ -colorable, then

$$\chi_{(n,m)}(G) \leq 1 + (2n + m) + (2n + m)^2 + \dots + (2n + m)^{t-1}$$

$$= \begin{cases} t & \text{if } n = 0 \text{ and } m = 1 \\ \frac{(2n + m)^t - 1}{2n + m - 1} & \text{otherwise.} \end{cases}$$

An *injective  $k$ -coloring* of a graph  $G$  is an assignment of  $k$  colors to the vertices of  $G$  so that vertices at distance two are assigned different colors. Adjacent vertices may be assigned the same color. The *injective chromatic number* of  $G$  is the least  $k$  for which there exists an injective  $k$ -coloring of  $G$ . The injective chromatic number is bounded above by the chromatic number of  $S(G)$  (see, for example, [17]).

The previous arguments work essentially as given with a small modification to take into account the fact that vertices of the same color may be adjacent. The graph  $H_{(n,m)}^t$  is redefined by replacing each “.” by one of “ $+_i$ ”, “ $-_i$ ”, or “ $\sim_j$ ”. This increases both the number of vertices of  $H_{(n,m)}^t$  and the bound on its  $(n,m)$ -mixed chromatic number by a factor of  $(2n + m)$ . Thus, one obtains:

**Corollary 3.4:** Let  $G$  be an  $(n, m)$ -colored mixed graph. If  $G$  has an injective  $t$ -coloring, then

$$\begin{aligned} \chi_{(n, m)}(G) &\leq (2n + m) + (2n + m)^2 + \dots + (2n + m)^t \\ &= \begin{cases} t & \text{if } n = 0 \text{ and } m = 1 \\ (2n + m) \frac{(2n + m)^t - 1}{2n + m - 1} & \text{otherwise.} \end{cases} \end{aligned}$$

■

#### 4. Oriented Colorings

When  $n = 1$  and  $m = 0$  we have an *oriented graph*, and the  $(n, m)$ -mixed chromatic number is equal to the oriented chromatic number. Hence, the above results imply bounds for the oriented chromatic number. In particular, if  $S(G)$  has a  $t$ -coloring, then  $\chi_o(G) \leq 2^t - 1$ .

An important property of a coloring of  $S(G)$ , or an injective  $t$ -coloring of  $G$ , used in the above arguments is that vertices joined by a directed path of length two must be assigned different colors. The *2-dipath chromatic number* of  $G$  is the smallest  $k$  for which there is a 2-dipath  $k$ -coloring of  $G$ . The 2-dipath chromatic number is at most the chromatic number of the square [16].

The argument in Section 3 goes through unchanged for oriented graphs if a  $k$ -coloring of  $S(G)$  is replaced by a 2-dipath  $k$ -coloring of  $G$ . This is the same as replacing  $S(G)$  by the square of  $G$  as a directed graph (two vertices are adjacent if they are joined by a directed path of length at most two) in the argument. Doing so, one obtains the following bound:

**Corollary 4.1:** Let  $G$  be an oriented graph. If  $G$  has a 2-dipath  $t$ -coloring, then

$$\chi_o(G) = \chi_{(1, 0)}(G) \leq 2^t - 1.$$

■

Finally, if the 2-dipath  $t$ -coloring need not assign different colors to adjacent vertices, then proceeding as discussed at the end of the previous section one obtains  $\chi_o(G) = \chi_{(1, 0)}(G) \leq 2^{t+1} - 1$  (see also [7]).

#### References

- [1] R. Campbell; *Oriented Colorings that are Injective on Neighbourhoods*. M.Sc. Thesis, Department of Mathematics and Statistics, University of Victoria, Canada (2009).
- [2] N. Clarke, and G. MacGillivray; Injective oriented colourings: In- and out-neighbourhoods separately. Manuscript (2009).
- [3] G. MacGillivray, A. Raspaud, and J. Swarts; *Injective Oriented Colourings*. Lecture Notes in Computer Science 5911, Montpellier, France. Proceedings of WG (2009).
- [4] G. MacGillivray, A. Raspaud, and J. Swarts; Obstructions to injective oriented colouring. *Electronic Notes in Discrete Math.*, **38**, 597–605 (2011).
- [5] G. MacGillivray, A. Raspaud, and J. Swarts; Obstructions to locally injective oriented improper colorings. Manuscript (2011).
- [6] G. MacGillivray and J. Swarts; The complexity of locally injective homomorphisms. *Discrete Mathematics*, **310**, 2685–2696 (2010).
- [7] J. Swarts; *The Complexity of Digraph Homomorphisms: Local Tournaments, Injective Homomorphisms and Polymorphisms*. Ph.D. Thesis, Department of Mathematics and Statistics, University of Victoria, Victoria, BC, Canada (2008).
- [8] K.M. Young; *2-Dipath and Proper 2-Dipath Colourings*. M.Sc. Thesis, University of Victoria (2011).
- [9] E. Sopena; The chromatic number of oriented graphs. *J. Graph Theory*, **25**, 191–205 (1997).
- [10] J. Nešetřil and A. Raspaud; Antisymmetric ows and strong colorings of oriented graphs. *Ann. Inst. Fourier, Grenoble*, **49**, 1037–1056 (1999).
- [11] J. Nešetřil and A. Raspaud; Coloured homomorphisms of colored mixed graphs. *J. Combinatorial Theory, Series B*, **80**, 147–155 (2000).
- [12] A. Montejano, P. Ochem, A. Pinlou, A. Raspaud, and E. Sopena; Homomorphisms of 2-edge-colored graphs. *Discrete Applied Mathematics*, **158**, 1365–1379 (2010).

- [13] A. Montejano, A. Pinlou, A. Raspaud, and E. Sopena; Chromatic number of sparse colored planar graphs. *Electronic Notes in Discrete Math.*, **34**, 363–367 (2009).
  - [14] J.A. Bondy and U. S. R.Murty; *Graph Theory*. Springer (2007).
  - [15] P. Hell and J. Nešetřil; *Graphs and Homomorphisms*. Oxford University Press (2004).
  - [16] C. Min and W. Wang, The 2-dipath chromatic number of Halin graphs. *Information Processing Letters*, **99**, 47–53 (2006).
  - [17] G. Hahn, J. Kratochvíl, J. Siran, and D. Sotteau; On the injective chromatic number of graphs. *Discrete Math.*, **256**, 179–192 (2002).
- 

Received: October 1, 2012  
Revised: February 16, 2013